

Producto: Prospecto/Leaflet
Codigo SAP/ SAP code:
Codigo anterior/ Previous code:
Formato/ format: 250 x 160 mm
Creado el/ Creation Date:
Gramaje/ Weight: 50g / m²

CONFORME
LABORATORIO:



Anverso:

160 mm

Package leaflet: Information for the user

Chloralieve™

Honey & Lemon Flavour

2mg / 0.6mg / 1.2mg lozenges

Lidocaine Hydrochloride / Amylmetacresol / 2,4-Dichlorobenzyl Alcohol

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 2 days.

1. What Chloralieve™ is and what it is used for

Chloralieve Honey & Lemon Flavour lozenges (referred to as Chloralieve in this leaflet) contains amylmetacresol and 2,4-dichlorobenzyl alcohol - both antiseptics, and lidocaine hydrochloride - a local anaesthetic for the throat. Chloralieve is indicated for the local relief of symptoms of sore throat in adults and children over 12 years of age.

You must talk to a doctor if you do not feel better or if you feel worse after 2 days.

2. What you need to know before you take Chloralieve™.

Do not take Chloralieve:

- if you are allergic to lidocaine hydrochloride or other amide-type local anaesthetic drugs, amylmetacresol, 2,4-dichlorobenzyl alcohol or any of the other ingredients of this medicine (listed in section 6).
- Children under 12 years old should not take this medicine.
- If you have a history of allergy to local anaesthetics such as lidocaine, bupivacaine, and ropivacaine.
- If you have a history of or are suspected to have a blood condition called methaemoglobinemia.

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Chloralieve. You should inform your doctor if:

- You suffer from asthma or narrowing of the airways (bronchospasm).
- Your symptoms do not improve after two days or if you have a fever, headaches, nausea, vomiting or skin rash.

This product may be harmful for the teeth.

- The anesthetic contained in this medicine may cause aspiration (coughing during meals or a choking sensation) while eating. Do not take food directly following the use of this medicine.
- This medicine may cause numbness of the tongue and may increase the danger of biting trauma. Therefore care should be taken in eating and drinking hot foods.
- Prolonged use of this medicine (more than 5 days) is not recommended, as it may alter the natural microbial balance of the throat.

What is in this leaflet

1. What Chloralieve is and what it is used for
2. What you need to know before you take Chloralieve
3. How to take Chloralieve
4. Possible side effects
5. How to store Chloralieve
6. Contents of the pack and other information

- Follow the indicated dosage: if taken in large quantities or over time this medicine may affect the heart or nervous system, and may cause convulsions (see section "If you take more Chloralieve than you should").
- Elderly persons or persons in a weakened condition are more sensitive to possible adverse reactions and should consult their doctor before taking this medicine.
- This medicine should not be used in the area of mouth and throat if greater acute wounds exist.

Children and adolescents

This medicine is not to be used in children under 12 years of age.

Other medicines and Chloralieve

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor or pharmacist if you are taking any of the following medicines, as it may be necessary to adjust their doses:

- Beta blockers (used to treat heart failure or arterial disease) or medicines containing cimetidine (used to treat stomach ulcers).
- Other local anaesthetics (amides).
- Medicines used to treat heart disorders, such as mexiletine or procainamide.
- Medicines such as fluvoxamine (used to treat depression).
- Antibiotics (drugs for certain bacterial infections), such as erythromycin or antimycotics (medicines against certain fungal infections), such as itraconazole.

Although no interactions should occur, do not take other mouth or throat antiseptics while using Chloralieve.

Chloralieve with food, drink and alcohol

Do not take this medicine before meals or before drinking.

Pregnancy, breast-feeding and fertility

The use of this medicine is not advised during pregnancy and lactation. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Driving and using machines

Observe your response to this medicine. Taken at the

NON PRINTABLE AREA

XXXXXXXXXX

PROCESS
BLACK

