

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Loperamide Hydrochloride Tablets 2mg

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 2 mg Loperamide hydrochloride.

Excipient with known effect: lactose

For a full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM

Tablet

White, round, biconvex tablet marked “aL”-on one side and plain on reverse.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

For the symptomatic treatment of acute diarrhoea, in adults and children aged 12 years and over.

For the symptomatic treatment of acute episodes of diarrhoea associated with irritable bowel syndrome in adults aged 18 years and over following initial diagnosis by a doctor.

4.2 Posology and method of administration

Posology:

Acute Diarrhoea

Adults and children over 12 years:

2 tablets (4 mg) initially followed by 1 tablet (2 mg) after every loose stool.

The maximum daily dose should not exceed 6 tablets (12 mg).

Symptomatic treatment of acute episodes of diarrhoea associated with irritable bowel syndrome in adults aged 18 years and over

Two tablets (4 mg) to be taken initially, followed by 1 tablet (2 mg) after every loose stool, or as previously advised by your doctor. The maximum daily dose should not exceed 6 tablets (12 mg).

Paediatric population

Contraindicated in children younger than 12 years of age

Use in elderly

No dose adjustment is required for the elderly.

Renal impairment

No dose adjustment is required for patients with renal impairment.

Hepatic impairment

Although no pharmacokinetic data are available in patients with hepatic impairment, Imodium should be used with caution in such patients because of reduced first pass metabolism. (See 4.4 Special warnings and special precautions for use).

Method of administration

Oral use. The tablets should be taken with liquid.

4.3 Contraindications

Loperamide HCl is contraindicated:

- in patients with known hypersensitivity to loperamide HCl or to any of the excipients,
- in children under 12 years of age,.

- in patients with acute dysentery, which is characterised by blood in stools and high fever,

- in patients with acute ulcerative colitis,

- in patients with bacterial enterocolitis caused by invasive organisms including Salmonella, Shigella, and Campylobacter,

- in patients with pseudomembranous colitis associated with the use of broad-spectrum antibiotics.

Loperamide HCl should not be used when inhibition of peristalsis is to be avoided due to the possible risk of significant sequelae including ileus, megacolon and toxic megacolon. Loperamide HCl must be discontinued promptly when constipation, abdominal distension or ileus develop

4.4 Special Warnings and Special Precautions for Use

Treatment of diarrhoea with loperamide HCl is only symptomatic. Whenever an underlying aetiology can be determined, specific treatment should be given when appropriate. In patients with diarrhoea, especially in children, frail and elderly patients, fluid and electrolyte depletion may occur. In such cases administration of appropriate fluid and electrolyte replacement therapy is the most important measure. Use of this medicine does not preclude the administration of appropriate fluid and electrolyte replacement therapy.

In acute diarrhoea, if clinical improvement is not observed within 48 hours, the administration of loperamide HCl should be discontinued and patients should be advised to consult their doctor.

Persistent diarrhoea can be an indicator of potentially more serious conditions, and this medicine should not be used for prolonged periods until the underlying cause of the diarrhoea has been investigated.

Patients with AIDS treated with loperamide HCl for diarrhoea should have therapy stopped at the earliest signs of abdominal distension. There have been isolated reports of obstipation with an increased risk for toxic megacolon in AIDS patients with infectious colitis from both viral and bacterial pathogens treated with loperamide HCl.

Although no pharmacokinetic data are available in patients with hepatic impairment, loperamide HCl should be used with caution in such patients because of reduced first pass metabolism, as it may result in a relative overdose leading to CNS toxicity.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine because it contains lactose.

If patients are taking this medicine to control episodes of diarrhoea associated with Irritable Bowel Syndrome previously diagnosed by their doctor, and clinical improvement is not observed within 48 hours, the administration of loperamide HCl should be discontinued and they should consult with their doctor. Patients should also return to their doctor if the pattern of their symptoms changes or if the repeated episodes of diarrhoea continue for more than two weeks.

Cardiac events including QT interval and QRS complex prolongation, torsade de pointes have been reported in association with overdose. Some cases had a fatal outcome (see section 4.9). Overdose can unmask existing Brugada syndrome. Patients should not exceed the recommended dose and/or the recommended duration of treatment

“Caution is needed in patients with a history of drug abuse. Loperamide is and opioid and addiction is observed with opioids as a class.”

For acute episodes of diarrhoea associated with irritable bowel syndrome
Special Warnings to be included in the leaflet:

Only take loperamide HCl to treat acute episodes of diarrhoea associated with irritable bowel syndrome (IBS) if your doctor has previously diagnosed IBS.

- If any of the following now apply, do not use the product without first consulting your doctor, even if you know you have IBS:
- If you are aged 40 years or over and it is some time since your last attack of IBS or the symptoms are different this time
- If you have recently passed blood from the bowel
- If you suffer from severe constipation
- If you are feeling sick or vomiting
- If you have lost your appetite or lost weight
- If you have difficulty or pain passing urine
- If you have a fever
- If you have recently travelled abroad

Consult your doctor if you develop new symptoms, or if your symptoms worsen, or if your symptoms have not improved over two weeks.

4.5 Interaction with other medicinal products and other forms of interaction

Non-clinical data have shown that loperamide is a P-glycoprotein substrate. Concomitant administration of loperamide (16 mg single dose) with quinidine, or ritonavir, which are both P-glycoprotein inhibitors, resulted in a 2 to 3-fold increase in loperamide plasma levels. The clinical relevance of this pharmacokinetic interaction with P-glycoprotein inhibitors, when loperamide is given at recommended doses, is unknown.

The concomitant administration of loperamide (4 mg single dose) and itraconazole, an inhibitor of CYP3A4 and P-glycoprotein, resulted in a 3 to 4-fold increase in loperamide plasma concentrations. In the same study a CYP2C8 inhibitor, gemfibrozil, increased loperamide by approximately 2-fold. The combination of itraconazole and gemfibrozil resulted in a 4-fold increase in peak plasma levels of loperamide and a 13-fold increase in total plasma exposure. These increases were not associated with central nervous system (CNS) effects as measured by psychomotor tests (i.e. subjective drowsiness and the Digit Symbol Substitution Test).

The concomitant administration of loperamide (16 mg single dose) and ketoconazole, an inhibitor of CYP3A4 and P-glycoprotein, resulted in a 5-fold increase in loperamide plasma concentrations. The increase was not associated with increased pharmacodynamic effects as measured by pupillometry.

Concomitant treatment with oral desmopressin resulted in a 2-fold increase of desmopressin plasma concentrations, presumably due to slower gastrointestinal motility.

It is expected that drugs with similar pharmacological properties may potentiate loperamide's effect and that drugs that accelerate gastrointestinal transit may decrease its effect

4.6 Fertility, pregnancy and lactation

Pregnancy

Safety in pregnancy has not been established, although studies in animals have not demonstrated any teratogenic or embryotoxic effects. As with other drugs, it is not advisable to administer Loperamide during pregnancy, especially during the first trimester.

Lactation

Small amounts of loperamide may appear in human breast milk. Therefore, this medicine is not recommended during breast-feeding.

Women who are pregnant or breast feeding infants should therefore be advised to consult their doctor for appropriate treatment.

Fertility

The effect on human fertility has not been evaluated.

4.7 Effects on ability to drive and use machines

Loss of consciousness, depressed level of consciousness, tiredness, dizziness or drowsiness may occur when diarrhoea is treated with loperamide HCl. Therefore, it is advisable to use caution when driving a car or operating machinery. See section 4.8 undesirable effects.

4.8 Undesirable effects

Adults and children aged ≥ 12 years

The safety of loperamide HCl was evaluated in 2755 adults and children aged ≥ 12 years who participated in 26 controlled and uncontrolled clinical trials of loperamide HCl used for the treatment of acute diarrhoea.

The most commonly reported (i.e. $\geq 1\%$ incidence) adverse drug reactions (ADRs) in clinical trials with loperamide HCl in acute diarrhoea were: constipation (2.7%), flatulence (1.7%), headache (1.2%) and nausea (1.1%).

Table 1 displays ADRs that have been reported with the use of loperamide HCl from either clinical trial (acute diarrhoea) or post-marketing experience.

The frequency categories use the following convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); and very rare ($< 1/10,000$).

Table 1: Adverse Drug Reactions

System Organ Class	Indication			
	Common	Uncommon	Rare	Unknown
Immune System Disorders			Hypersensitivity reaction ^a Anaphylactic	

			reaction (including Anaphylactic shock) ^a Anaphylactoid reaction ^a	
Nervous System Disorders	Headache	Dizziness Somnolence ^a	Loss of consciousness ^a Stupor ^a Depressed level of consciousness ^a Hypertonia ^a Coordination abnormality ^a	
Eye Disorders			Miosis ^a	
Gastrointestinal Disorders	Constipation Nausea Flatulence	Abdominal pain Abdominal discomfort Dry mouth Abdominal pain upper Vomiting Dyspepsia ^a	Ileus ^a (including paralytic ileus) Megacolon ^a (including toxic megacolon ^b) Abdominal distension	acute pancreatitis
Skin and Subcutaneous Tissue Disorders		Rash	Bullous eruption ^a (including Stevens-Johnson syndrome, Toxic epidermal necrolysis and Erythema multiforme) Angioedema ^a Urticaria ^a Pruritus ^a	
Renal and Urinary Disorders			Urinary retention ^a	
General Disorders and Administration Site Conditions			Fatigue ^a	

a) Inclusion of this term is based on post-marketing reports for loperamide HCl. As the process for determining post marketing ADRs did not differentiate between chronic and acute indications or adults and children, the frequency is estimated from all clinical trials with loperamide HCl (acute and chronic), including trials in children ≤ 12 years (N=3683).

b) See section 4.4 Special Warnings and Special Precautions for use.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at:

www.mhra.gov.uk/yellowcard.

The following adverse reaction(s) should be added under the SOC Gastrointestinal disorder with a frequency “not known”:
acute pancreatitis

4.9 Overdose

Symptoms:

In cases of overdose (including relative overdose due to hepatic dysfunction) the following effects may occur – constipation, ileus, urinary retention and neurological symptoms (CNS depression, stupor, coordination abnormality, somnolence, bradypnoea, miosis, muscular hypertonía and respiratory depression).

Children and patients with hepatic dysfunction may be more sensitive to CNS effects than adults.

In individuals who have ingested overdoses of loperamide HCl, cardiac events such as QT interval and QRS complex prolongation, torsade de pointes, other serious ventricular arrhythmias, cardiac arrest and syncope have been observed (see section 4.4). Fatal cases have also been reported. Overdose can unmask existing Brugada syndrome.

Treatment:

In cases of overdose, ECG monitoring for QT interval prolongation should be initiated.

If CNS symptoms of overdose occur, naloxone may be given as an antidote.

Since the duration of action of Loperamide is longer than that of naloxone, (1 to 3 hours), repeated treatment with naloxone might be indicated. The patient should be kept under constant observation for at least 48 hours in order to detect any possible CNS depression

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antipropulsives: ATC code A07DA03

Loperamide binds to the opiate receptor in the gut wall, reducing propulsive peristalsis, increasing intestinal transit time and enhancing resorption of water and electrolytes. Loperamide increases the tone of the anal sphincter, which helps reduce faecal incontinence and urgency.

In a double blind randomised clinical trial in 56 patients with acute diarrhoea receiving loperamide, onset of anti-diarrhoeal action was observed within one hour following a single 4 mg dose. Clinical comparisons with other antidiarrhoeal drugs confirmed this exceptionally rapid onset of action of loperamide.

5.2 Pharmacokinetic Properties:

Absorption: Most ingested loperamide is absorbed from the gut, but as a result of significant first pass metabolism, systemic bioavailability is only approximately 0.3%.

Distribution: Studies on distribution in rats show a high affinity for the gut wall with a preference for binding to receptors of the longitudinal muscle layer. The plasma protein binding of loperamide is 95%, mainly to albumin. Non-clinical data have shown that loperamide is a P-glycoprotein substrate.

Metabolism: loperamide is almost completely extracted by the liver, where it is predominantly metabolized, conjugated and excreted via the bile. Oxidative N-demethylation is the main metabolic pathway for loperamide, and is mediated mainly through CYP3A4 and CYP2C8. Due to this very high first pass effect, plasma concentrations of unchanged drug remain extremely low.

Elimination: The half-life of loperamide in man is about 11 hours with a range of 9-14 hours. Excretion of the unchanged loperamide and the metabolites mainly occurs through the faeces.

5.3 Preclinical safety data

Acute and chronic studies on loperamide showed no specific toxicity. Results of *in vivo* and *in vitro* studies carried out indicated that loperamide is not genotoxic. In reproduction studies, very high doses (40 mg/kg/day – 240 times the maximum human use level) loperamide impaired fertility and foetal survival in association with maternal toxicity in rats. Lower doses had no effects on maternal or foetal health and did not affect peri- and post-natal development.

Non-clinical in vitro and in vivo evaluation of loperamide indicates no significant cardiac electrophysiological effects within its therapeutically relevant concentration range and at significant multiples of this range (up to 47-fold). However, at extremely high concentrations associated with overdoses (see section 4.4), loperamide has cardiac electrophysiological actions consisting of inhibition of potassium (hERG) and sodium currents, and arrhythmias.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Ludipress (consisting of lactose monohydrate 94%, polyvinylpyrrolidone 6%, 50:50 mixture of kollidon 30 and kollidon CL)

Magnesium stearate

6.2 Incompatibilities

None known.

6.3 Shelf life

24 months unopened.

6.4 Special precautions for storage

Store below 25°C, protect from light and moisture.

6.5 Nature and contents of container

PVC/aluminium foil blister pack.

Pack size: 12 tablets per carton.

6.6 Special precautions for disposal

Not applicable.

7 MARKETING AUTHORISATION HOLDER

Aspar Pharmaceuticals Ltd
Albany House,
Acrewood way,
St Albans,
AL4 0JY,
United Kingdom

Formatted: Indent: Left:
0.76 cm

Formatted: Indent: Left:
0.76 cm

Deleted: ¶

8 MARKETING AUTHORISATION NUMBER(S)

PL 08977/0044

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

08/10/97

10 DATE OF REVISION OF THE TEXT

31/01/2025