

Package Leaflet: Information for the user

Sebco®

Coal tar solution 12% w/w
Salicylic acid 2% w/w
Sulfur for external use 4% w/w

Read all of this leaflet carefully before you start using this medicine because it contains important information for you. Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

- | | |
|--|---|
| 1. What Sebco® is and what it is used for | 4. Possible side effects |
| 2. What you need to know before you use Sebco® | 5. How to store Sebco® |
| 3. How to use Sebco® | 6. Contents of the pack and other information |

1. WHAT SEBCO® IS AND WHAT IT IS USED FOR

Sebco® is an ointment containing coal tar, salicylic acid and sulfur mixed in coconut oil. The mixture of coal tar, salicylic acid and sulfur reduces itch, helps remove scaly skin and has antiseptic properties. Sebco® is used in the treatment of skin diseases which cause dry scaly skin on the scalp. These include psoriasis, eczema, seborrhoeic dermatitis and dandruff.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE SEBCO®

Do not use Sebco® if:

- You know you are allergic to any of the ingredients in Sebco® (see section 6).
- You have an acute infection of an area of skin. Swelling, pain, redness and maybe pus are signs of this.

Make sure your doctor or pharmacist knows if any of the above applies to you.

Take special care with Sebco®:

- Avoid contact with the mouth, mucous membranes, eyes and any areas of broken skin.
- Sebco® can stain clothes and jewellery. Remove or protect these before application.

Other medicines and Sebco®

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine. You should not use Sebco® during pregnancy except for short-term treatment of a small area of the scalp. Oral forms (e.g. tablets) of salicylic acid can cause adverse effects in your unborn baby. It is not known if the same risks apply to Sebco® when it is used on the scalp.

Driving and using machines

Sebco® should not affect your ability to drive or use machinery.

Sebco® contains cetostearyl alcohol

Cetostearyl alcohol may cause a sensitivity rash (e.g. contact dermatitis).

Do not smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

3. HOW TO USE SEBCO®

Sebco® should be used only on your skin.

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

When first opening Sebco® do not use if the tube membrane is already broken.

In severe scaly scalp conditions it is usually necessary to apply Sebco® daily for 3-7 days until control has been obtained.

With mild scaliness use intermittently as necessary, e.g. once a week.



[Illustration 1] Pierce the tube membrane by inverting the cap. Affix the scalp applicator to the threaded portion of the tube.



[Illustration 2] Part the hair and apply a thin ribbon of Sebco® to the affected areas of the scalp. Take care to avoid contact with the eyes. Gently rub in the ointment and leave in contact with scalp for approximately one hour. Wash the hands immediately afterwards.



[Illustration 3] After one hour remove the ointment by showering or rinsing with warm water and shampoo. If after the first week there is no improvement or after 4 weeks symptoms persist, consult your doctor. **If you accidentally get some into your eyes wash it out immediately with plenty of cold tap water.** If irritation of the eyes develops consult your doctor or pharmacist.

Sebco® is not recommended for children under 6 years of age.

If you use more Sebco® than you should or if you accidentally swallow some Sebco®

If you use more Sebco® than you should or if you accidentally swallow some Sebco® contact your nearest hospital casualty department or tell your doctor immediately.

If you forget to use Sebco®

If it is within the day, apply when you remember. If it is the next day do not use a double application, just continue as before.

If you have any further questions on the use of this product ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines Sebco® can sometimes cause side effects, although not everybody gets them.

The following side effects have been reported:

- Irritation of the skin
- Inflammation of the hair follicles causing soreness and small pus filled spots around the base of the hairs
- Acne like eruptions
- Discolouration of skin and hair
- Hair loss
- Rarely, a sensitivity rash to sunlight
- If you have asthma or if anyone in your family is allergic to aspirin, Sebco® may cause difficulty in breathing or a tightness of your chest. This is very rare.

Reporting of side effects

If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE SEBCO®

Keep out of the sight and reach of children. Your ointment could harm them.

Do not use your ointment after the expiry date which is stated on the carton. The expiry date refers to the last day of that month.

Keep your medicine at room temperature, in a dry place, and not above 25°C.

Discard tube no later than 4 weeks after opening.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

The active ingredients are coal tar 12% w/w, salicylic acid 2% w/w and sulfur for external use 4% w/w. The other ingredients are coconut oil, white soft paraffin, cetostearyl alcohol, glycerol, liquid paraffin, polyoxyethylene glycerol monostearate, hard paraffin.

What Sebco® looks like and contents of pack

Sebco® is a buff coloured ointment. It is licensed for sale in tubes of 5, 15, 40 and 100 grams.

Availability may be limited in some sizes.

Marketing authorisation holder

Derma UK Ltd, The Toffee Factory, Lower Steenbergs Yard, Ouseburn, Newcastle upon Tyne NE1 2DF, UK.

Manufacturer

Ovelle Limited, Coes Road, Dundalk, Co. Louth, Ireland. 'Sebco®' is a registered trademark.

This leaflet was last revised in April 2025. xxxx

CLIENT / PRODUCT	Derma UK Sebco PIL	SIZE mm	208 x 147	PROOF
BRIEF		MOD		COLOURS PMS
PHARMA CODE	-	PROGS	Illustrator CS6	Black
COMPONENT	-	FONTS	Helvetica Neue 45,55,75	
BAR CODE	-		(ITC BT Franklin Bk, Med, Demi)	

Package Leaflet: Information for the user

Sebco®

Coal tar solution 12% w/w
Salicylic acid 2% w/w
Sulfur for external use 4% w/w

Read all of this leaflet carefully before you start using this medicine because it contains important information for you. Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Sebco® is and what it is used for
2. What you need to know before you use Sebco®
3. How to use Sebco®
4. Possible side effects
5. How to store Sebco®
6. Contents of the pack and other information

1. WHAT SEBCO® IS AND WHAT IT IS USED FOR

Sebco® is an ointment containing coal tar, salicylic acid and sulfur mixed in coconut oil. The mixture of coal tar, salicylic acid and sulfur reduces itch, helps remove scaly skin and has antiseptic properties. Sebco® is used in the treatment of skin diseases which cause dry scaly skin on the scalp. These include psoriasis, eczema, seborrhoeic dermatitis and dandruff.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE SEBCO®

Do not use Sebco® if:

- You know you are allergic to any of the ingredients in Sebco® (see section 6).
- You have an acute infection of an area of skin. Swelling, pain, redness and maybe pus are signs of this.

Make sure your doctor or pharmacist knows if any of the above applies to you.

Take special care with Sebco®:

- Avoid contact with the mouth, mucous membranes, eyes and any areas of broken skin.
- Sebco® can stain clothes and jewellery. Remove or protect these before application.

Other medicines and Sebco®

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine. You should not use Sebco® during pregnancy except for short-term treatment of a small area of the scalp. Oral forms (e.g. tablets) of salicylic acid can cause adverse effects in your unborn baby. It is not known if the same risks apply to Sebco® when it is used on the scalp.

Driving and using machines

Sebco® should not affect your ability to drive or use machinery.

Sebco® contains cetostearyl alcohol

Cetostearyl alcohol may cause a sensitivity rash (e.g. contact dermatitis).

Do not smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

3. HOW TO USE SEBCO®

Sebco® should be used only on your skin.

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

When first opening Sebco® do not use if the tube membrane is already broken.

In severe scaly scalp conditions it is usually necessary to apply Sebco® daily for 3-7 days until control has been obtained.

With mild scaliness use intermittently as necessary, e.g. once a week.



[Illustration 1] Pierce the tube membrane by inverting the cap. Affix the scalp applicator to the threaded portion of the tube.



[Illustration 2] Part the hair and apply a thin ribbon of Sebco® to the affected areas of the scalp. Take care to avoid contact with the eyes. Gently rub in the ointment and leave in contact with scalp for approximately one hour. Wash the hands immediately afterwards.



[Illustration 3] After one hour remove the ointment by showering or rinsing with warm water and shampoo. If after the first week there is no improvement or after 4 weeks symptoms persist, consult your doctor. **If you accidentally get some into your eyes wash it out immediately with plenty of cold tap water.** If irritation of the eyes develops consult your doctor or pharmacist.

Sebco® is not recommended for children under 6 years of age.

If you use more Sebco® than you should or if you accidentally swallow some Sebco®

If you use more Sebco® than you should or if you accidentally swallow some Sebco® contact your nearest hospital casualty department or tell your doctor immediately.

If you forget to use Sebco®

If it is within the day, apply when you remember. If it is the next day do not use a double application, just continue as before.

If you have any further questions on the use of this product ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines Sebco® can sometimes cause side effects, although not everybody gets them.

The following side effects have been reported:

- Irritation of the skin
- Inflammation of the hair follicles causing soreness and small pus filled spots around the base of the hairs
- Acne like eruptions
- Discolouration of skin and hair
- Hair loss
- Rarely, a sensitivity rash to sunlight
- If you have asthma or if anyone in your family is allergic to aspirin, Sebco® may cause difficulty in breathing or a tightness of your chest. This is very rare.

Reporting of side effects

If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE SEBCO®

Keep out of the sight and reach of children. Your ointment could harm them.

Do not use your ointment after the expiry date which is stated on the carton. The expiry date refers to the last day of that month.

Keep your medicine at room temperature, in a dry place, and not above 25°C.

Discard tube no later than 4 weeks after opening.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

The active ingredients are coal tar 12% w/w, salicylic acid 2% w/w and sulfur for external use 4% w/w. The other ingredients are coconut oil, white soft paraffin, cetostearyl alcohol, glycerol, liquid paraffin, polyoxyethylene glycerol monostearate, hard paraffin.

What Sebco® looks like and contents of pack

Sebco® is a buff coloured ointment. It is licensed for sale in tubes of 5, 15, 40 and 100 grams.

Availability may be limited in some sizes.

Marketing authorisation holder

Derma UK Ltd, The Toffee Factory, Lower Steenbergs Yard, Ouseburn, Newcastle upon Tyne NE1 2DF, UK.

Manufacturer

Bioglan AB, PO Box 50310, S-202, 13 Malmö, Sweden. 'Sebco®' is a registered trademark.

This leaflet was last revised in April 2025. xxxx

CLIENT / PRODUCT	Derma UK Sebco PIL	SIZE mm	208 x 147	PROOF	
BRIEF		MOD		COLOURS PMS	
PHARMA CODE	-	PROGS	Illustrator CS6	Black	
COMPONENT	-	FONTS	Helvetica Neue 45,55,75		
BAR CODE	-		(ITC BT Franklin Bk, Med, Demi)		