



Back Page

- If you take a large overdose of Mentino, contact your doctor or get medical advice, as you may need medical attention.

If you forget to take Mentino

- If you find you have forgotten to take your dose of Mentino, wait and take your next dose at the usual time.
- Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

In general, the observed side effects are mild to moderate.

Common (affects 1 to 10 users in 100):

- Headache, sleepiness, constipation, elevated liver function tests, dizziness, balance disorders, shortness of breath, high blood pressure and drug hypersensitivity

Uncommon (affects 1 to 10 users in 1,000):

- Tiredness, fungal infections, confusion, hallucinations, vomiting, abnormal gait, heart failure and venous blood clotting (thrombosis/thromboembolism)

Very Rare (affects less than 1 user in 10,000):

- Seizures

Not known (frequency cannot be estimated from the available data):

- Inflammation of the pancreas, inflammation of the liver (hepatitis) and psychotic reactions

Alzheimer’s disease has been associated with depression, suicidal ideation and suicide. These events have been reported in patients treated with Mentino.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

United Kingdom

Yellow Card Scheme Website:  
www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store  
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Mentino

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and the blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Mentino contains

- The active substance is memantine hydrochloride.  
Each tablet contains 5 mg of memantine hydrochloride equivalent to 4.15 mg memantine.  
Each tablet contains 10 mg of memantine hydrochloride equivalent to 8.31 mg memantine.  
Each tablet contains 15 mg of memantine hydrochloride equivalent to 12.46 mg memantine.  
Each tablet contains 20 mg of memantine hydrochloride equivalent to 16.62 mg memantine.
- The other ingredients for Mentino 5 mg, 10 mg, 15 mg and 20 mg Orodispersible tablets are Polacrilin, Sodium hydroxide (E 524), Water purified, Lactose monohydrate, Cellulose microcrystalline (E 460), Mannitol (E 421), Croscarmellose sodium (E 468), Aspartame (E 951), Silica colloidal anhydrous, Iron oxide red (E 172), Flavor peppermint, Magnesium stearate (E 572)

What Mentino looks like and contents of the Pack

Mentino 5 mg Orodispersible tablets are presented as light pink round, flat, speckled tablets with beveled edges, a diameter of 7 mm and engraved with “5” on one side.  
Mentino 10 mg Orodispersible tablets are presented as light pink, round, flat, speckled tablets with beveled edges, a diameter of 9 mm and engraved with “10” on one side.  
Mentino 15 mg Orodispersible tablets are presented as light pink, round, flat, speckled tablets with beveled edges, a diameter of 11 mm and engraved with “15” on one side.  
Mentino 20 mg Orodispersible tablets are presented as Light pink, round, flat, speckled tablets with beveled edges, a diameter of 12 mm and engraved with “20” on one side.

5 mg:

Aluminium/ peelable Aluminium blisters

Each pack contains 7 orodispersible tablets.

10 mg:

Aluminium/ peelable Aluminium blisters

Pack sizes of 7, 14, 28, 30, 42, 50, 56, 70, 84, 98, 100,112 Orodispersible tablets.

Multipack containing 980 (10 packs of 98) and 1000 (20 packs of 50) Orodispersible tablets.

Perforated unit dose blister: Aluminium/ peelable Aluminium blisters

Pack sizes 49 x 1, 56 x1, 98 x 1 and 100 x 1 Orodispersible tablets  
x

Not all pack sizes may be marketed.

15 mg:

Aluminium/ peelable Aluminium blisters

Each pack contains 7 orodispersible tablets.

20 mg:

Aluminium/ peelable Aluminium blisters

Pack sizes of 7, 14, 28, 42, 56, 70, 84, 98, 112 Orodispersible tablets.

Multipack containing 840 (20 x 42) Orodispersible tablets.

Perforated unit dose blister: Aluminium/ peelable Aluminium blisters

Pack sizes 49 x 1, 56 x1, 98 x 1 and 100 x 1 Orodispersible tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer  
Marketing Authorisation Holder

Novumgen Limited  
20-22 Wenlock Road, London,  
N1 7GU, United Kingdom

Manufacturer

GENEPHARM S.A.  
18 km Marathonos Avenue  
15351 Pallini Attiki  
Greece

Rontis Hellas S.A.  
Medical and Pharmaceutical Products S.A.  
P.O. Box 3012 Larisa Industrial Area, Larisa,  
41004, Greece

“If you would like a leaflet with larger text, please contact +44 7483273058”.

This leaflet was last revised in SEP 2024.



code

Artwork Number may be added

|                                                                           |        |
|---------------------------------------------------------------------------|--------|
|                                                                           |        |
| Product :<br>Mentino 5 mg, 10 mg, 15 mg, 20 mg,<br>Orodispersible Tablets |        |
| Version                                                                   | : V2   |
| Sequence Number                                                           | : 0001 |