

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if your symptoms worsen or do not improve.

1. What Lyclear Dermal Cream is and what it is used for
2. What you need to know before you use Lyclear Dermal Cream
3. How to use Lyclear Dermal Cream
4. Possible side effects
5. How to store Lyclear Dermal Cream
6. Contents of the pack and other information

Lyclear Dermal Cream is used to treat infestations of scabies and crab lice on the body. It contains the ingredient permethrin which belongs to a group of medicines called pyrethroids which kill mites such as scabies and crab lice.

Do not use Lyclear Dermal Cream:

- If you are allergic to Permethrin or any of the other ingredients of this medicine (listed in section 6).

Talk to your doctor or pharmacist before using Lyclear Dermal Cream:

- Do not smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.
- For cutaneous use only! Do not swallow this medicine.

Avoid contact with eyes or mucous membranes (inside the nose or throat, genital region) or open wounds.

Carers who apply permethrin are advised to wear gloves.

Do not use Lyclear Dermal Cream in newborns and infants less than 2 months of age, unless your doctor tells you so. There is no adequate experience in infants and toddlers. Treatment to children up to 23 months of age should only be given under close medical supervision.

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. Corticosteroids (steroids) should not be used to treat skin conditions like eczema when Lyclear Dermal Cream is being used because they may make the infestation of scabies and lice worse by interfering with the body's natural defence to scabies and crab lice. Lyclear Dermal Cream is not known to interact with any medicines.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

No effects are known on ability to drive and use machines.

Lyclear Dermal Cream contains butylated hydroxytoluene (E321) which may cause local skin reactions (e.g. contact dermatitis), or irritation to the eyes and mucous membranes (lining of the nose, mouth and genital areas). Formaldehyde which may cause local skin reactions (e.g. contact dermatitis). Lyclear Dermal Cream contains paraffins. These excipients of the cream can reduce the efficiency and hence the reliability of latex products (e.g. condoms, diaphragms) used at the same time.

Lyclear Dermal Cream contains lanolin which may cause local skin reactions (e.g. contact dermatitis).

Lyclear Dermal Cream is harmful to all types of insects and also to animals living in water e. g. fish.

Take care that Lyclear Dermal Cream does not get into aquaria or terraria.

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. To pierce tube, remove the cap and reverse it over the nozzle, push firmly and turn. Carefully apply a thin layer of cream to the skin (see "How and when should you use Lyclear Dermal Cream?").

Apply the cream to clean, dry, cool skin. Do not take a hot bath or shower immediately before treatment.

Adults and adolescents over 12 years of age. Apply up to 30 g of cream (corresponding to one tube of 30 g). The application should be repeated 7 – 14 days after the first treatment.

Children aged from 6 - 12 years. Apply up to 15 g of cream (corresponding to ½ tube of 30 g). The application should be repeated 7 - 14 days after the first treatment.

Children aged from 2 months - 5 years. Apply up to 7.5 g of cream (corresponding to ¼ tube of 30 g). The application should be repeated 7 - 14 days after the first treatment.

Newborns and infants under 2 months of age. There is only limited amount of data available in this age group and no dose can be recommended (see also section 2 under “warnings and precautions”).

The above information is merely a guide. The dose can be adjusted according to the needs of the individual patient and the individual body surface area. For example, some adults require a larger amount of cream.

Brand:				Application:				Item code(s):		Market:		Master text version:				Barcode 1 type:								Component size (flat):				Cutter guide nr.		Manufacturer:	
Lyclear				Cream				LF00260F		United Kingdom						Barcode 1 nr:								200x135mm							
Variant:				Size:		Unit of measurement:		Languages on the component:				Packaging type:				Barcode 2 type:								Folded size:				PO nr:		Printer:	
Dermal cream 5%				30g				English				leaflet				Barcode 2 nr:												CR08501			
				<div>1: Black</div> <div>2: PANTONE 300 C</div> <div>3: Cutter guide</div> <div>4:</div>								<div>5:</div> <div>6:</div> <div>7:</div> <div>8:</div>				<div>Marketing Approval:</div> <div>Name:</div> <div>Date: ... / ... / ...</div> <div>Signature:</div>				<div>Regulatory Approval:</div> <div>Name:</div> <div>Date: ... / ... / ...</div> <div>Signature:</div>				WW							
Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:	Pass nr:	Pass date:														
0	29-08-2023	1	23-01-2024	2	-	3		4		5		6		7		8															

Adults over 18 years of age: Apply up to 30 g of cream (corresponding to one tube of 30 g).

Lyclear Dermal Cream is for cutaneous use only.

For treatment of scabies: Adults, adolescents and children over 2 years should apply the cream to the whole body including the neck, palms of the hands and soles of the feet. The head and face can be spared unless this area includes places affected by scabies (itch mites).

Lyclear Dermal Cream is a vanishing cream and when rubbed gently into the skin it will disappear. Therefore, there is no need to continue to apply cream to the skin until it remains detectable on the surface.

neck, face, ears, and scalp. Parts of the skin around the mouth (because the cream could be licked off) and the eyes should be spared. Keep your child from licking the cream from the hands. If necessary, children should wear gloves.

There is no adequate experience in infants and toddlers. Treatment to children up to the age of 23 months should therefore only be treated under close medical supervision.

Elderly: Elderly patients (over 65 years) should use the cream in the same way as adults but in addition, the face, ears and scalp should also be treated. Care should be taken to avoid applying the cream to areas of skin around the eyes.

For treatment of crab lice in adults: Any facial hair (beards moustaches) and eyelashes should be checked for the presence of live lice and eggs. Cream should be applied to any facial hair, carefully avoiding eyes. If live lice or eggs are found on the eyelashes they should be removed carefully using tweezers. Cream should not be applied to the eye lashes because the cream can cause moderate eye irritation.

Sufficient cream should be applied to cover the pubic region, peri-anal (around the anus), inner thighs down to the knees and any hair that grows up from the pubic area to the chest/stomach.

For scabies: Leave the cream on the skin for at least eight hours, for example, overnight. Avoid bathing, showering or washing during this period, because this could endanger the success of treatment. If, by way of an exception, you have to wash your hands within the 8-12 hour period, then reapply the cream to the hands and wrist area. The same applies if you have to wash other parts of treated skin (buttocks, external genitalia).

After at least eight hours, take a shower or wash the skin with soap and water. The application should be repeated 7 – 14 days after the first treatment.

For crab lice: The cream should be left on for 24 hours and then washed off with soap and water.

If too much cream is applied, it should be washed off with soap and warm water. If there are any signs of irritation, talk to your doctor.

If the contents of the tube are swallowed by a child get medical aid from the doctor or Accident and Emergency Department of the nearest hospital immediately.

Do not apply a double quantity to make up for a forgotten application.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If severe hypersensitivity reactions occur, please consult a doctor immediately! In this case you must not use Lyclear Dermal Cream any more.

Common: may affect up to 1 in 10 people.

itching (pruritus), reddening of the skin or unusual sensations on the skin (paraesthesias) such as tingling, pricking, skin burning sensation as well as dry skin are common. However, such symptoms can also occur as a result of the disease itself. The Itching and a skin rash (post-scabies eczema) may persist for up to four weeks after the end of treatment. This is caused by a reaction to the killed scabies mites. If after using Lyclear Dermal Cream you have the impression that the disease is persisting, please speak to your doctor before applying it again.

Rare: may affect up to 1 in 1,000 people. Headache can occur rarely.

Very rare: may affect up to 1 in 10,000 people.

Very rarely, skin lesions due to skin picking disorder (excoriations), inflammation of the hair follicles (folliculitis) and reduced skin pigmentation have been reported at the time Lycleair Dermal Cream is used.

Sensitive/allergic persons have reported breathing difficulties at the time substances from the pyrethrin group were being used.

Not known: frequency cannot be estimated from the available data.

Intolerability reactions may occur on the skin (contact allergy reactions) that are expressed as itching, reddening, blisters or nettle rash (urticaria). These reactions may also spread beyond the area of skin treated. Nausea may appear. Vomiting was not reported after the use of Lyclear Dermal Cream but is known in connection with other permethrin-containing drugs.

Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

Keep this medicine out of the sight and reach of children. Do not store above 25°C. Do not freeze. Do not use this medicine after the expiry date which is stated on the carton after "Exp". The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

What Lyclear Dermal Cream contains

- The active substance is Permethrin (5%w/w)
- The other ingredients are Coconut oil, glyceryl monostearate, macrogol cetyl ether, cetomacrogol 1000, isopropyl myristate, lanolin alcohols/liquid paraffin mixture, butylated hydroxytoluene (E321), glycerol, formaldehyde solution, carbomer, sodium hydroxide and purified water.

Lyclear Dermal Cream is a white cream in a 30 g tube.

Marketing Authorisation holder: Omega Pharma Ltd, Wrafton, Braunton, Devon, EX33 2DL, UK. Send all enquiries to this address.

Manufacturer: Medgenix Benelux NV, Vliegveld 21, BE-8560 Wevelgem, Belgium.

Lyclear is a registered trademark. Date of text revision: July 2023.

8100002972
LF00260F

FOR APPROVAL • Please valid this proof by signature and send it back to us. Thank you. Please verify this proof carefully concerning lay-out, size, texts and bar code. For the PMS colours please refer to the latest PMS colour guide. After your approval on the above we can proceed to the next stage of your project.