



88 mm

217 mm

118 mm

11 mm

8 mm

Direction of reading

10 mm

75 mm

39.99 mm

31042293

For the treatment of eczema and other dry skin conditions

•Moisturising and protective

•Relieves and soothes

Diprob

base

Cream

Emollient

500g

Store below 25°C.

Contains: white soft paraffin, cetostearyl alcohol, liquid paraffin, macrogol cetostearyl ether, chlorocresol, sodium dihydrogen phosphate, sodium hydroxide, phosphoric acid and purified water.

See leaflet for further information.

PL 35104/0054 PA 1113/030/001

MAH (UK and Ireland)

Phoenix Labs

Suite 12, Bunkilla Plaza, Bracetown Business Park, Clonee, Co. Meath, Ireland

5 010605 240225

BN:

EXP:

PHOENIX LABS

20 mm

113 mm

358 mm

129 mm

4

FLX

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed on this label. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

UK: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)

Ireland: [www.hpra.ie](http://www.hpra.ie)

**5. HOW TO STORE DIPROBASE**

The cream should not be stored above 25°C.

Keep out of the sight and reach of children.

Do not use the cream after the expiry date “EXP” which is stamped on the pack. This refers to the last date of that month.

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

**6. FURTHER INFORMATION**

**What Diprob**

The ingredients are white soft paraffin, cetostearyl alcohol, liquid paraffin, macrogol cetostearyl ether, chlorocresol, sodium dihydrogen phosphate, sodium hydroxide, phosphoric acid and purified water.

**What Diprob**

Diprob

The holder of the Marketing Authorisation is:

**MAH (UK and Ireland):** Phoenix Labs, Suite 12, Bunkilla Plaza, Bracetown Business Park, Clonee, Co. Meath, Ireland.

**The manufacturer is:**

Famar A.V.E. Avlon Plant, 49th km National Road Athens-Lamia, Avlona Attiki, 19011, Greece

**This leaflet was last revised in February 2024.**

For the treatment of eczema and other dry skin conditions. Diprob

red, inflamed, damaged, dry or chapped skin. Can also be used to protect raw skin and can be applied before bathing.

**Directions for use:**

Apply a thin layer of cream and massage thoroughly into affected areas of skin.

**For cutaneous use.**

**Keep out of the sight and reach of children.**

For external use only

PHOENIX LABS

Store below 25°C.

Contains: white soft paraffin, cetostearyl alcohol, liquid paraffin, macrogol cetostearyl ether, chlorocresol, sodium dihydrogen phosphate, sodium hydroxide, phosphoric acid and purified water.

See leaflet for further information.

PL 35104/0054 PA 1113/030/001

**MAH (UK and Ireland)**

Phoenix Labs

Suite 12, Bunkilla Plaza, Bracetown Business Park, Clonee, Co. Meath, Ireland

5 010605 240225

129 mm

358 mm

116 mm

113 mm

75 mm

63 mm

12 mm

**PACKAGE LEAFLET: INFORMATION FOR THE USER**

**Diprob**

**Read all of this leaflet carefully before you start using this medicine.**

Keep this leaflet. You may need to read it again.

If you have any further questions, ask your doctor or pharmacist.

This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What Diprob

2. Before you use Diprob

3. How to use Diprob

4. Possible side effects

5. How to store Diprob

6. Further information

**1. WHAT DIPROBASE IS AND WHAT IT IS USED FOR**

Diprob

In adults and children, Diprob

In adults and children, Diprob

Eczema is a common skin disease, which causes the skin to become red and itchy.

**2. BEFORE YOU USE DIPROBASE**

**Do not use DIPROBASE**

if you, or your child, have ever had an allergic reaction to Diprob

Ask your doctor or pharmacist if you are not sure.

**Important information about some of the ingredients of DIPROBASE**

Diprob

If you are worried about this or any other effects, you should contact your doctor or pharmacist.

**3. HOW TO USE DIPROBASE**

Always use Diprob

For adults and children, the cream should be applied thinly to cover the affected area completely, massaging gently and thoroughly into the skin.

If the symptoms persist or the condition worsens, stop using the cream and consult your doctor or pharmacist.

**If you forget to use DIPROBASE**

If you forget to use Diprob

on as before.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, Diprob

If your symptoms persist or the condition worsens, stop using the cream and consult your doctor or pharmacist.

Skin reactions including itching, rash, redness, peeling, burning, pain, dryness and skin inflammation (dermatitis) have been reported with product use.

If you are worried by this or any other effects, you should tell your doctor or pharmacist.

11 mm

