



**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**

Sodium alginate, Sodium bicarbonate, Calcium carbonate

Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.

**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:  
 1) Neutralising excess stomach acid to relieve the pain and discomfort.  
 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.  
 Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).

**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**  
 • You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).  
**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.  
 The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.  
 You can take this product if you are pregnant or breast feeding.  
 This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).

**Turning the leaflet for the further information**  
**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor  
**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat.  
 If you experience these or any other side-effects stop taking the product and consult your doctor immediately.  
 Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.

**How to take Gaviscon Double Action Mint**  
**Do not take this product** within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines.  
 This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).  
**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product. For oral use. Shake well before use.  
**Adults and children 12 years and over:** Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice.  
 If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

**How to store Gaviscon Double Action Mint**  
 • Do not use this product after the expiry date (EXP: month/year) shown  
 • Use within 6 months of opening • Keep out of the reach and sight of children  
 • Do not store above 30°C. Do not refrigerate or freeze.

**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients.  
 The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten. Gaviscon Double Action Mint is available in bottles of 150ml, 300ml & 600ml.  
 Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
 Export distributors: Reckitt & Colman (Overseas) Ltd., Hull, HU8 7DS, UK.  
 Gaviscon is a trade mark of the RB group of companies  
 Text Revised: October 2020



**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate, Sodium bicarbonate, Calcium carbonate  
 Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.  
**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and alginate and works in two ways:  
 1) Neutralising excess stomach acid to relieve the pain and discomfort.  
 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.  
 Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).  
 page 1

**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**  
 • You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).  
**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in salt, please consult your doctor.  
 The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.  
 You can take this product if you are pregnant or breast feeding.  
 This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).  
 page 2

**Taking Other Medicines**  
 Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines.  
 This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).  
**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product. For oral use. Shake well before use.  
**Adults and children 12 years and over:** Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice.  
 If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.  
 page 3

Turn the leaflet for the further information  
**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor  
**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat.  
 If you experience these or any other side effects stop taking the product and consult your doctor immediately.  
 Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.  
 page 4

**How to store Gaviscon Double Action Mint**  
 • Do not use this product after the expiry date (EXP: month/year) shown  
 • Use within 6 months of opening • Keep out of the reach and sight of children  
 • Do not store above 30°C. Do not refrigerate or freeze.  
**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients.  
 The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar.  
 Gaviscon Double Action Mint is available in bottles of 150ml, 300ml & 600ml.  
 Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
 Export distributors: Reckitt & Colman (Overseas) Ltd., Hull, HU8 7DS, UK.  
 Gaviscon is a trade mark of the RB group of companies  
 Text Revised: October 2020  
 page 5


Page 6  
 page 6



**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate  
 Sodium bicarbonate  
 Calcium carbonate  
 Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.  
**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:  
 1) Neutralising excess stomach acid to relieve the pain and discomfort.  
 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.  
 Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal

reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).  
**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**  
 • You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).  
**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these


please consult your doctor.  
 The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.  
 You can take this product if you are pregnant or breast feeding.  
 This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).  
**Taking Other Medicines**  
 Do **not** take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines.



antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).  
**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice. If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor.  
**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.  
**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown  
 • Use within 6 months of opening

• Keep out of the reach and sight of children  
 • Do not store above 30°C  
 • Do not refrigerate or freeze  
**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients.  
 The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten.  
 Gaviscon Double Action Mint is available in 150ml, 200ml, 250ml, 300ml, 500ml and 600ml bottles. Not all pack sizes may be marketed.  
 Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
 Gaviscon is a trade mark of the Reckitt group.  
 Text Revised: August 2023



English TBR2662-D8340700

gaviscon  
 double action  
 mint

BARCODE INFO	
Barcode Type:	EAN 13
Code Value:	0000000000000
Magnification:	100 %
Overall Height:	25.98 mm
Bar Height:	22.9 mm
BWR Applied:	20 Micrometre



**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate  
 Sodium bicarbonate  
 Calcium carbonate  
 Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.  
**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:  
 1) Neutralising excess stomach acid to relieve the pain and discomfort.  
 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.  
 Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).  
**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**  
 • You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).  
**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.  
 The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.  
 You can take this product if you are pregnant or breast feeding.  
 This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).  
**Taking Other Medicines**  
 Do **not** take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines.

antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).  
**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice. If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.  
**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor.  
**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.  
**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown  
 • Use within 6 months of opening  
 • Keep out of the reach and sight of children  
 • Do not store above 30°C  
 • Do not refrigerate or freeze  
**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients.  
 The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten.  
 Gaviscon Double Action Mint is available in 150ml, 200ml, 250ml, 300ml, 500ml and 600ml bottles. Not all pack sizes may be marketed.  
 Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
 Gaviscon is a trade mark of the Reckitt group.  
 Text Revised: August 2023

English TBR2662-D8340700

gaviscon  
 double action  
 mint

177.8 mm

70 mm



PPTD1022/3  
PRINTING READS This Way


66.3mm

55mm

30 mm

**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
Sodium alginate  
Sodium bicarbonate  
Calcium carbonate  
Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.  
**What is Gaviscon Double Action Mint and what is it used for?**  
Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:  
1) Neutralising excess stomach acid to relieve the pain and discomfort.  
2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.  
Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).  
**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**  
• You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).  
**Take special care before treatment with Gaviscon Double Action Mint:**  
This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.  
The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.  
You can take this product if you are pregnant or breast feeding.  
This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).  
**Taking Other Medicines**  
Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines.

antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).  
**How to take Gaviscon Double Action Mint**  
Check that the cap seal is unbroken before first using this product.  
For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice. If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.  
**After taking Gaviscon Double Action Mint:**  
If symptoms persist after 7 days consult your doctor.  
**Possible side effects**  
Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.  
**How to store Gaviscon Double Action Mint**  
Do not use this product after the expiry date (EXP: month/year) shown  
• Use within 6 months of opening  
• Keep out of the reach and sight of children  
• Do not store above 30°C  
• Do not refrigerate or freeze  
**Further information**  
Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients.  
The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten.  
Gaviscon Double Action Mint is available in 150ml, 300ml & 600ml bottles.  
Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
Gaviscon is a trade mark of the RB group of companies  
Text Revised: October 2020




**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate  
 Sodium bicarbonate  
 Calcium carbonate  
 Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.  
**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:  
 1) Neutralising excess stomach acid to relieve the pain and discomfort.  
 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.  
 Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).  
**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**  
 • You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).  
**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.  
 The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.  
 You can take this product if you are pregnant or breast feeding.  
 This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).  
**Taking Other Medicines**  
 Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines.

antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).  
**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice. If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.  
**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor.  
**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.  
**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown  
 • Use within 6 months of opening  
 • Keep out of the reach and sight of children  
 • Do not store above 30°C  
 • Do not refrigerate or freeze  
**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients.  
 The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten.  
 Gaviscon Double Action Mint is available in 150ml, 300ml & 600ml bottles.  
 Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
 Gaviscon is a trade mark of the RB group of companies  
 Text Revised: October 2020


197mm

Do not store above 30°C. Do not refrigerate or freeze. Use within 6 months of opening.  
**KEEP OUT OF THE SIGHT AND REACH OF CHILDREN.**  
Glass bottle, handle with care. Do not drink directly from the bottle.



Peel here. Do not remove.


e500ml



**DISPENSING PACK ONLY.  
NOT FOR RETAIL SALE.**

• Sodium alginate • Sodium bicarbonate • Calcium carbonate

**WHAT IS THIS PRODUCT?** Each 10ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate.  
For oral use. Shake well before use. If symptoms persist after 7 days consult your doctor.  
Contains sodium, methyl (E218) and propyl (E216) parahydroxybenzoates. High in sodium.  
**See leaflet for further information.**  
Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
PL 00063/0552 Gaviscon is a trade mark of the RB group of companies




80.5mm
60mm
56.5mm
30mm

75mm
27.5mm

<p><b>PATIENT INFORMATION LEAFLET</b> <b>GAVISCON DOUBLE ACTION MINT</b> Sodium alginate Sodium bicarbonate Calcium carbonate</p> <p>Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.</p> <p><b>What is Gaviscon Double Action Mint and what is it used for?</b> Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways: (1) Neutralising excess stomach acid to relieve the pain and discomfort. (2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest. Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid</p>	<p>regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).</p> <p><b>Before taking Gaviscon Double Action Mint</b> <b>Do not take this product if:</b></p> <ul style="list-style-type: none"> <li>You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).</li> </ul> <p><b>Take special care before treatment with Gaviscon Double Action Mint:</b> This medicine contains calcium (3.25 mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please</p>	<p>consult your doctor. The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet. You can take this product if you are pregnant or breast feeding. This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).</p> <p><b>Taking Other Medicines</b> Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as</p>	
--	---	---	--

<p>norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).</p> <p><b>How to take Gaviscon Double Action Mint</b> Check that the cap seal is unbroken before first using this product. For oral use. Shake well before use.</p> <p><b>Adults and children 12 years and over:</b> Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day, or as directed by your doctor. <b>Children under 12 years:</b> Should only be taken on medical advice. If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.</p> <p><b>After taking Gaviscon Double Action</b></p>	<p><b>Mint:</b> If symptoms persist after 7 days consult your doctor.</p> <p><b>Possible side effects</b> Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <a href="http://www.mhra.gov.uk/yellowcard">http://www.mhra.gov.uk/yellowcard</a> or search for MHRA Yellow Card in the Google Play or Apple App Store.</p> <p><b>How to store Gaviscon Double Action Mint</b> Do not use this product after the expiry date (EXP: month/year) shown</p>	<ul style="list-style-type: none"> <li>Use within 6 months of opening</li> <li>Keep out of the sight and reach of children</li> <li>Do not store above 30°C.</li> <li>Do not refrigerate or freeze</li> </ul> <p><b>Further information</b> Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients. The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten. Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS. Gaviscon is a trade mark of the RB group of companies Text Revised: October 2020</p>
---	--	--





**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate, Sodium bicarbonate, Calcium carbonate

Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.

**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:

- 1) Neutralising excess stomach acid to relieve the pain and discomfort.
- 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.

Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).

**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.

The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet. You can take this product if you are pregnant or breast feeding.

This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).

**Taking Other Medicines**  
 Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as

norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).

**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice.  
 If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist.  
 If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor.

**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat.  
 If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.

**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown


- Use within 6 months of opening

- Keep out of the sight and reach of children
- Do not store above 30°C.
- Do not refrigerate or freeze


**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients. The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten. Gaviscon Double Action Mint is available in bottles of 150ml, 300ml, 500ml & 600ml. Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS. Gaviscon is a trade mark of the RB group of companies  
 Text Revised: October 2020

TBR2682-D8340705 - English

Gaviscon  
 Double Action  
 Mint



**CUSTOMER INFO:**  
 Minimum Point Size = 7.42pt



**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate, Sodium bicarbonate, Calcium carbonate

Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.

**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:

- 1) Neutralising excess stomach acid to relieve the pain and discomfort.
- 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.

Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).

**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.

The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet. You can take this product if you are pregnant or breast feeding. This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).

**Taking Other Medicines**  
 Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as 0000000)

norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).

**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice.  
 If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist.  
 If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

**After taking Gaviscon Double Action Mint:**  
 If symptoms persist after 7 days consult your doctor.

**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.

**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown

- Use within 6 months of opening


- Keep out of the sight and reach of children
- Do not store above 30°C.
- Do not refrigerate or freeze

**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients. The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water. This product does not contain sugar or gluten. Gaviscon Double Action Mint is available in bottles of 150ml, 300ml, 500ml & 600ml. Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS. Gaviscon is a trade mark of the RB group of companies  
 Text Revised: October 2020


0000000

TBR2682-D8340705 - English

**Gaviscon  
 Double Action  
 Mint**



**CUSTOMER INFO:**  
 Minimum Point Size = 7.42pt



**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate, Sodium bicarbonate, Calcium carbonate

Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.

**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:

- 1) Neutralising excess stomach acid to relieve the pain and discomfort.
- 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.

Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).

**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.

The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.

You can take this product if you are pregnant or breast feeding.

This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).

**Taking Other Medicines**  
 Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and

0000000

beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).

**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.  
**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.  
**Children under 12 years:** Should only be taken on medical advice.  
 If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist.  
 If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

**After taking Gaviscon Double Action Mint:**

If symptoms persist after 7 days consult your doctor.

**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat.  
 If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.


**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown


- Use within 6 months of opening
- Keep out of the sight and reach of children

• Do not store above 30°C.  
 • Do not refrigerate or freeze

**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients. The other ingredients are sodium carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water.  
 This product does not contain sugar or gluten.  
 Gaviscon Double Action Mint is available in bottles of 150ml, 300ml & 600ml.  
 Manufacturer and PL Holder in UK:  
 Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.  
 Gaviscon is a trade mark of the RB group of companies  
 Text Revised: October 2020

0000000





**PATIENT INFORMATION LEAFLET**  
**GAVISCON DOUBLE ACTION MINT**  
 Sodium alginate, Sodium bicarbonate, Calcium carbonate

Please read this leaflet carefully before you take this medicine. If you are not sure about anything ask your pharmacist or doctor.

**What is Gaviscon Double Action Mint and what is it used for?**  
 Gaviscon Double Action Mint is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate and works in two ways:

- 1) Neutralising excess stomach acid to relieve the pain and discomfort.
- 2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest.

Gaviscon Double Action Mint is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy and for symptoms of excess stomach acidity (hyperacidity).

**Before taking Gaviscon Double Action Mint**  
**Do not take this product if:**

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

**Take special care before treatment with Gaviscon Double Action Mint:**  
 This medicine contains calcium (3.25mmol per 10ml). This medicine contains 127.88 mg sodium (main component of cooking/table salt) in each 10 ml. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. If you have been advised to follow a diet restricted in either of these please consult your doctor.

The maximum recommended daily dose of this medicinal product contains 1023.04 mg sodium (found in table salt). This is equivalent to 51.15 % of the adult recommended maximum daily dietary intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period of time, especially if you have been advised to follow a low salt (sodium) diet.

You can take this product if you are pregnant or breast feeding.

This product contains methyl (E218) and propyl (E216) parahydroxybenzoates which may cause allergic reactions (possibly delayed).

**Taking Other Medicines**  
 Do not take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antihistamines, antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparation, antifungals such as ketoconazole, digoxin and

0000000

beta-blockers (for heart conditions), neuroleptics (for mental illness), thyroxine, chloroquine (for malaria) and bisphosphonates (for osteoporosis).

**How to take Gaviscon Double Action Mint**  
 Check that the cap seal is unbroken before first using this product.  
 For oral use. Shake well before use.

**Adults and children 12 years and over:**  
 Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.

**Children under 12 years:** Should only be taken on medical advice.

If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist.

If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

**After taking Gaviscon Double Action Mint:**

If symptoms persist after 7 days consult your doctor.

**Possible side effects**  
 Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness, or swelling of the face, lips, tongue or throat. If you experience these or any other side-effects stop taking the product and consult your doctor immediately. Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.

**How to store Gaviscon Double Action Mint**  
 Do not use this product after the expiry date (EXP: month/year) shown

- Use within 6 months of opening
- Keep out of the sight and reach of children

• Do not store above 30°C.  
 • Do not refrigerate or freeze

**Further information**  
 Each 10 ml of oral suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as the active ingredients. The other ingredients are sodium, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, mint flavour, sodium hydroxide and water.

This product does not contain sugar or gluten. Gaviscon Double Action Mint is available in bottles of 150ml, 300ml & 600ml.

Manufacturer and PL Holder in UK: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS.

Gaviscon is a trade mark of the RB group of companies

Text Revised: October 2020

0000000