

Quetiapine Northumbria 50 mg, 150 mg, 200 mg, 300 mg, 400 mg prolonged-release tablets

quetiapine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Quetiapine Northumbria is and what it is used for
2. What you need to know before you take Quetiapine Northumbria
3. How to take Quetiapine Northumbria
4. Possible side effects
5. How to store Quetiapine Northumbria
6. Contents of the pack and other information

1. What Quetiapine Northumbria is and what it is used for

Quetiapine Northumbria contains a substance called quetiapine. This belongs to a group of medicines called antipsychotics. Quetiapine Northumbria can be used to treat several illnesses, such as:

- Bipolar depression and major depressive episodes in major depressive disorder: where you feel sad. You may find that you feel depressed, feel guilty, lack energy, lose your appetite or can't sleep.
- Mania: where you may feel very excited, elated, agitated, enthusiastic or hyperactive or have poor judgment including being aggressive or disruptive.
- Schizophrenia: where you may hear or feel things that are not there, believe things that are not true or feel unusually suspicious, anxious, confused, guilty, tense or depressed.

When Quetiapine Northumbria is being taken to treat major depressive episodes in major depressive disorder, it will be taken in addition to another medicine being used to treat this illness.

Your doctor may continue to prescribe Quetiapine Northumbria even when you are feeling better.

2. What you need to know before you take Quetiapine Northumbria

Do not take Quetiapine Northumbria

- If you are allergic to quetiapine or any of the other ingredients of this medicine (listed in section 6)
- If you are taking any of the following medicines:
 - some medicines for HIV
 - azole medicines (for fungal infections)
 - erythromycin or clarithromycin (for infections)
 - nefazodone (for depression)

If you are not sure, talk to your doctor or pharmacist before taking Quetiapine Northumbria.

Warnings and precautions

Talk to your doctor or pharmacist before taking Quetiapine Northumbria if:

- You have depression or other conditions that are treated with antidepressants. The use of these medicines together with Quetiapine Northumbria can lead to serotonin syndrome, a potentially life-threatening condition (see "Other medicines and Quetiapine Northumbria").
- You, or someone in your family, have or have had any heart problems, for example heart rhythm problems, weakening of the heart muscle or inflammation of the heart or if you are taking any medicines that may have an impact on the way your heart beats.
- You have low blood pressure.
- You have had a stroke, especially if you are elderly.
- You have problems with your liver.
- You have ever had a fit (seizure).
- You have diabetes or have a risk of getting diabetes. If you do, your doctor may check your blood sugar levels while you are taking Quetiapine Northumbria.
- You know that you have had low levels of white blood cells in the past (which may or may not have been caused by other medicines).
- You are an elderly person with dementia (loss of brain function). If you are, Quetiapine Northumbria should not be taken because the group of medicines that Quetiapine Northumbria belongs to may increase the risk of stroke, or in some cases the risk of death, in elderly people with dementia.
- You are an elderly person with Parkinson's disease/parkinsonism.
- You or someone else in your family has a history of blood clots, as medicines like these have been associated with formation of blood clots.
- You have or have had a condition where you stop breathing for short periods during your normal nightly sleep (called "sleep apnoea") and are taking medicines that slow down the normal activity of the brain ("depressants").
- You have or have had a condition where you can't completely empty your bladder (urinary retention), have an enlarged prostate, a blockage in your intestines, or increased pressure inside your eye. These conditions are sometimes caused by medicines (called "anti-cholinergics") that affect the way nerve cells function in order to treat certain medical conditions.
- You have or have had problems with alcohol or drug abuse.

Tell your doctor immediately if you experience any of the following after taking Quetiapine Northumbria:

- A combination of fever, severe muscle stiffness, sweating or a lowered level of consciousness (a disorder called "neuroleptic malignant syndrome"). Immediate medical treatment may be needed.
- Uncontrollable movements, mainly of your face or tongue.
- Dizziness or a severe sense of feeling sleepy. This could increase the risk of accidental injury (fall) in elderly patients.
- Fits (seizures).
- A long-lasting and painful erection (Priapism).
- Have a fast and irregular heartbeat, even when you are at rest, palpitations, breathing problems, chest pain or unexplained tiredness. Your doctor will need to check your heart and if necessary, refer you to a cardiologist immediately.

These conditions can be caused by this type of medicine.

Tell your doctor as soon as possible if you have:

- A fever, flu-like symptoms, sore throat, or any other infection, as this could be a result of a very low white blood cell count, which may require Quetiapine Northumbria to be stopped and/or treatment to be given.
- Constipation along with persistent abdominal pain, or constipation which has not responded to treatment, as this may lead to a more serious blockage of the bowel.

Thoughts of suicide and worsening of your depression

If you are depressed or you may sometimes have thoughts of harming or killing yourself. These may be increased when first starting treatment, since these medicines all take time to work, usually about two weeks but sometimes longer. These thoughts may also be increased if you suddenly stop taking your medication. You may be more likely to think like this if you are a young adult. Information from clinical trials has shown an increased risk of suicidal thoughts and/or suicidal behaviour in young adults aged less than 25 years with depression.

If you have thoughts of harming or killing yourself at any time, contact your doctor or go to a hospital straight away. You may find it helpful to tell a relative or close friend that you are depressed, and ask them to read this leaflet. You might ask them to tell you if they think your depression is getting worse, or if they are worried about changes in your behaviour.

Severe cutaneous adverse reactions (SCARs)

Severe cutaneous adverse reactions (SCARs) which can be life-threatening or fatal have been reported very rarely with treatment of this medicine. These are commonly manifested by:

- Stevens-Johnson syndrome (SJS), a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals
- Toxic Epidermal Necrolysis (TEN), a more severe form causing extensive peeling of the skin
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) consists of flu-like symptoms with a rash, fever, swollen glands, and abnormal blood test results (including increased white blood cells (eosinophilia) and liver enzymes)
- Acute Generalized Exanthematous Pustulosis (AGEP), small blisters filled with pus
- Erythema Multiforme (EM), skin rash with itchy-red irregular spots

Stop using Quetiapine Northumbria if you develop these symptoms and contact your doctor or seek medical attention immediately.

Weight gain

Weight gain has been seen in patients taking Quetiapine Northumbria. You and your doctor should check your weight regularly.

Children and adolescents

Quetiapine Northumbria is not for use in children and adolescents below 18 years of age.

Other medicines and Quetiapine Northumbria

Tell your doctor if you are taking, have recently taken or might take any other medicines. Do not take Quetiapine Northumbria if you are taking any of the following medicines:

- Some medicines for HIV.
- Azole medicines (for fungal infections).
- Erythromycin or clarithromycin (for infections).
- Nefazodone (for depression).

Tell your doctor if you are taking any of the following medicines:

- Epilepsy medicines (like phenytoin or carbamazepine).
- High blood pressure medicines.
- Barbiturates (for difficulty sleeping).
- Thioridazine or Lithium (other anti-psychotic medicines).
- Medicines that have an impact on the way your heart beats, for example, medicines that can cause an imbalance in electrolytes (low levels of potassium or magnesium) such as diuretics (water pills) or certain antibiotics (medicines to treat infections).
- Medicines that can cause constipation.
- Medicines (called "anti-cholinergics") that affect the way nerve cells function in order to treat certain medical conditions.
- Anti-depressants. These medicines may interact with Quetiapine Northumbria and you may experience symptoms such as involuntary, rhythmic contractions of muscles, including the muscles that control movement of the eye, agitation, hallucinations, coma, excessive sweating, tremor, exaggeration of reflexes, increased muscle tension, body temperature above 38°C (serotonin syndrome). Contact your doctor when experiencing such symptoms.

Before you stop taking any of your medicines, please talk to your doctor first.

Quetiapine Northumbria with food, drink and alcohol

- Quetiapine Northumbria can be affected by food and you should therefore take your tablets at least one hour before a meal or prior to bedtime.
- Be careful how much alcohol you drink. This is because the combined effect of Quetiapine Northumbria and alcohol can make you sleepy.
- Do not drink grapefruit juice while you are taking Quetiapine Northumbria. It can affect the way the medicine works.

Pregnancy and breastfeeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

You should not take Quetiapine Northumbria during pregnancy unless this has been discussed with your doctor. Quetiapine Northumbria should not be taken if you are breastfeeding.

The following symptoms which can represent withdrawal may occur in newborn babies, of mothers that have used Quetiapine Northumbria in the last trimester (last three months of their pregnancy): shaking, muscle stiffness and/or weakness, sleepiness, agitation, breathing problems, and difficulty in feeding. If your baby develops any of these symptoms you may need to contact your doctor.

Driving and using machines

Your tablets may make you feel sleepy. Do not drive or use any tools or machines until you know how the tablets affect you.

Effect on Urine Drug Screens

If you are having a urine drug screen, taking Quetiapine Northumbria may cause positive results for methadone or certain drugs for depression called tricyclic antidepressants (TCAs) when some test methods are used, even though you may not be taking methadone or TCAs. If this happens, a more specific test can be performed.

Quetiapine Northumbria contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Quetiapine Northumbria

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will decide on your starting dose. The maintenance dose (daily dose) will depend on your illness and needs, but will usually be between 150 mg and 800 mg.

- You will take your tablets once a day.
- Do not split, chew or crush the tablets.
- Swallow your tablets whole with a drink of water.
- Take your tablets without food (at least one hour before a meal or at bedtime, your doctor will tell you when).
- Do not drink grapefruit juice while you are taking Quetiapine Northumbria. It can affect the way the medicine works.
- Do not stop taking your tablets even if you feel better, unless your doctor tells you to.

Liver problems

If you have liver problems your doctor may change your dose.

Elderly people

If you are elderly your doctor may change your dose.

Use in children and adolescents

Quetiapine Northumbria should not be used by children and adolescents aged under 18 years.

If you take more Quetiapine Northumbria than you should

If you take more Quetiapine Northumbria than prescribed by your doctor, you may feel sleepy, feel dizzy and experience abnormal heart beats. Contact your doctor or nearest hospital straight away. Take the Quetiapine Northumbria tablets with you.

If you forget to take Quetiapine Northumbria

If you forget to take a dose, take it as soon as you remember. If it is almost time to take the next dose, wait until then. Do not take a double dose to make up for a forgotten tablet.

If you stop taking Quetiapine Northumbria

If you suddenly stop taking Quetiapine Northumbria, you may be unable to sleep (insomnia), or you may feel sick (nausea), or you may experience headache, diarrhoea, being sick (vomiting), dizziness or irritability.

Your doctor may suggest you reduce the dose gradually before stopping treatment. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Very common (may affect more than 1 in 10 people):

- Dizziness (may lead to falls), headache, dry mouth.
- Feeling sleepy (this may go away with time, as you keep taking Quetiapine Northumbria) (may lead to falls).
- Discontinuation symptoms (symptoms which occur when you stop taking Quetiapine Northumbria) include not being able to sleep (insomnia), feeling sick (nausea), headache, diarrhoea, being sick (vomiting), dizziness and irritability. Gradual withdrawal over a period of at least 1 to 2 weeks is advisable.
- Putting on weight.
- Abnormal muscle movements. These include difficulty starting muscle movements, shaking, feeling restless or muscle stiffness without pain.
- Changes in the amount of certain fats (triglycerides and total cholesterol).

Common (may affect up to 1 in 10 people):

- Rapid heartbeat.
- Feeling like your heart is pounding, racing or has skipped beats.
- Constipation, upset stomach (indigestion).
- Feeling weak.
- Swelling of arms or legs.
- Low blood pressure when standing up. This may make you feel dizzy or faint (may lead to falls).
- Increased levels of sugar in the blood.
- Blurred vision.
- Abnormal dreams and nightmares.
- Feeling more hungry.
- Feeling irritated.
- Disturbance in speech and language.
- Thoughts of suicide and worsening of your depression.
- Shortness of breath.
- Vomiting (mainly in the elderly).
- Fever.
- Changes in the amount of thyroid hormones in your blood.
- Decreases in the number of certain types of blood cells.
- Increases in the amount of liver enzymes measured in the blood.
- Increases in the amount of the hormone prolactin in the blood. Increases in the hormone prolactin could in rare cases lead to the following:
 - Men and women to have swelling of breasts and unexpectedly produce breast milk.
 - Women to have no monthly period or irregular periods.

Uncommon (may affect up to 1 in 100 people):

- Fits or seizures.
- Allergic reactions that may include raised lumps (weals), swelling of the skin and swelling around the mouth.
- Unpleasant sensation in the legs (also called restless legs syndrome).
- Difficulty swallowing.
- Uncontrollable movements, mainly of your face or tongue.
- Sexual dysfunction.
- Diabetes.
- Change in electrical activity of the heart seen on ECG (QT prolongation).
- A slower than normal heart rate which may occur when starting treatment and which may be associated with low blood pressure and fainting.
- Difficulty in passing urine.
- Fainting (may lead to falls).
- Stuffy nose.
- Decrease in the amount of red blood cells.
- Decrease in the amount of sodium in the blood.
- Worsening of pre-existing diabetes.
- Confusion.

Rare (may affect up to 1 in 1,000 people):

- A combination of high temperature (fever), sweating, stiff muscles, feeling very drowsy or faint (a disorder called "neuroleptic malignant syndrome").
- Yellowing of the skin and eyes (jaundice).
- Inflammation of the liver (hepatitis).
- A long-lasting and painful erection (priapism).
- Swelling of breasts and unexpected production of breast milk (galactorrhoea).
- Menstrual disorder.
- Blood clots in the veins especially in the legs (symptoms include swelling, pain and redness in the leg), which may travel through blood vessels to the lungs causing chest pain and difficulty in breathing. If you notice any of these symptoms seek medical advice immediately.
- Walking, talking, eating or other activities while you are asleep.
- Body temperature decreased (hypothermia).
- Inflammation of the pancreas.
- A condition (called "metabolic syndrome") where you may have a combination of 3 or more of the following: an increase in the fat around your abdomen, a decrease in "good cholesterol" (HDL-C), an increase in a type of fat in your blood called triglycerides, high blood pressure and an increase in your blood sugar.
- Combination of fever, flu-like symptoms, sore throat, or any other infection with very low white blood cell count, a condition called agranulocytosis.
- Bowel obstruction.
- Increased blood creatine phosphokinase (a substance from muscles).

Very rare (may affect up to 1 in 10,000 people):

- Severe rash, blisters, or red patches on the skin.
- A severe allergic reaction (called anaphylaxis) which may cause difficulty in breathing or shock.
- Rapid swelling of the skin, usually around the eyes, lips and throat (angioedema).
- A serious blistering condition of the skin, mouth, eyes and genitals (Stevens-Johnson syndrome).
- Inappropriate secretion of a hormone that controls urine volume.
- Breakdown of muscle fibers and pain in muscles (rhabdomyolysis).

Not known (frequency cannot be estimated from the available data):

- Skin rash with irregular red spots (erythema multiforme) See section 2.
- Rapid appearance of areas of red skin studded with small pustules (small blisters filled with white/yellow fluid called Acute Generalised Exanthematous Pustulosis (AGEP) See section 2.
- Serious, sudden allergic reaction with symptoms such as fever and blisters on the skin and peeling of the skin (toxic epidermal necrolysis) See section 2.
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) with symptoms of flu-like symptoms with a rash, fever, swollen glands, and abnormal blood test results (including increased white blood cells (eosinophilia) and liver enzymes) See section 2.
- Symptoms of withdrawal may occur in newborn babies of mothers that have used quetiapine during their pregnancy.
- Stroke.
- Disorder of the heart muscle (cardiomyopathy).
- Inflammation of the heart muscle (myocarditis).
- Inflammation of blood vessels (Vasculitis), often with skin rash with small red or purple bumps.

The class of medicines to which Quetiapine Northumbria belongs can cause heart rhythm problems, which can be serious and in severe cases fatal.

Some side effects are only seen when a blood test is taken. These include changes in the amount of certain fats (triglycerides and total cholesterol) or sugar in the blood, changes in the amount of thyroid hormones in your blood, increased liver enzymes, decreases in the number of certain types of blood cells, decrease in the amount of red blood cells, increased blood creatine phosphokinase (a substance in the muscles), decrease in the amount of sodium in the blood, and increases in the amount of the hormone prolactin in the blood. Increases in the hormone prolactin could in rare cases lead to the following:

- Men and women to have swelling of breasts and unexpectedly produce breast milk.
- Women to have no monthly period or irregular periods.

Your doctor may ask you to have blood tests from time to time.

Additional side effects in children and adolescents

The same side effects that may occur in adults may also occur in children and adolescents.

The following side effects have been seen more often in children and adolescents or have not been seen in adults:

Very Common (may affect more than 1 in 10 people):

- Increase in the amount of a hormone called prolactin, in the blood. Increases in the hormone prolactin could in rare cases lead to the following:
 - Boys and girls to have swelling of breasts and unexpectedly produce breast milk.
 - Girls to have no monthly period or irregular periods.
- Increased appetite.
- Vomiting.
- Abnormal muscle movements. These include difficulty starting muscle movements, shaking, feeling restless or muscle stiffness without pain.
- Increase in blood pressure.

Common (may affect up to 1 in 10 people):

- Feeling weak, fainting (may lead to falls).
- Stuffy nose.
- Feeling irritated.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Quetiapine Northumbria

Keep this medicine out of the sight and reach of children.

Do not use this medicine Quetiapine Northumbria after the expiry date which is stated on the container after EXP. The expiry date refers to the last day of that month.

Quetiapine Northumbria does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Quetiapine Northumbria contains

- The active substance is quetiapine. Quetiapine Northumbria tablets contain 50 mg, 150 mg, 200 mg, 300 mg or 400 mg of quetiapine (as quetiapine fumarate).
- The other ingredients are:
 - Tablet core: lactose, Methacrylic acid – ethyl acrylate copolymer (1:1), type A crystalline maltose, magnesium stearate and talc.
 - Tablet coating: Methacrylic acid – ethyl acrylate copolymer (1:1), type A, triethyl citrate.

What Quetiapine Northumbria looks like and contents of the pack

- The 50 mg prolonged-release tablets are white to off white, round biconvex, engraved with '50' on one side, 7.1 mm in diameter and 3.2 mm in thickness.
- The 150 mg prolonged-release tablets are white to off white, oblong biconvex and engraved with '150' on one side, 13.6 mm in length, 6.6 mm in width and 4.2 mm in thickness.
- The 200 mg prolonged-release tablets are white to off white, oblong biconvex and engraved with '200' on one side, 15.2 mm in length, 7.7 mm in width and 4.8 mm in thickness.
- The 300 mg prolonged-release tablets are white to off white, oblong biconvex and engraved with '300' on one side and 18.2 mm in length, 8.2 mm in width and 5.4 mm in thickness.
- The 400 mg prolonged-release tablets are white to off white, oval biconvex and engraved with '400' on one side and 20.7 mm in length, 10.2 mm in width and 6.3 mm in thickness.
- Quetiapine Northumbria prolonged-release tablets are available as PVC/PCTFE- Aluminium foil blisters packed in a cardboard box.

Pack sizes are:

Quetiapine Northumbria 50 mg: 10, 20, 30, 50, 60 and 100 tablets.
Quetiapine Northumbria 150 mg: 10, 20, 30, 50, 60 and 100 tablets.
Quetiapine Northumbria 200 mg: 10, 20, 30, 50, 60 and 100 tablets.
Quetiapine Northumbria 300 mg: 10, 20, 30, 50, 60 and 100 tablets.
Quetiapine Northumbria 400 mg: 10, 20, 30, 50, 60 and 100 tablets.

Not all pack sizes may be marketed

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:

Fontus Health Ltd.
60 Lichfield Street
Walsall
WS4 2BX
United Kingdom

Manufacturer:

Pharmathen S.A.
6 Dervenakion Str,
Pallini
Attiki
15351
Greece

Or

Pharmathen International S.A.
Sapes Industrial Park Block 5
Rodopi
69300
Greece

Distributed by:

Northumbria Pharma Ltd.
NETpark
Thomas Wright Way
Sedgefield
Stockton-on-Tees
TS21 3FD
United Kingdom

Marketing Authorisation Numbers(s)

50mg: PL GB 42924/0024
150mg: PL GB 42924/0025
200mg: PL GB 42924/0026
300mg: PL GB 42924/0027
400mg: PL GB 42924/0028

This leaflet was last revised in 08/2025