

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Sumatriptan 100mg film-coated tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 100 mg of sumatriptan as the succinate salt.

Excipient(s) with known effect:

Lactose

For the full list of the excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film-coated (tablet).

White, oval, biconvex, 8.2x17mm coated tablets, embossed with 'SN' on one side and "100" on the other side.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Sumatriptan 100 mg Tablets are indicated for the acute relief of migraine attacks, with or without aura. Sumatriptan 100 mg Tablets should only be used where there is a clear diagnosis of migraine.

4.2 Posology and method of administration

Posology

Sumatriptan 50 mg Tablets are indicated for the acute intermittent treatment of migraine. Sumatriptan should not be used prophylactically. The recommended dose of sumatriptan should not be exceeded.

Sumatriptan is recommended as monotherapy for the acute treatment of a migraine attack and should not be given concomitantly with ergotamine or derivatives of ergotamine (including methysergide) (see section 4.3).

It is advisable that sumatriptan be given as early as possible after the onset of migraine attack but it is equally effective at whatever stage of the attack it is administered.

Adults

The recommended dose of oral sumatriptan is a single 50 mg tablet. Some patients may require 100 mg.

If a patient does not respond to the first dose of sumatriptan, a second dose should not be taken for the same attack. In these cases, the attack can be treated with paracetamol, acetylsalicylic acid, or non-steroidal anti-inflammatory drugs. Sumatriptan tablets may be taken for subsequent attacks.

If the patient has responded to the first dose but the symptoms recur a second dose may be given in the next 24 hours provided that there is a minimum interval of two hours between the two doses and no more than 300mg is taken in any 24 hour period.

Paediatric population

The efficacy and safety of sumatriptan tablets in children aged less than 10 years have not been established. No clinical data are available in this age group.

The efficacy and safety of sumatriptan tablets in children 10 to 17 years of age have not been demonstrated in the clinical studies performed in this age group. Therefore, the use of sumatriptan tablets in children 10 to 17 years of age is not recommended (see section 5.1).

Elderly (Over 65 years of age):

Experience of the use of sumatriptan in patients aged over 65 years is limited. The pharmacokinetics do not differ significantly from a younger population but until further clinical data are available, the use of Sumatriptan 50 mg Tablets in patients aged over 65 years is not recommended.

Method of administration

For oral use.

The tablets should be swallowed whole with water.

4.3 Contraindications

Hypersensitivity to sumatriptan or to any of the excipients listed in section 6.1.

Sumatriptan should not be given to patients who have had myocardial infarction or have ischaemic heart disease, coronary vasospasm (Prinzmetal's angina), peripheral vascular disease or patients who have symptoms or signs consistent with ischaemic heart disease.

Sumatriptan should not be administered to patients with a history of cerebrovascular accident (CVA) or transient ischaemic attack (TIA).

Sumatriptan should not be administered to patients with severe hepatic impairment.

The use of sumatriptan in patients with moderate and severe hypertension and mild uncontrolled hypertension is contraindicated.

The concomitant administration of ergotamine or derivatives of ergotamine (including methysergide) or any triptan/5-hydroxytryptamine₁ (5-HT₁) receptor agonist is contraindicated. (See section 4.5)

Concurrent administration of monoamine oxidase inhibitors (MAOIs) and sumatriptan is contraindicated.

Sumatriptan tablets must not be used within two weeks of discontinuation of therapy with monoamine oxidase inhibitors.

4.4 Special warnings and precautions for use

Sumatriptan tablets should only be used where there is a clear diagnosis of migraine.

Sumatriptan is not indicated for use in the management of hemiplegic, basilar or ophthalmoplegic migraine.

The recommended doses of sumatriptan should not be exceeded. As with other migraine therapies, before treating headaches in patients not previously diagnosed as migraineurs, and in migraineurs who present atypical symptoms, care should be taken to exclude other potentially serious neurological conditions (e.g. CVA, TIA). If the patient presents with atypical symptoms or if they have not received an appropriate diagnosis for sumatriptan use.

It should be noted that migraineurs may be at risk of certain cerebrovascular events (e.g. cerebrovascular accident, transient ischaemic attack).

Following administration, sumatriptan can be associated with transient symptoms including chest pain and tightness which may be intense and involve the throat (See section 4.8). Where such symptoms are thought to indicate ischaemic heart disease, no further doses of sumatriptan should be given and appropriate evaluation should be carried out.

Sumatriptan should not be given to patients with risk factors for ischaemic heart disease, including those patients who are heavy smokers or users of nicotine substitution therapies, without prior cardiovascular evaluation (See Section 4.3). Special consideration should be given to postmenopausal women and males over 40 with these risk factors. These evaluations however, may not identify every patient who has cardiac disease, and in very rare cases, serious cardiac events have occurred in patients without underlying cardiovascular disease.

Sumatriptan should be administered with caution to patients with controlled hypertension as transient increases in blood pressure and peripheral vascular resistance have been observed in a small proportion of patients (see section 4.3).

There have been rare post-marketing reports describing patients with serotonin syndrome (including altered mental status, autonomic instability and neuromuscular abnormalities) following the use of a selective serotonin reuptake inhibitor (SSRI) and sumatriptan. Serotonin syndrome has been reported following concomitant treatment with triptans and serotonin noradrenaline reuptake inhibitors (SNRIs).

If concomitant treatment with sumatriptan and an SSRI/SNRI is clinically warranted, appropriate observation of the patient is advised (see section 4.5).

If serotonin syndrome is suspected, a dose reduction or discontinuation of therapy should be considered depending on the severity of the symptoms.

Sumatriptan should be administered with caution to patients with conditions which may affect significantly the absorption, metabolism or excretion of drugs, e.g. impaired hepatic (Child Pugh grade A or B; see section 5.2) or renal function (see section 5.2). A 50mg dose should be considered in patients with hepatic impairment.

Sumatriptan should be used with caution in patients with a history of seizures or other risk factors which lower the seizure threshold, as seizures have been reported in association with sumatriptan (see section 4.8).

Patients with known hypersensitivity to sulphonamides may exhibit an allergic reaction following administration of sumatriptan. Reactions may range from cutaneous hypersensitivity to anaphylaxis. Evidence of cross-sensitivity is limited however, caution should be exercised before using sumatriptan in these patients.

Undesirable effects may be more common during concomitant use of triptans and herbal preparations containing St John's Wort (*Hypericum perforatum*).

Prolonged use of any type of painkiller for headaches can make them worse. If this situation is experienced or suspected, medical advice should be obtained, and treatment should be discontinued. The diagnosis of medication overuse headache (MOH) should be suspected in patients who have frequent or daily headaches despite (or because of) the regular use of headache medications.

Lactose

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine as it contains lactose.

Sodium

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Studies in healthy subjects show that sumatriptan does not interact with propranolol, flunarizine, pizotifen or alcohol.

There are limited data on an interaction with preparations containing ergotamine or another triptan/5-HT₁ receptor agonist. The increased risk of coronary vasospasm is a theoretical possibility and concomitant administration is contraindicated. (see section 4.3).

The period of time that should elapse between the use of sumatriptan and ergotamine-containing preparations or another triptan/5-HT₁ receptor agonist is not known. This will also depend on the doses and types of products used. The effects may be additive. It is advised to wait at least 24 hours following the use of ergotamine-containing preparations or another triptan/5-HT₁ receptor agonist before administering sumatriptan. Conversely, it is advised to wait at least 6 hours following use of sumatriptan before administering an ergotamine-containing product and at least 24 hours before administering another triptan/5-HT₁ receptor agonist.

An interaction may occur between sumatriptan and monoamine oxidase inhibitors and concomitant administration is contraindicated (see section 4.3).

There have been rare post-marketing reports describing patients with serotonin syndrome (including altered mental status, autonomic instability and neuromuscular abnormalities) following the use of SSRIs and sumatriptan. Serotonin syndrome has also been reported following concomitant treatment with triptans and SNRIs (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

Post-marketing data from the use of sumatriptan during the first trimester in over 1,000 women are available. Although these data contain insufficient information to draw definitive conclusions, they do not point to an increased risk of congenital defects. Experience with the use of sumatriptan in the second and third trimester is limited.

Evaluation of experimental animal studies does not indicate direct teratogenic effects or harmful effects on peri- and postnatal development. However, embryofetal viability might be affected in the rabbit (see section 5.3). Administration of sumatriptan should only be considered if the expected benefit to the mother is greater than any possible risk to the foetus.

Lactation

Sumatriptan is secreted into breast milk, with average relative infant doses of < 4% following administration of a single dose of sumatriptan. Infant exposure can be minimised by avoiding breast feeding for 12 hours after treatment, during which time any breast milk expressed should be discarded.

There have been reports of breast pain and/or nipple pain following sumatriptan intake in breastfeeding women (see section 4.8). The pain was usually transient and disappeared in 3 to 12 hours.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. Drowsiness may occur as a result of migraine or its treatment with sumatriptan. This may influence the ability to drive and to operate machinery.

4.8 Undesirable effects

Adverse events are listed below by system organ class and frequency. Frequencies are defined as: very common (>1/10), common (>1/100, <1/10), uncommon (>1/1000, <1/100), rare (>1/10,000, <1/1000), very rare (<1/10,000), not known (cannot be estimated from the available data). Some of the symptoms reported as undesirable effects may be associated symptoms of migraine.

Clinical Trial Data Nervous System Disorders

Common: Dizziness, drowsiness, sensory disturbance including paraesthesia and hypoaesthesia.

Vascular Disorders

Common: Transient increases in blood pressure arising soon after treatment. Flushing.

Respiratory, Thoracic and Mediastinal Disorders

Common: Dyspnoea.

Gastrointestinal Disorders

Common: Nausea and vomiting occurred in some patients but it is unclear if this is related to sumatriptan or the underlying condition.

Musculoskeletal and Connective Tissue Disorders

Common: Sensations of heaviness (usually transient and may be intense and can affect any part of the body including the chest and throat).

Myalgia.

General Disorders and Administration Site Conditions

Common: Pain, sensations of heat or cold, pressure or tightness (these events are usually transient and may be intense and can affect any part of the body including the chest and throat).

Feelings of weakness, fatigue (both events are mostly mild to moderate in intensity and transient).

Investigations

Very rare: Minor disturbances in liver function tests have occasionally been observed.

Post-Marketing Data

Immune System Disorders

Not known: Hypersensitivity reactions ranging from cutaneous hypersensitivity to anaphylaxis.

Nervous System Disorders

Not known: Seizures, although some have occurred in patients with either a history of seizures or concurrent conditions predisposing to seizures there are also reports in patients where no such predisposing factors are apparent.

Tremor, dystonia, nystagmus, scotoma.

Eye Disorders

Not known: Flickering, diplopia, reduced vision. Loss of vision including reports of permanent defects. However, visual disorders may also occur during a migraine attack itself.

Cardiac Disorders

Not known: Bradycardia, tachycardia, palpitations, cardiac arrhythmias, transient ischaemic ECG changes, coronary artery vasospasm, angina, myocardial infarction (see sections 4.3 and 4.4).

Vascular Disorders

Not known: Hypotension, Raynaud's phenomenon.

Gastrointestinal Disorders

Not known: Ischaemic colitis, diarrhoea, dysphagia.

Musculoskeletal, Connective Tissue and Bone Disorders

Not known: Neck stiffness.

Arthralgia.

Reproductive system and breast disorders

Rare: Breast pain.

General Disorders and Administration Site Conditions

Not known: Pain trauma activated, pain inflammation activated.

Psychiatric disorders

Not known: Anxiety.

Skin and subcutaneous tissue disorders

Not known: Hyperhidrosis.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in Google Play or Apple App store.

4.9 Overdose

Symptoms and signs

There have been some reports of over dosage with sumatriptan tablets. Doses in excess of 400mg orally were not associated with side effects other than those mentioned.

Treatment

If overdose occurs, the patient should be monitored for at least ten hours and standard supportive treatment applied as required.

It is unknown what effect haemodialysis or peritoneal dialysis has on the plasma concentrations of sumatriptan.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Analgesics: Selective 5-HT₁ receptor agonists.
ATC code: N02CC01

Sumatriptan has been demonstrated to be a specific and selective 5-Hydroxytryptamine₁ (5HT_{1D}) receptor agonist with no effect on other 5HT receptor (5-HT₂-5-HT₇) subtypes. The vascular 5-HT_{1D} receptor is found predominantly in cranial blood vessels and mediates vasoconstriction. In animals, sumatriptan selectively constricts the carotid arterial circulation but does not alter cerebral blood flow. The carotid arterial circulation supplies blood to the extracranial and intracranial tissues such as the meninges and dilatation of and/or oedema formation in these vessels is thought to be the underlying mechanism of migraine in man.

In addition, evidence from animal studies suggests that sumatriptan inhibits trigeminal nerve activity. Both these actions (cranial vasoconstriction and inhibition of trigeminal nerve activity) may contribute to the anti-migraine action of sumatriptan in humans.

Sumatriptan is effective in treating menstrual migraine i.e. migraine without aura that occurs between 3 days prior and up to 5 days post onset of menstruation. Sumatriptan should be taken as soon as possible in an attack.

Clinical response begins around 30 minutes following a 100mg oral dose.

Although the recommended dose of oral sumatriptan is 50mg, migraine attacks vary in severity both within and between patients. Doses of 25-100mg have shown greater efficacy than placebo in clinical trials, but 25mg is statistically significantly less effective than 50 and 100mg.

A number of placebo-controlled clinical studies assessed the safety and efficacy of oral sumatriptan in over 650 children and adolescent migraineurs aged 10-17 years. These studies failed to demonstrate relevant differences in headache relief at 2 hours between placebo and any sumatriptan dose. The undesirable effects profile of oral sumatriptan in adolescents aged 10-17 years was similar to that reported from studies in the adult population.

5.2 Pharmacokinetic properties

Following oral administration, sumatriptan is rapidly absorbed, 70% of maximum concentration occurring at 45 minutes. After 100mg dose, the maximum plasma concentration is 54ng/ml. Mean absolute oral bioavailability is 14% partly due to presystemic metabolism and partly due to incomplete absorption. The elimination phase half-life is approximately 2

hours, although there is an indication of a longer terminal phase. Plasma protein binding is low (14-21%), mean volume of distribution is 170 litres. Mean total plasma clearance is approximately 1160ml/min and the mean renal plasma clearance is approximately 260ml/min. Non-renal clearance accounts for about 80% of the total clearance. Sumatriptan is eliminated primarily by oxidative metabolism mediated by monoamine oxidase A.

Special patient populations

Hepatic Impairment

Sumatriptan pharmacokinetics after an oral dose (50 mg) and a subcutaneous dose (6 mg) were studied in 8 patients with mild to moderate hepatic impairment matched for sex, age, and weight with 8 healthy subjects.

Following an oral dose, sumatriptan plasma exposure (AUC and C_{max}) almost doubled (increased approximately 80%) in patients with mild to moderate hepatic impairment compared to the control subjects with normal hepatic function. There was no difference between the patients with hepatic impairment and control subjects after the s.c. dose. This indicates that mild to moderate hepatic impairment reduces presystemic clearance and increases the bioavailability and exposure to sumatriptan compared to healthy subjects.

Following oral administration, pre-systemic clearance is reduced in patients with mild to moderate hepatic impairment and systemic exposure is almost doubled.

The pharmacokinetics in patients with severe hepatic impairment have not been studied (see Section 4.3 Contraindications and Section 4.4 Warnings and Precautions).

The major metabolite, the indole acetic acid analogue of Sumatriptan is mainly excreted in the urine, where it is present as a free acid and the glucuronide conjugate. It has no known 5HT₁ or 5HT₂ activity. Minor metabolites have not been identified. The pharmacokinetics of oral Sumatriptan do not appear to be significantly affected by migraine attacks.

In a pilot study, no significant differences were found in the pharmacokinetic parameters between the elderly and young healthy volunteers.

5.3 Preclinical safety data

Sumatriptan was devoid of genotoxic and carcinogenic activity in *in-vitro* systems and animal studies.

In a rat fertility study oral doses of sumatriptan resulting in plasma levels approximately 200 times those seen in man after a 100 mg oral dose were associated with a reduction in the success of insemination.

This effect did not occur during a subcutaneous study where maximum plasma levels achieved approximately 150 times those in man by the oral route.

In rabbits embryoletality, without marked teratogenic defects, was seen. The relevance for humans of these findings is unknown.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Lactose monohydrate
Croscarmellose sodium
Lactose anhydrous
Microcrystalline cellulose
Magnesium stearate

Tablet coat

Lactose monohydrate
Mannitol
Titanium dioxide E171
Talc
Triacetin

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store below 30° C

6.5 Nature and contents of container

PVC/al blister or PVC-PvDC/Al blister.

Pack sizes: 2, 3, 6, 7, 12, 14 Tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Torrent Pharma (UK) Ltd.
3rd Floor, Nexus Building,
4 Gatwick Road,
Crawley,
West Sussex,
RH10 9BG,
United Kingdom

8 MARKETING AUTHORISATION NUMBER(S)

PL 36687/0366

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10/01/2025

10 DATE OF REVISION OF THE TEXT

05/02/2026

11 DOSIMETRY (IF APPLICABLE)

12 INSTRUCTIONS FOR PREPARATION OF RADIOPHARMACEUTICALS (IF APPLICABLE)