

 MICRO LABS LIMITED Bangalore, INDIA	1) Product	: Tafuprost eye drops, solution, in single-dose container-BBUK	Colours used ■ Black Colours not for Printing ■ Keylines		
	2) Strength	: 15 micrograms/ml			
	3) Component	: Leaflet			
	4) No of colours	: 01			
	5) Dimension	: 150(L) x 530(W) mm			
	6) Artwork Code	: NA			
	7) Pharma Code	: NA			
	8) Font Type and Size	: Arial Narrow, 9 pt			
	9) Technical Specification	: NA			
Code No. : - WOA ✓ WA		NA	Revision: 00		
Reason for Change		NA	Date: 24-10-2024		
	Reviewed by	Reviewed by	Reviewed by	Reviewed by	Approved by
Department	Contract Giver/ PL holder	Regulatory Affairs	Packaging Development	Head Production/ Packing (Site)	Head QA (Site)
Sign / Date					

PHARMACODE READING DIRECTION

150 mm




BROWN & BURK

Package leaflet: Information for the user
Tafuprost 15 micrograms/ml eye drops, solution, in single-dose container
 Tafuprost

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- What Tafuprost is, and what it is used for
- What you need to know before you use Tafuprost
- How to use Tafuprost
- Possible side effects
- How to store Tafuprost
- Contents of the pack and other information

1. What Tafuprost is and what it is used for

What kind of medicine is it and how does it work?
Tafuprost eye drops contain tafuprost, which belongs to a group of medicines called prostaglandin analogues. Tafuprost lowers the pressure in the eye. It is used when the pressure in the eye is too high.

What is your medicine for?
Tafuprost is used to treat a type of glaucoma called open angle glaucoma and also a condition known as ocular hypertension in adults. Both of these conditions are linked with an increase in the pressure within your eye and eventually they may affect your eyesight.

2. What you need to know before you use Tafuprost

Do not use Tafuprost:

- If you are allergic to tafuprost or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions
Talk to your doctor, pharmacist or nurse before using Tafuprost.

Please note that tafuprost may have the following effects and that some of them may be permanent:

- Tafuprost may increase the length, thickness, colour and/or number of your eyelashes and may cause unusual hair growth on your eyelids.
- Tafuprost may cause darkening of the colour of the skin around the eyes. Wipe off any excess solution from the skin. This will reduce the risk of skin darkening.
- Tafuprost may change the colour of your iris (the coloured part of your eye). If Tafuprost is used in one eye only, the colour of the treated eye may permanently become different from the colour of the other eye.
- Tafuprost may cause hair growth in areas where the solution comes repeatedly in contact with the skin surface.

Tell your doctor

- If you have kidney problems
- If you have liver problems
- If you have asthma
- If you have other eye diseases.

Children and adolescents
Tafuprost is not recommended for children and adolescents below 18 years due to a lack of data on safety and efficacy.

Other medicines and Tafuprost
Tell your doctor or pharmacist if you are taking, have recently taken or might take other medicines. If you use other medicines in the eye, leave at least 5 minutes between putting in Tafuprost and the other medication.

Pregnancy, breast-feeding and fertility
If you may become pregnant, you must use an effective method of birth control during Tafuprost therapy. Do not use tafuprost if you are pregnant. You should not use tafuprost if you are breast-feeding. Ask your doctor for advice.

Driving and using machines
Tafuprost has no or negligible influence on the ability to drive and use machines. You may find that your vision is blurred for a time just after you put Tafuprost in your eye. Do not drive or use any tools or machines until your vision is clear.

Tafuprost contains phosphates
This medicine contains approximately 0.04 mg phosphates in each drop which is equivalent to 1.2 mg/ml. If you suffer from severe damage to the clear layer at the front of the eye (the cornea), phosphates may cause in very rare cases cloudy patches on the cornea due to calcium build-up during treatment.

3. How to use Tafuprost

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 1 drop of Tafuprost in the eye or eyes, once daily in the evening. Do not instil more drops or use more often than as instructed by your doctor. This may make Tafuprost less effective.


Only use Tafuprost in both eyes if your doctor told you to.

For use as eye drops only. Do not swallow.


Instructions for use:
When you start a new pouch:
Do not use the single-dose containers if the pouch is broken. Open the pouch along the dashed line. Write down the date you opened the pouch in the space reserved for the date on the pouch.

Every time you use Tafuprost:

- Wash your hands.
- Take the strip of containers from the pouch.
- Detach one single-dose container from the strip.
- Put the remaining strip back in the pouch and fold the edge to close the pouch.
- Make sure that the solution is in the bottom part of the single-dose container.




6. To open the container, twist off the tab.




7. Tilt your head backwards.

8. Place the tip of the container close to your eye.




9. Pull the lower eyelid downwards and look up.

10. Gently squeeze the container and let one drop fall into the space between the lower eyelid and the eye.



11. Close your eye for a moment and press the inner corner of the eye with your finger for about one minute. This helps to prevent the eye drop from draining down the tear duct.

12. Wipe off any excess solution from the skin around the eye.



530 mm

Size: 150(L) x 530(W) mm
Front Side

150 mm

If a drop misses your eye, try again.

If your doctor has told you to use drops in both eyes, repeat steps 7 to 12 for your other eye.
The contents of one single-dose container are sufficient for both eyes. Discard the opened container with any remaining contents immediately after use.

If you use other medicines in the eye, leave at least 5 minutes between putting in Tafuprost and the other medication.

If you use more Tafuprost than you should, it is unlikely to cause you any serious harm. Put in your next dose at the usual time.

If the medicine is accidentally swallowed, please contact a doctor for advice.

If you forget to use Tafuprost, use a single drop as soon as you remember, and then go back to your regular routine. Do not use a double dose to make up for a forgotten dose.

Do not stop using Tafuprost without asking your doctor. If you stop using Tafuprost, the pressure in the eye will increase again. This may cause a permanent injury to your eye.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most side effects are not serious.

Common side effects
The following may affect up to 1 in 10 people:

Effects on the nervous system:

- headache.

Effects on the eye:

- itching of the eye
- irritation in the eye
- eye pain
- redness of the eye
- changes in the length, thickness and number of eyelashes
- dry eye
- foreign body sensation in the eye
- discolouration of eyelashes
- redness of the eyelids
- small spot like areas of inflammation on the surface of the eye
- sensitivity to light
- watery eyes
- blurred vision
- reduction in the eye's ability to see details
- change of colour of the iris (may be permanent).

Uncommon side effects
The following may affect up to 1 in 100 people:

Effects on the eye:

- change of colour of the skin around the eyes
- puffy eyelids
- tired eyes
- swelling of the eye's surface membranes
- eye discharge
- inflammation of the eyelids
- signs of inflammation inside the eye
- discomfort in the eye
- pigmentation of the eye's surface membranes
- follicles in the surface membranes of the eye
- allergic inflammation
- abnormal sensation in the eye.

Effects on the skin and tissue under the skin:

- unusual hair growth on eyelids.

Not known: frequency cannot be estimated from the available data

Effects on the eye:

- inflammation of the iris/lvea (middle layer of the eye)
- eyes appear sunken
- macular oedema/cystoid macular oedema (swelling of the retina within the eye leading to worsening vision).

Effects on the respiratory system:

- worsening of asthma, shortness of breath

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly (see details below). By reporting side effects, you can help provide more information on the safety of this medicine.
United Kingdom: Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

5. How to store Tafuprost

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the single-dose container, pouch and the carton after 'Exp'. The expiry date refers to the last day of that month.

Store the unopened foil pouches in a refrigerator (2°C – 8°C). Do not open the pouch until you are about to start using the eye drops as unused containers in an open pouch must be discarded 28 days after first opening the pouch.

After opening the foil pouch:

- Keep the single-dose containers in the original foil pouch.
- Do not store above 25°C
- Discard unused single-dose containers after 28 days from date of first opening of the foil pouch
- Discard an opened single-dose container with any remaining solution immediately after use.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Tafuprost contains

The active substance is tafuprost. 1 ml of solution contains 15 micrograms of tafuprost. One single-dose container (0.3 ml) contains 4.5 micrograms of tafuprost. One drop (about 30 µl) contains about 0.45 micrograms of Tafuprost.

The other ingredients are glycerol, sodium dihydrogen phosphate dihydrate, disodium edetate, polysorbate 80, water for injections, hydrochloric acid and/or sodium hydroxide (to adjust the pH value).

What Tafuprost looks like and contents of the pack
Tafuprost 15 micrograms/ml eye drops, solution is a clear, colourless liquid (solution) supplied in single-dose plastic containers, each containing 0.3 ml of solution.

Pack sizes: 30 x 0.3 ml (6 pouches with 5 single-dose containers each)
90 x 0.3 ml (18 pouches with 5 single-dose containers each)

Not all pack sizes may be marketed.


Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder
Brown & Burk UK Limited
Micro House
Bury street
Ruislip, H4A 7TL
United Kingdom

Manufacturer (Batch release)
Brown & Burk UK Limited
Micro House
Bury street
Ruislip, H4A 7TL
United Kingdom

This leaflet was last revised in {10/2024}.

BBUK Code Artwork Code



530 mm

Size: 150(L) x 530(W) mm
Back Side

PHARMACODE READING DIRECTION