

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Zolpidem 5 mg film-coated tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains: 5 mg zolpidem tartrate, equivalent to 4.02 mg zolpidem.

Excipient(s) with known effect: Each tablet contains 47.55 mg lactose monohydrate.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film-coated tablet

Red-pink, round, biconvex film-coated tablet without bisection line.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Zolpidem is indicated for the short-term treatment of insomnia **in adults** in situations where the insomnia is debilitating or is causing severe distress for the patient.

4.2 Posology and method of administration

Posology

Prior to starting treatment with zolpidem tartrate, a discussion should be held with patients to put in place a strategy for ending treatment with zolpidem tartrate in order to minimise the risk of dependence, addiction and drug withdrawal syndrome (see section 4.4). Treatment should be given for the shortest possible duration.

The treatment should be taken in a single intake and not be re-administered during the same night.

The recommended daily dose for adults is 10 mg to be taken immediately at bedtime. The lowest effective daily dose of zolpidem should be used and must not exceed 10 mg.

The duration of treatment should usually vary from a few days to two weeks with a maximum of four weeks including tapering off where clinically appropriate.

Treatment should be as short as possible. It should not exceed four weeks including the period of tapering off. In certain cases, extension beyond the maximum treatment period may be necessary; if so, extension beyond the maximum treatment period should not take place without re-evaluation of the patient's status, since the risk of abuse and dependence increases with the duration of treatment (see section 4.4).

Special Populations

Paediatric population

Zolpidem is not recommended for use in children and adolescents below 18 years of age, due to a lack of data to support use in this age group. The available evidence from placebo-controlled clinical trials is presented in section 5.1.

Elderly

Elderly or debilitated patients may be especially sensitive to the effects of zolpidem, in these subjects a 5 mg dose is recommended. These recommended doses should not exceed 10 mg in this population.

Hepatic impairment

Severe Hepatic Impairment

Zolpidem is contraindicated in patients with severe hepatic impairment as it may contribute to encephalopathy (see section 4.3).

Mild to Moderate Hepatic Impairment

As clearance and metabolism of zolpidem tartrate is reduced in hepatic impairment, dosage should begin at 5 mg in these patients with particular caution being exercised in elderly patients. In adults (under 65 years) dosage may be increased to 10 mg only where the clinical response is inadequate and the drug is well tolerated.

Method of administration

For oral use.

Zolpidem tartrate acts rapidly and therefore should be taken immediately before retiring, or in bed.

4.3 Contraindications

Zolpidem tartrate is contraindicated in patients:

- With hypersensitivity to zolpidem or any of the inactive ingredients (listed in section 6.1),
- With severe hepatic insufficiency,
- With acute and/or severe respiratory insufficiency,
- Known to have previously experienced complex sleep behaviour after taking zolpidem, see section 4.4.

4.4 Special warnings and precautions for use

Warnings:

Zolpidem should be used with caution in patients with sleep apnoea syndrome, and myasthenia gravis.

- **Respiratory insufficiency**
As hypnotics have the capacity to depress respiratory drive precautions should be observed if zolpidem is prescribed to patients with compromised respiratory function.
- **Risk from concomitant use of opioids**
Concomitant use of zolpidem and opioids may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing of sedative medicines such as benzodiazepines or related drugs such as zolpidem with opioids should be reserved for patients for whom alternative treatment options are not possible.
If a decision is made to prescribe zolpidem concomitantly with opioids, the lowest effective dose should be used, and the duration of treatment should be as short as possible (see also general dose recommendation in section 4.2). The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and their environment to be aware of these symptoms (see section 4.5).
- **Hepatic Insufficiency**
Mild to moderate Hepatic Impairment/ insufficiency- See dose recommendations (see section 4.2, 4.3 and 4.8).
- **Drug dependence, tolerance and potential for abuse**
Use of zolpidem may lead to the development of abuse and/or physical and psychological dependence.

Drug addiction comprises behavioural, cognitive and physiological phenomena that may include a strong desire to take the drug, difficulties in controlling drug use and possible tolerance or physical dependence. Physical dependence is a state that develops as a result of physiological adaptation in response to repeated drug use, which manifests as withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug. Addiction and dependence are related but distinct presentations and in discussing these themes, terminology that apportion blame to the individual should be avoided.

For all patients, prolonged use of this product may lead to drug dependence and addiction, but can occur with short-term use at recommended therapeutic doses. The risks are increased in individuals with current or past history of substance misuse disorder (including alcohol misuse) or mental health disorder (e.g., major depression).
Additional support and monitoring may be necessary when prescribing for patients at risk of drug misuse.

A comprehensive patient history should be taken to document concomitant medications, including over-the-counter medicines and medicines obtained on-line, and past and present medical and psychiatric conditions.

Tolerance

Some loss of efficacy to the hypnotic effects of sedative/hypnotic agents like zolpidem may develop after repeated use for a few weeks.

Patients may find that treatment is less effective with chronic use and express a need to increase the dose to obtain the same level of symptom control as initially experienced. Patients may also supplement their treatment with additional medications to achieve the same effect. These could be signs that the patient is developing tolerance. The risks of developing tolerance should be explained to the patient.

Overuse or misuse may result in overdose and/or death. It is important that patients only use medicines that are prescribed for them at the dose they have been prescribed and do not give this medicine to anyone else.

Patients should be closely monitored for signs of misuse, abuse, or addiction.

The clinical need for treatment with zolpidem tartrate should be reviewed regularly, with frequent assessments of patients being undertaken during the course of their treatment.

Drug withdrawal syndrome

Prior to starting treatment with zolpidem tartrate, a discussion should be held with patients to explain the risk of dependence, addiction, and drug withdrawal syndrome. A withdrawal strategy for ending treatment with zolpidem tartrate should also be put in place with the patient before starting treatment (there may be exceptions to this in specific clinical situations such as symptom management in end of life palliative care).

Drug withdrawal syndrome may occur upon abrupt cessation of therapy or dose reduction.

Once physical dependence has developed, abrupt termination of treatment will be accompanied by withdrawal symptoms. These may consist of headaches or muscle pain, extreme anxiety and tension, restlessness, confusion and irritability. In severe cases the following symptoms may occur: derealisation, depersonalisation, hyperacusis, numbness and tingling of the extremities, hypersensitivity to light, noise and physical contact, hallucinations, delirium, or epileptic seizures.

When a patient no longer requires therapy, it is advisable to taper the dose gradually to minimise symptoms of withdrawal. Tapering from a high dose may take in excess of weeks or months. Patients should be informed of this when the medication is first prescribed.

The reduction schedule for a patient should be tailored to the individual and should be modified to allow intolerable withdrawal symptoms to improve before making the next reduction. If using a published withdrawal schedule,

apply it flexibly to accommodate the person's preferences, changes to their circumstances and the response to dose reductions.

A slow stepwise rate of reduction proportionate to the existing dose should be suggested, so that decrements become smaller as the dose is lowered, unless clinical risk is such that rapid withdrawal is needed.

Precautions:

The cause of insomnia should be identified wherever possible and the underlying factors treated before a hypnotic is prescribed. The failure of insomnia to remit after a 7-14 day course of treatment may indicate the presence of a primary psychiatric or physical disorder, and the patient should be carefully re-evaluated at regular intervals.

Specific patient groups

- **Elderly**
See dose recommendations.
Due to the myorelaxant effect, there is a risk of falls and consequent injury, particularly for elderly patients when they get up at night.
- **Psychotic illness**
Hypnotics such as zolpidem are not recommended for the primary treatment of psychotic illness.
- **Paediatric Patients**
Safety and efficacy of zolpidem in patients below the age of 18 years have not been established. In an 8-week study in paediatric patients (aged 6-17 years) with insomnia associated with attention-deficit/hyperactivity disorder (ADHD), psychiatric and nervous system disorders comprised the most frequent treatment emergent adverse events observed with zolpidem versus placebo and included dizziness (23.5% vs 1.5%), headache (12.5% vs 9.2%), and hallucinations (7.4% vs 0%) (see section 4.2).
- **Use in patients with a history of drug or alcohol abuse**
Extreme caution should be exercised when prescribing for patients with a history of drug or alcohol abuse. These patients should be under careful surveillance when receiving zolpidem tartrate **or any other hypnotic**, since they are at risk of habituation and psychological dependence.
- **Next-day psychomotor impairment**
Like other sedative/hypnotic drugs, zolpidem has CNS-depressant effects. The risk of next-day psychomotor impairment, including impaired driving ability, is increased if:
 - zolpidem is taken within less than 8 hours before performing activities that require mental alertness (see section 4.7);
 - a dose higher than the recommended dose is taken;
 - zolpidem is co-administered with other CNS depressants or with other drugs that increase the blood levels of zolpidem, or with alcohol or illicit drugs (see section 4.5).Zolpidem should be taken in a single intake immediately at bedtime and not be re-administered during the same night.

- **Amnesia**
Sedative/hypnotic agents such as zolpidem may induce anterograde amnesia. The condition occurs most often several hours after ingesting the product. In order to reduce the risk, patients should ensure that they will be able to have an uninterrupted sleep of 8 hours (see section 4.8).
- **Depression and Suicidality**
Some epidemiological studies show an increased incidence of completed suicide and suicide attempt in patients with or without depression, treated with hypnotics such as zolpidem. A causal relationship has not been established. Although no clinically significant pharmacokinetic and pharmacodynamic interactions with SSRIs have been demonstrated (see section 4.5), as with other sedative/hypnotic drugs, zolpidem tartrate should be administered with caution in patients exhibiting symptoms of depression. Suicidal tendencies may be present therefore the least amount of zolpidem that is feasible should be supplied to these patients to avoid the possibility of intentional overdose by the patient.
Pre-existing depression may be unmasked during use of zolpidem. Since insomnia may be a symptom of depression, the patient should be re-evaluated if insomnia persists.

General information

- **Rebound insomnia**
A transient syndrome whereby the symptoms that led to treatment with sedative/hypnotic agents recur in an enhanced form may occur on withdrawal of hypnotic treatment. It may be accompanied by other reactions including mood changes, anxiety and restlessness. Since the risk of withdrawal phenomena or rebound has been shown to be greater after abrupt discontinuation of treatment, it is recommended that the dosage is decreased gradually where clinically appropriate.
It is important that the patient should be aware of the possibility of rebound phenomena, thereby minimising anxiety over such symptoms should they occur when the medicinal product is discontinued.
In the case of sedative/hypnotic agents with a short duration of action, withdrawal phenomena can become manifest within the dosage interval.
- **Severe Injuries**
Due to its pharmacological properties, zolpidem can cause drowsiness and a decreased level of consciousness, which may lead to falls and consequently to severe injuries, see also section 4.8.
- **Patients with Long QT syndrome**
An *in vitro* cardiac electrophysiological study showed that under experimental conditions using very high concentration and pluripotent stem cells zolpidem may reduce the hERG related potassium currents. The potential consequence in patients with congenital long QT syndrome is unknown. As a precaution, the benefit/risk ratio of zolpidem treatment in patients with known congenital long QT syndrome should be carefully considered.

- **Other psychiatric and "paradoxical" reactions**
Other psychiatric and paradoxical reactions like restlessness, insomnia exacerbated, agitation, irritability, aggression, delusion, anger, nightmares, hallucinations, abnormal behaviour, delirium, and other adverse behavioural effects are known to occur when using sedative/hypnotic agents like zolpidem. Should this occur, use of the product should be discontinued. These reactions are more likely to occur in the elderly.
- **Somnambulism and associated behaviours**
Complex sleep behaviours, including sleep walking, and other associated behaviours such as “sleep driving”, preparing and eating food, making phone calls or having sex, with amnesia for the event, have been reported in patients who had taken zolpidem and were not fully awake. These events may occur following the first or any subsequent use of zolpidem. Discontinue treatment immediately if a patient experiences a complex sleep behaviour, due to the risk to the patient and others (see section 4.3). The use of alcohol and other CNS-depressants with zolpidem appears to increase the risk of such behaviours, as does the use of zolpidem at doses exceeding the maximum recommended dose.

The medicinal product contains lactose.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

The medicinal product contains sodium

This medicinal product contains less than 1 mmol Sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

4.5 Interaction with other medicinal products and other forms of interaction

- **Alcohol**
Concomitant intake with alcohol is not recommended.
The sedative effect may be enhanced when the product is used in combination with alcohol. This affects the ability to drive or use machines.
- **CNS depressants**
Enhancement of the central depressive effect may occur in cases of concomitant use with antipsychotics (neuroleptics), hypnotics, anxiolytics/sedatives, antidepressant agents, narcotic analgesics, antiepileptic drugs, anaesthetics and sedative antihistamines.
Therefore, concomitant use of zolpidem with these drugs may increase drowsiness and next-day psychomotor impairment, including impaired driving ability (see section 4.4 and section 4.7). Also, isolated cases of visual hallucinations were reported in patients taking zolpidem with antidepressants including bupropion, desipramine, fluoxetine, sertraline and venlafaxine.
Co- administration of fluvoxamine may increase blood levels of zolpidem, concurrent use is not recommended.

In the case of narcotic analgesics enhancement of euphoria may also occur leading to an increase in psychological dependence.

- **Opioids**

The concomitant use of sedative medicines such as benzodiazepines or related drugs such as zolpidem with opioids increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. The dosage and duration of concomitant use should be limited (see section 4.4).

- **CYP450 inhibitors and inducers**

Compounds which inhibit cytochrome P450 may enhance the activity of some hypnotics like zolpidem.

Zolpidem is metabolised via several hepatic cytochrome P450 enzymes, the main enzyme being CYP3A4 with the contribution of CYP1A2. The pharmacodynamic effect of zolpidem is decreased when it is administered with a CYP3A4 inducer such as rifampicin and St. John's Wort. Co-administration of St. John's Wort may decrease blood levels of zolpidem, concurrent use is not recommended.

However, when zolpidem was administered with itraconazole (a CYP3A4 inhibitor), its pharmacokinetics and pharmacodynamics were not significantly modified. The clinical relevance of these results is unknown.

Co-administration of zolpidem with ketoconazole (200mg twice daily), a potent CYP3A4 inhibitor, prolonged zolpidem elimination half-life, increased total AUC, and decreased apparent oral clearance when compared to zolpidem plus placebo. The total AUC for zolpidem was increased modestly, when co-administered with ketoconazole, it increased by a factor of 1.83 when compared to zolpidem alone. A routine dosage adjustment of zolpidem is not considered necessary, but patients should be advised that use of zolpidem with ketoconazole may enhance the sedative effects.

Co-administration of ciprofloxacin may increase blood levels of zolpidem, concurrent use is not recommended.

- **Other drugs**

When zolpidem was administered with warfarin, digoxin, ranitidine, no significant pharmacokinetic interactions were observed.

4.6 Fertility, pregnancy and lactation

Pregnancy

The use of zolpidem is not recommended during pregnancy.

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity.

Zolpidem crosses the placenta.

A large amount of data on pregnant women (more than 1000 pregnancy outcomes) collected from cohort studies has not demonstrated evidence of the occurrence of malformations following exposure to benzodiazepines or benzodiazepine-like substances during the first

trimester of pregnancy. However, certain case-control studies reported an increased incidence of cleft lip and palate associated with use of benzodiazepines during pregnancy.

Cases of reduced foetal movement and foetal heart rate variability have been described after administration of benzodiazepines or benzodiazepine-like substances during the second and/or third trimester of pregnancy. Administration of zolpidem during the late phase of pregnancy, or during labour has been associated with effects on the neonate, such as hypothermia, hypotonia, feeding difficulties ('floppy infant syndrome'), and respiratory depression, due to the pharmacological action of the product. Cases of severe neonatal respiratory depression have been reported.

Moreover, infants born to mothers who took sedative/hypnotic agents chronically during the latter stages of pregnancy may have developed physical dependence and may be at risk of developing withdrawal symptoms in the postnatal period. Appropriate monitoring of the newborn in the postnatal period is recommended.

If zolpidem is prescribed to a woman of childbearing potential, she should be warned to contact her physician about stopping the product if she intends to become or suspects that she is pregnant.

Breastfeeding

Small quantities of zolpidem appear in breast milk. The use of zolpidem in nursing mothers is therefore, not recommended.

4.7 Effects on ability to drive and use machines

Zolpidem has major influence on the ability to drive and use machines.

Vehicle drivers and machine operators should be warned that, as with other hypnotics, there may be a possible risk of drowsiness, prolonged reaction time, dizziness, sleepiness, blurred/double vision and reduced alertness and impaired driving the morning after therapy (see section 4.8). In order to minimise this risk a resting period of at least 8 hours is recommended between taking zolpidem and driving, using machinery and working at heights. Driving ability impairment and behaviours such as 'sleep-driving' have occurred with zolpidem alone at therapeutic doses.

Furthermore, the co-administration of zolpidem with alcohol and other CNS depressants increases the risk of such behaviours (see section 4.4 and 4.5). Patients should be warned not to use alcohol or other psychoactive substances when taking zolpidem.

4.8 Undesirable effects

The following CIOMS frequency rating is used, when applicable:

Very common ($\geq 1/10$)

Common ($\geq 1/100$ to $< 1/10$)

Uncommon ($\geq 1/1\ 000$ to $< 1/100$)

Rare ($\geq 1/10\ 000$ to $< 1/1\ 000$)

Very rare ($< 1/10\ 000$)

Not known (cannot be estimated based on available data).

There is evidence of a dose-relationship for adverse effects associated with zolpidem use, particularly for certain CNS events. As recommended in section 4.2, they should, in theory, be less if zolpidem is taken immediately before retiring, or in bed. They occur most frequently in elderly patients.

Nervous system disorders

Common: somnolence, headache, dizziness, exacerbated insomnia, cognitive disorders such as memory disorders (memory impairment, amnesia, anterograde amnesia).

Uncommon: paraesthesia, tremor, disturbance in attention, speech disorder;

Rare: depressed level of consciousness.

Psychiatric disorders

Common: hallucination, agitation, nightmare, depression (see section 4.4);

Uncommon: confusional state, irritability, restlessness, aggression, somnambulism (see section 4.4), euphoric mood, complex sleep behaviour (see section 4.4);

Rare: libido disorder;

Very rare: delusion, dependence (withdrawal symptoms or rebound effects may occur after treatment discontinuation) (see section 4.4);

Not known: anger, abnormal behaviour, complex sleep behaviour, delirium (see section 4.4).

Most of these psychiatric undesirable effects are related to paradoxical reactions.

General disorders and administration site conditions

Common: fatigue;

Rare: gait disturbance, fall (predominantly in elderly patients and when zolpidem was not taken in accordance with prescribing recommendation) (see section 4.4);

Not known: drug tolerance, withdrawal syndrome. Withdrawal syndrome has been reported upon discontinuation of zolpidem tartrate (see section 4.4). Withdrawal symptoms vary and may include rebound insomnia, muscle pain, anxiety, tremor, sweating, agitation, confusion, headache, palpitations, tachycardia, delirium, nightmares, hallucinations, panic attacks, muscle aches/cramps, gastrointestinal disturbances and irritability. In severe cases the following symptoms may occur: derealisation, depersonalisation, hyperacusis, numbness and tingling of the extremities, hypersensitivity to light, noise and physical contact, hallucinations. In very rare cases, seizures may occur.

Eye disorders

Uncommon: diplopia, vision blurred;

Rare: visual impairment.

Respiratory, thoracic and mediastinal disorders

Very rare: respiratory depression (see section 4.4).

Gastro-intestinal Disorders

Common: diarrhoea, nausea, vomiting, abdominal pain.

Musculoskeletal and connective tissue disorders

Common: back pain;

Uncommon: arthralgia, myalgia, muscle spasms, neck pain, muscular weakness.

Infections and infestations

Common: upper respiratory tract infection, lower respiratory tract infection.

Skin and subcutaneous tissue disorders

Uncommon: rash, pruritus, hyperhidrosis;

Rare: urticaria.

Hepatobiliary disorders

Uncommon: liver enzymes elevated;

Rare: hepatocellular, cholestatic or mixed liver injury (see sections 4.2, 4.3 and 4.5).

Immune system disorders

Not known: angioneurotic oedema.

Metabolism and nutrition disorders

Uncommon: appetite disorder.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system listed in Appendix V***.

4.9 Overdose

Patients should be informed of the signs and symptoms of overdose and ensure that family and friends are also aware of these signs and to seek immediate medical help if they occur.

Signs and symptoms

In cases of overdose involving zolpidem alone or with other CNS-depressant agents (including alcohol), impairment of consciousness up to coma, and more severe symptomatology, including fatal outcomes have been reported.

Management

General symptomatic and supportive measures should be used. If there is no advantage in emptying the stomach, activated charcoal should be given to reduce absorption. Special attention should be paid to respiratory and cardiovascular functions in intensive care. Sedating drugs should be withheld even if excitation occurs. Use of flumazenil may be considered where serious symptoms are observed. However, flumazenil administration may contribute to the appearance of neurological symptoms (convulsions).

Zolpidem is not dialyzable.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Hypnotics and Sedatives, Benzodiazepine related drugs
ATC Code: N05CF02

Zolpidem tartrate is an imidazopyridine which selectively binds the omega-1 receptor subtype (also known as the benzodiazepine-1 subtype) which is the alpha unit of the GABA-A receptor complex. Whereas benzodiazepines non-selectively bind all three omega receptor subtypes, zolpidem tartrate preferentially binds the omega-1 subtype. The clinical relevance is not known. The modulation of the chloride anion channel via this receptor leads to the specific sedative effects demonstrated by zolpidem tartrate. These effects are reversed by the benzodiazepine antagonist flumazenil.

In animals: The selective binding of zolpidem to omega-1 receptors may explain the virtual absence at hypnotic doses of myorelaxant and anticonvulsant effects in animals which are normally exhibited by benzodiazepines which are not selective for omega-1 sites.

In humans: The preservation of deep sleep (stages 3 and 4 - slow-wave sleep) may be explained by the selective omega-1 binding by zolpidem tartrate. All identified effects of zolpidem tartrate are reversed by the benzodiazepine antagonist flumazenil.

Preliminary single dose studies did not reveal respiratory depressant effects in normal subjects or in mild or moderate COPD.

The randomized trials only showed convincing evidence of efficacy of 10 mg zolpidem. In a randomized double-blind trial in 462 non-elderly healthy volunteers with transient insomnia, zolpidem 10 mg decreased the mean time to fall asleep by 10 minutes compared to placebo, while for 5 mg zolpidem this was 3 minutes. In a randomized double-blind trial in 114 non-elderly patients with chronic insomnia, zolpidem 10 mg decreased the mean time to fall asleep by 30 minutes compared to placebo, while for 5mg zolpidem this was 15 minutes. In some patients, a lower dose of 5 mg could be effective.

Paediatric population

Safety and efficacy of zolpidem have not been established in children aged less than 18 years. A randomized placebo-controlled study in 201 children aged 6-17 years with insomnia associated with Attention Deficit Hyperactivity Disorder (ADHD) failed to demonstrate efficacy of zolpidem 0.25 mg/kg/day (with a maximum of 10 mg/day) as compared to placebo. Psychiatric and nervous system disorders comprised the most frequent treatment emergent adverse events observed with zolpidem versus placebo and included dizziness (23.5% versus 1.5%), headache (12.5% versus 9.2%), and hallucinations (7.4% versus 0%) (see section 4.2).

5.2 Pharmacokinetic properties

Zolpidem tartrate has both a rapid absorption and onset of hypnotic action. Bioavailability is 70% following oral administration and demonstrates linear kinetics in the therapeutic dose range. Peak plasma concentration is reached at between 0.5 and 3 hours.

The elimination half-life is short, with a mean of 2.4 hours (0.7-3.5) and a duration of action of up to 6 hours.

Protein binding amounts to $92.5\% \pm 0.1\%$. First pass metabolism by the liver amounts to approximately 35%. Repeated administration has been shown not to modify protein binding indicating a lack of competition between zolpidem tartrate and its metabolites for binding sites.

The distribution volume in adults is 0.54 ± 0.02 l/kg and decreases to 0.34 ± 0.05 l/kg in the very elderly.

All metabolites are pharmacologically inactive and are eliminated in the urine (56%) and in the faeces (37%).

Zolpidem tartrate has been shown in trials to be non-dialysable.

Plasma concentrations in elderly subjects and those with hepatic impairment are increased. In patients with renal insufficiency, whether dialysed or not, there is a moderate reduction in clearance.

The other pharmacokinetic parameters are unaffected.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, cancerogenic potential, toxicity to reproduction and development.

Foetal developmental retardations and foetotoxic effects in rats and rabbits were observed only at doses well above the maximum human dosage. There was no evidence of a teratogenic potential.

Fertility:

In a rat reproduction study, there was no effect on fertility in males or females after daily oral doses of 4 to 100 mg base/kg or 5 to 130 times the recommended human dose in mg/m².

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Lactose, monohydrate

Cellulose, microcrystalline

Sodium starch glycolate type A

Hypromellose

Magnesium stearate

Sepisperse dry 5212 Rose, containing:

Hypromellose

Cellulose, microcrystalline

Titanium dioxide E 171

Red Iron oxide E 172

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

5 (five) years.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions. Store in the original package to protect from light.

6.5 Nature and contents of container

The film-coated tablets are packed in PVC clear /Al press-through blister.
Pack size of 10 and 20 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

ALKALOID-INT d.o.o.
Šlandrova ulica 4
1231 Ljubljana-Črnuče
Slovenia

8 MARKETING AUTHORISATION NUMBER(S)

PL 34088/0003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 13/12/2011
Renewal of the authorisation: 14/04/2018

10 DATE OF REVISION OF THE TEXT

26/11/2025