

Package leaflet: Information for the patient

Zonisamide Mylan 100 mg hard capsules

(zonisamide)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is as above but will be referred to as Zonisamide Mylan throughout this leaflet. Zonisamide Mylan 25 mg hard capsules and Zonisamide Mylan 50 mg hard capsules are also available.

What is in this leaflet

1. What Zonisamide Mylan is and what it is used for
2. What you need to know before you take Zonisamide Mylan
3. How to take Zonisamide Mylan
4. Possible side effects
5. How to store Zonisamide Mylan
6. Contents of the pack and other information

1. What Zonisamide Mylan is and what it is used for

Zonisamide Mylan contains the active substance zonisamide, and is used as an antiepileptic medicine.

Zonisamide Mylan is used to treat seizures that affect one part of the brain (partial seizure), which may or may not be followed by a seizure affecting all of the brain (secondary generalisation).

Zonisamide Mylan may be used:

- on its own to treat seizures in adults
- with other antiepileptic medicines to treat seizures in adults, adolescents, and children aged 6 years and above.

2. What you need to know before you take Zonisamide Mylan

Do not take Zonisamide Mylan

- if you are allergic to zonisamide or any of the other ingredients of this medicine (listed in section 6).
- if you are allergic to other sulfonamide medicines. Examples include: sulfonamide antibiotics, thiazide diuretics, and sulfonylurea antidiabetes medicines.

Warnings and precautions

Zonisamide Mylan belongs to a group of medicines (sulfonamides) which can cause severe allergic reactions, severe skin rashes, and blood disorders, which very rarely can be fatal (see section 4. Possible Side Effects).

A small number of people being treated with antiepileptics such as Zonisamide Mylan have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

Serious rashes occur in association with zonisamide therapy, including cases of Stevens-Johnson syndrome.

The use of Zonisamide Mylan may lead to high levels of ammonia in the blood which could lead to a change in brain function, especially if you are also taking other medicines which can increase ammonia levels (for example valproate), have a genetic disorder causing build-up of too much ammonia in the body (urea cycle disorder), or if you have liver problems. Tell your doctor immediately if you become unusually drowsy or confused.

Talk to your doctor or pharmacist before taking Zonisamide Mylan if you:

- are younger than 12 years old, as you may be at greater risk of decreased sweating, heat stroke, pneumonia and liver problems. If you are younger than 6 years old, Zonisamide Mylan is not recommended for you
- are elderly, as your dose of Zonisamide Mylan may need adjusting, and you may be more likely to develop an allergic reaction, severe skin rash, swelling of the feet and legs, and itchiness when taking Zonisamide Mylan (see section 4 Possible Side Effects)
- suffer from liver problems, as your dose of Zonisamide Mylan may need adjusting
- have eye problems such as glaucoma
- suffer from kidney problems as your dose of Zonisamide Mylan may need adjusting
- have previously suffered from kidney stones, as you may be at increased risk of developing more kidney stones. **Reduce the risk of kidney stones by drinking sufficient water**
- live in a place or are on holiday in a place where the weather is warm. Zonisamide Mylan can make you perspire less, which can cause your body temperature to increase. **Reduce the risk of overheating by drinking sufficient water and keeping cool**
- are underweight, or have lost a lot of weight as Zonisamide Mylan can cause you to lose more weight. Tell your doctor as this may need to be monitored.
- are pregnant or could become pregnant (see section 'pregnancy, breast-feeding and fertility' for further information).

If any of these applies to you, tell your doctor before you take Zonisamide Mylan.

Children and adolescents

Talk to your doctor about the following risks:

Preventing overheating and dehydration in children

Zonisamide Mylan can cause your child to sweat less and overheat and if your child is not treated this can lead to brain damage and death. Children are most at risk especially in hot weather.

When your child is taking Zonisamide Mylan:

- keep your child cool especially in hot weather
- your child must avoid heavy exercise especially when the weather is hot
- give your child plenty of cold water to drink
- your child must not take these medicines:

carbonic anhydrase inhibitors (like topiramate and acetazolamide), and anticholinergic agents (like clomipramine, hydroxyzine, diphenhydramine, haloperidol, imipramine and oxybutynin).

If your child's skin feels very hot with little or no sweating, becomes confused, has muscle cramps, or your child's heartbeat or breathing becomes rapid:

- take your child to a cool, shaded place
- sponge your child's skin with cool (not cold) water
- give your child cold water to drink
- seek urgent medical assistance.

- Body weight: You should monitor your child's weight every month and see your doctor as soon as possible if your child is not gaining enough weight. Zonisamide Mylan is not recommended for children who are underweight or have a small appetite, and should be used with caution in those below 20 kg.
- Increased acid level in the blood and kidney stones: Reduce these risks by ensuring that your child drinks enough water and is not taking any other medicine which could cause kidney stones (see Other medicines). Your doctor will monitor your child's blood bicarbonate levels and kidneys (see also section 4).

Do not give this medicine to children below the age of 6 years because it is not known for this age group whether the potential benefits are greater than the risks.

Other medicines and Zonisamide Mylan

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

- Zonisamide Mylan should be used carefully in adults when taken with medicines that can cause kidney stones, like topiramate or acetazolamide. In children, this combination is not recommended.
- Zonisamide Mylan could possibly increase your blood levels of medicines like digoxin and quinidine, and so a reduction in their dose may be required.
- Other medicines like phenytoin, carbamazepine, phenobarbitone and rifampicin can decrease your blood levels of zonisamide, which may require an adjustment of your dose of Zonisamide Mylan.

Zonisamide Mylan with food and drink

Zonisamide Mylan can be taken with or without food.

Pregnancy, breast-feeding and fertility

- If you are pregnant, or think you may be pregnant, you must tell your doctor straight away and discuss possible risks the epilepsy medicine you are taking might pose to your unborn baby.
- If you are planning to become pregnant you should discuss your epilepsy treatment with your doctor as early as possible before you become pregnant.
- You should not stop your treatment without discussing this with your doctor. Suddenly stopping may lead to breakthrough seizures which may harm you and your unborn baby. It is important that your epilepsy remains well controlled.

Pregnancy

You must only take zonisamide during your pregnancy if your doctor tells you to. If you are a woman of childbearing age you must use adequate contraception while taking zonisamide and for one month after stopping zonisamide.

Studies have shown an increased risk of physical birth abnormalities in children of women taking epilepsy medicines during pregnancy. The risk of physical birth abnormalities may increase when more than one epilepsy medicine is used at the same time. Where possible, your doctor should consider using one epilepsy medicine to control your epilepsy. More research is needed to better understand whether taking zonisamide during pregnancy increases the risk of having a baby born with a physical birth abnormality or a learning or thinking disability. Studies have shown that babies born to mothers using zonisamide during pregnancy were smaller than expected for their age at birth, compared with babies born to mothers treated with lamotrigine monotherapy

Do not breast-feed whilst taking, or for one month after stopping Zonisamide Mylan.

There are no clinical data available on the effects of zonisamide on human fertility. Studies in animals have shown changes in fertility parameters.

Driving and using machines

Zonisamide Mylan may affect your concentration, ability to react/respond, and may make you feel sleepy, particularly at the beginning of your treatment or after your dose is increased. Be especially careful while driving or operating machinery if Zonisamide Mylan affects you in this way.

Zonisamide Mylan contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take Zonisamide Mylan

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended adult dose

When you take Zonisamide Mylan on its own:

- The starting dose is 100 mg taken once a day.
- This may be increased by up to 100 mg at intervals of two weeks.
- The recommended dose is 300 mg once a day.

When you take Zonisamide Mylan with other antiepileptic medicines:

- The starting dose is 50 mg daily taken in two equal doses of 25 mg.
- This may be increased by up to 100 mg at intervals of one to two weeks.
- The recommended daily dose is between 300 mg and 500 mg.
- Some people respond to lower doses. The dose may be increased more slowly if you experience side effects, are elderly or if you suffer from kidney or liver problems.

Use in children (aged 6 to 11 years) and adolescents (aged 12 to 17 years) weighing at least 20 kg:

- The starting dose is 1 mg per kg of body weight taken once a day.
- This may be increased by 1 mg per kg of body weight at intervals of one to two weeks.
- The recommended daily dose is 6 to 8 mg per kg for a child with a body weight of up to 55 kg or 300 to 500 mg for a child with a body weight more than 55 kg (which ever dose is lower) taken once a day.

Example: A child who weighs 25 kg should take 25 mg once a day for the first week, and then increase the daily dose by 25 mg at the start of each week until a daily dose between 150 to 200 mg is reached.

If you feel that the effect of Zonisamide Mylan is too strong or too weak, talk to your doctor or pharmacist.

- Zonisamide Mylan capsules must be swallowed whole with water.
- Do not chew the capsules.
- Zonisamide Mylan can be taken once or twice daily, as instructed by your doctor.
- If you take Zonisamide Mylan twice a day, take half the daily dose in the morning and half in the evening.

If you take more Zonisamide Mylan than you should

If you may have taken more Zonisamide Mylan than you should, tell a carer (relative or friend), your doctor or pharmacist immediately, or contact your nearest hospital casualty department, taking your medicine with you. You may become sleepy and could lose consciousness. You might also feel sick, have a sore stomach, muscle twitches, eye movement, feel faint, have a slowed heartbeat, and reduced breathing and kidney function. Do not try to drive.

If you forget to take Zonisamide Mylan

- If you forget to take a dose, don't worry: take the next dose when it is due.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Zonisamide Mylan

- Zonisamide Mylan is meant to be taken as a long-term medicine. Do not reduce your dose or stop your medicine unless your doctor tells you to.
- If your doctor advises you to stop taking Zonisamide Mylan your dose will be reduced gradually to lower the risk of more seizures.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Zonisamide Mylan belongs to a group of medicines (sulfonamides) that can cause severe allergic reactions, severe skin rashes, and blood disorders, which very rarely can be fatal.

Contact your doctor immediately if you:

- have difficulty breathing, a swollen face, lips or tongue, or a severe skin rash as these symptoms may indicate that you are having a severe allergic reaction
- have signs of overheating - high body temperature but little or no sweating, rapid heartbeat and breathing, muscle cramps, and confusion
- have thoughts of harming or killing yourself. A small number of people being treated with anti-epileptics such as Zonisamide Mylan have had thoughts of harming or killing themselves
- have pain in your muscles or a feeling of weakness, as this may be a sign of abnormal muscle breakdown which can lead to kidney problems
- get a sudden pain in your back or stomach, have pain on urinating (passing water) or notice blood in your urine, as this may be a sign of kidney stones
- develop visual problems such as eye pain or blurred vision while taken Zonisamide Mylan.

Contact your doctor as soon as possible if you:

- have an unexplained skin rash, as this could develop into a more severe skin rash or skin peeling
- feel unusually tired or feverish, have a sore throat, swollen glands, or find that you bruise more easily, as this may mean you have a blood disorder
- have signs of increased acid level in the blood- headaches, drowsiness, shortness of breath and loss of appetite. Your doctor may need to monitor or treat this.

Your doctor may decide that you should stop using Zonisamide Mylan.

The most common side effects of Zonisamide Mylan are mild. They occur during the first month of treatment and usually decrease with continued treatment. In children ages 6 – 17 years old, side effects were consistent with those described below with the following exceptions: pneumonia, dehydration, sweating decreased (common), abnormal liver enzymes (uncommon), middle ear infection, sore throat, sinus and chest infections, cough, nosebleeds, runny nose, stomach pain, vomiting, rash, eczema and fever.

Very common (may affect more than 1 in 10 people):

- agitation, irritability, confusion, depression
- poor muscle coordination, dizziness, poor memory, sleepiness, double vision
- loss of appetite, decreased blood levels of bicarbonate (a substance that prevents your blood from becoming acidic).

Common (may affect up to 1 in 10 people):

- difficulty sleeping, strange or unusual thoughts, feeling anxious or emotional
- slowed thoughts, loss of concentration, speech abnormalities, abnormal skin sensation (pins and needles), tremor, involuntary movement of the eyes
- kidney stones
- skin rashes, itching, allergic reactions, fever, tiredness, flu-like symptoms, hair loss
- ecchymosis (a small bruise caused by blood leaking from broken blood vessels in the skin)

- loss of weight, nausea, indigestion, stomach pains, diarrhoea (loose stools), constipation
- swelling of the feet and legs
- vomiting
- mood swings
- increased blood levels of creatinine (a waste product that your kidneys should normally remove)
- increased levels of liver enzymes in the blood.

Uncommon (may affect up to 1 in 100 people):

- anger, aggression, thoughts of suicide, suicide attempt
- gall bladder inflammation, gallstones
- urinary stones
- lung infection / inflammation, urinary tract infections
- low blood potassium levels, convulsions/seizures
- breathing disorders
- hallucinations
- abnormal urine tests.

Very rare (may affect up to 1 in 10,000 people):

- memory loss, coma, neuroleptic malignant syndrome (inability to move, sweating, fever, incontinence), status epilepticus (prolonged or repeated seizures)
- shortness of breath, inflammation of the lungs
- inflammations of the pancreas (severe pain in the stomach or back)
- liver problems, kidney failure
- severe rashes or skin peeling (at the same time you may feel unwell or develop a fever)
- abnormal muscle breakdown (you may feel pain or weakness in your muscles) which can lead to kidney problems
- swollen glands, blood disorders (reduction in the number of blood cells, which can make infection more likely and can make you look pale, feel tired and feverish, and bruise more easily)
- decreased sweating, overheating
- problems with your urine
- increased blood levels of creatine phosphokinase or urea which can be seen in a blood test
- abnormal results from liver function tests
- glaucoma, which is a blockage of fluid in the eye causing increased pressure in the eye. Eye pain, blurred vision or decreased vision may occur and can be signs of glaucoma.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Zonisamide Mylan

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after EXP.

The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use Zonisamide Mylan if the capsules look damaged in any way. If your capsules become discoloured or show any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Zonisamide Mylan contains:

Zonisamide Mylan 100 mg hard capsules:

The active substance is zonisamide. Each capsule contains 100 mg of zonisamide.

The other ingredients are:

- capsule contents: microcrystalline cellulose, hydrogenated vegetable oil and sodium laurilsulfate
- capsule shell: gelatin and titanium dioxide (E171)
- printing ink: shellac, black iron oxide (E172) and potassium hydroxide.

What Zonisamide Mylan looks like and contents of the pack

Zonisamide Mylan 100 mg hard capsules have a white body and white cap, marked 'Z 100' in black and contain a white/almost white powder.

Zonisamide Mylan 100 mg are available in blister packs of 56 capsules.

Manufacturer:

J. Uriach y Compania S.A.,
Av Cami Reial 51-57,
08184 Palau-Solita i Plegamans – Barcelona,
Spain.

Procured from within the EU.

Repackaged by the Product Licence Holder:
Cross Healthcare Ltd., Unit 2a Banded Industrial Estate, Throsk,
Stirling, FK7 7NP.

PLGB 20504/0544 Zonisamide Mylan 100 mg hard capsules

POM

This leaflet was last revised: 17/07/2023

**Blind or partially sighted? Is this leaflet hard to see or read?
Call 01786-817707 (Regulatory department), to obtain the leaflet in a format suitable for you.**

Package leaflet: Information for the patient Zonisamide Cross Healthcare 100 mg hard capsules (zonisamide)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is as above but will be referred to as Zonisamide Cross Healthcare throughout this leaflet.

What is in this leaflet

1. What Zonisamide Cross Healthcare is and what it is used for
2. What you need to know before you take Zonisamide Cross Healthcare
3. How to take Zonisamide Cross Healthcare
4. Possible side effects
5. How to store Zonisamide Cross Healthcare
6. Contents of the pack and other information

1. What Zonisamide Cross Healthcare is and what it is used for

Zonisamide Cross Healthcare contains the active substance zonisamide, and is used as an antiepileptic medicine.

Zonisamide Cross Healthcare is used to treat seizures that affect one part of the brain (partial seizure), which may or may not be followed by a seizure affecting all of the brain (secondary generalisation).

Zonisamide Cross Healthcare may be used:

- on its own to treat seizures in adults
- with other antiepileptic medicines to treat seizures in adults, adolescents, and children aged 6 years and above.

2. What you need to know before you take Zonisamide Cross Healthcare

Do not take Zonisamide Cross Healthcare

- if you are allergic to zonisamide or any of the other ingredients of this medicine (listed in section 6).
- if you are allergic to other sulfonamide medicines. Examples include: sulfonamide antibiotics, thiazide diuretics, and sulfonylurea antidiabetes medicines.

Warnings and precautions

Zonisamide Cross Healthcare belongs to a group of medicines (sulfonamides) which can cause severe allergic reactions, severe skin rashes, and blood disorders, which very rarely can be fatal (see section 4. Possible Side Effects).

A small number of people being treated with antiepileptics such as Zonisamide Cross Healthcare have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

Serious rashes occur in association with zonisamide therapy, including cases of Stevens-Johnson syndrome.

The use of Zonisamide Cross Healthcare may lead to high levels of ammonia in the blood which could lead to a change in brain function, especially if you are also taking other medicines which can increase ammonia levels (for example valproate), have a genetic disorder causing build-up of too much ammonia in the body (urea cycle disorder), or if you have liver problems. Tell your doctor immediately if you become unusually drowsy or confused.

Talk to your doctor or pharmacist before taking Zonisamide Cross Healthcare if you:

- are younger than 12 years old, as you may be at greater risk of decreased sweating, heat stroke, pneumonia and liver problems. If you are younger than 6 years old, Zonisamide Cross Healthcare is not recommended for you
- are elderly, as your dose of Zonisamide Cross Healthcare may need adjusting, and you may be more likely to develop an allergic reaction, severe skin rash, swelling of the feet and legs, and itchiness when taking Zonisamide Cross Healthcare (see section 4 Possible Side Effects)
- suffer from liver problems, as your dose of Zonisamide Cross Healthcare may need adjusting
- have eye problems such as glaucoma
- suffer from kidney problems as your dose of Zonisamide Cross Healthcare may need adjusting
- have previously suffered from kidney stones, as you may be at increased risk of developing more kidney stones. **Reduce the risk of kidney stones by drinking sufficient water**
- live in a place or are on holiday in a place where the weather is warm. Zonisamide Cross Healthcare can make you perspire less, which can cause your body temperature to increase. **Reduce the risk of overheating by drinking sufficient water and keeping cool**
- are underweight, or have lost a lot of weight as Zonisamide Cross Healthcare can cause you to lose more weight. Tell your doctor as this may need to be monitored.
- are pregnant or could become pregnant (see section 'pregnancy, breast-feeding and fertility' for further information).

If any of these applies to you, tell your doctor before you take Zonisamide Cross Healthcare.

Children and adolescents

Talk to your doctor about the following risks:

Preventing overheating and dehydration in children

Zonisamide Cross Healthcare can cause your child to sweat less and overheat and if your child is not treated this can lead to brain damage and death. Children are most at risk especially in hot weather.

When your child is taking Zonisamide Cross Healthcare:

- keep your child cool especially in hot weather
- your child must avoid heavy exercise especially when the weather is hot
- give your child plenty of cold water to drink
- your child must not take these medicines:

carbonic anhydrase inhibitors (like topiramate and acetazolamide), and anticholinergic agents (like clomipramine, hydroxyzine, diphenhydramine, haloperidol, imipramine and oxybutynin).

If your child's skin feels very hot with little or no sweating, becomes confused, has muscle cramps, or your child's heartbeat or breathing becomes rapid:

- take your child to a cool, shaded place
- sponge your child's skin with cool (not cold) water
- give your child cold water to drink
- seek urgent medical assistance.

- Body weight: You should monitor your child's weight every month and see your doctor as soon as possible if your child is not gaining enough weight. Zonisamide Cross Healthcare is not recommended for children who are underweight or have a small appetite, and should be used with caution in those below 20 kg.
- Increased acid level in the blood and kidney stones: Reduce these risks by ensuring that your child drinks enough water and is not taking any other medicine which could cause kidney stones (see Other medicines). Your doctor will monitor your child's blood bicarbonate levels and kidneys (see also section 4).

Do not give this medicine to children below the age of 6 years because it is not known for this age group whether the potential benefits are greater than the risks.

Other medicines and Zonisamide Cross Healthcare

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

- Zonisamide Cross Healthcare should be used carefully in adults when taken with medicines that can cause kidney stones, like topiramate or acetazolamide. In children, this combination is not recommended.
- Zonisamide Cross Healthcare could possibly increase your blood levels of medicines like digoxin and quinidine, and so a reduction in their dose may be required.
- Other medicines like phenytoin, carbamazepine, phenobarbitone and rifampicin can decrease your blood levels of zonisamide, which may require an adjustment of your dose of Zonisamide Cross Healthcare.

Zonisamide Cross Healthcare with food and drink

Zonisamide Cross Healthcare can be taken with or without food.

Pregnancy, breast-feeding and fertility

- If you are pregnant, or think you may be pregnant, you must tell your doctor straight away and discuss possible risks the epilepsy medicine you are taking might pose to your unborn baby.
- If you are planning to become pregnant you should discuss your epilepsy treatment with your doctor as early as possible before you become pregnant.
- You should not stop your treatment without discussing this with your doctor. Suddenly stopping may lead to breakthrough seizures which may harm you and your unborn baby. It is important that your epilepsy remains well controlled.

Pregnancy

You must only take zonisamide during your pregnancy if your doctor tells you to. If you are a woman of childbearing age you must use adequate contraception while taking zonisamide and for one month after stopping zonisamide.

Studies have shown an increased risk of physical birth abnormalities in children of women taking epilepsy medicines during pregnancy. The risk of physical birth abnormalities may increase when more than one epilepsy medicine is used at the same time. Where possible, your doctor should consider using one epilepsy medicine to control your epilepsy. More research is needed to better understand whether taking zonisamide during pregnancy increases the risk of having a baby born with a physical birth abnormality or a learning or thinking disability. Studies have shown that babies born to mothers using zonisamide during pregnancy were smaller than expected for their age at birth, compared with babies born to mothers treated with lamotrigine monotherapy

Do not breast-feed whilst taking, or for one month after stopping Zonisamide Cross Healthcare.

There are no clinical data available on the effects of zonisamide on human fertility. Studies in animals have shown changes in fertility parameters.

Driving and using machines

Zonisamide Cross Healthcare may affect your concentration, ability to react/respond, and may make you feel sleepy, particularly at the beginning of your treatment or after your dose is increased. Be especially careful while driving or operating machinery if Zonisamide Cross Healthcare affects you in this way.

Zonisamide Cross Healthcare contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take Zonisamide Cross Healthcare

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended adult dose

When you take Zonisamide Cross Healthcare on its own:

- The starting dose is 100 mg taken once a day.
- This may be increased by up to 100 mg at intervals of two weeks.
- The recommended dose is 300 mg once a day.

When you take Zonisamide Cross Healthcare with other antiepileptic medicines:

- The starting dose is 50 mg daily taken in two equal doses of 25 mg.
- This may be increased by up to 100 mg at intervals of one to two weeks.
- The recommended daily dose is between 300 mg and 500 mg.
- Some people respond to lower doses. The dose may be increased more slowly if you experience side effects, are elderly or if you suffer from kidney or liver problems.

Use in children (aged 6 to 11 years) and adolescents (aged 12 to 17 years) weighing at least 20 kg:

- The starting dose is 1 mg per kg of body weight taken once a day.
- This may be increased by 1 mg per kg of body weight at intervals of one to two weeks.
- The recommended daily dose is 6 to 8 mg per kg for a child with a body weight of up to 55 kg or 300 to 500 mg for a child with a body weight more than 55 kg (which ever dose is lower) taken once a day.

Example: A child who weighs 25 kg should take 25 mg once a day for the first week, and then increase the daily dose by 25 mg at the start of each week until a daily dose between 150 to 200 mg is reached.

If you feel that the effect of Zonisamide Cross Healthcare is too strong or too weak, talk to your doctor or pharmacist.

- Zonisamide Cross Healthcare capsules must be swallowed whole with water.
- Do not chew the capsules.
- Zonisamide Cross Healthcare can be taken once or twice daily, as instructed by your doctor.
- If you take Zonisamide Cross Healthcare twice a day, take half the daily dose in the morning and half in the evening.

If you take more Zonisamide Cross Healthcare than you should

If you may have taken more Zonisamide Cross Healthcare than you should, tell a carer (relative or friend), your doctor or pharmacist immediately, or contact your nearest hospital casualty department, taking your medicine with you. You may become sleepy and could lose consciousness. You might also feel sick, have a sore stomach, muscle twitches, eye movement, feel faint, have a slowed heartbeat, and reduced breathing and kidney function. Do not try to drive.

If you forget to take Zonisamide Cross Healthcare

- If you forget to take a dose, don't worry: take the next dose when it is due.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Zonisamide Cross Healthcare

- Zonisamide Cross Healthcare is meant to be taken as a long-term medicine. Do not reduce your dose or stop your medicine unless your doctor tells you to.
- If your doctor advises you to stop taking Zonisamide Cross Healthcare your dose will be reduced gradually to lower the risk of more seizures.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Zonisamide Cross Healthcare belongs to a group of medicines (sulfonamides) that can cause severe allergic reactions, severe skin rashes, and blood disorders, which very rarely can be fatal.

Contact your doctor immediately if you:

- have difficulty breathing, a swollen face, lips or tongue, or a severe skin rash as these symptoms may indicate that you are having a severe allergic reaction
- have signs of overheating - high body temperature but little or no sweating, rapid heartbeat and breathing, muscle cramps, and confusion
- have thoughts of harming or killing yourself. A small number of people being treated with anti-epileptics such as Zonisamide Cross Healthcare have had thoughts of harming or killing themselves
- have pain in your muscles or a feeling of weakness, as this may be a sign of abnormal muscle breakdown which can lead to kidney problems
- get a sudden pain in your back or stomach, have pain on urinating (passing water) or notice blood in your urine, as this may be a sign of kidney stones
- develop visual problems such as eye pain or blurred vision while taken Zonisamide Cross Healthcare.

Contact your doctor as soon as possible if you:

- have an unexplained skin rash, as this could develop into a more severe skin rash or skin peeling
- feel unusually tired or feverish, have a sore throat, swollen glands, or find that you bruise more easily, as this may mean you have a blood disorder
- have signs of increased acid level in the blood- headaches, drowsiness, shortness of breath and loss of appetite. Your doctor may need to monitor or treat this.

Your doctor may decide that you should stop using Zonisamide Cross Healthcare.

The most common side effects of Zonisamide Cross Healthcare are mild. They occur during the first month of treatment and usually decrease with continued treatment. In children ages 6 – 17 years old, side effects were consistent with those described below with the following exceptions: pneumonia, dehydration, sweating decreased (common), abnormal liver enzymes (uncommon), middle ear infection, sore throat, sinus and chest infections, cough, nosebleeds, runny nose, stomach pain, vomiting, rash, eczema and fever.

Very common (may affect more than 1 in 10 people):

- agitation, irritability, confusion, depression
- poor muscle coordination, dizziness, poor memory, sleepiness, double vision
- loss of appetite, decreased blood levels of bicarbonate (a substance that prevents your blood from becoming acidic).

Common (may affect up to 1 in 10 people):

- difficulty sleeping, strange or unusual thoughts, feeling anxious or emotional
- slowed thoughts, loss of concentration, speech abnormalities, abnormal skin sensation (pins and needles), tremor, involuntary movement of the eyes
- kidney stones

- skin rashes, itching, allergic reactions, fever, tiredness, flu-like symptoms, hair loss
- ecchymosis (a small bruise caused by blood leaking from broken blood vessels in the skin)
- loss of weight, nausea, indigestion, stomach pains, diarrhoea (loose stools), constipation
- swelling of the feet and legs
- vomiting
- mood swings
- increased blood levels of creatinine (a waste product that your kidneys should normally remove)
- increased levels of liver enzymes in the blood.

Uncommon (may affect up to 1 in 100 people):

- anger, aggression, thoughts of suicide, suicide attempt
- gall bladder inflammation, gallstones
- urinary stones
- lung infection / inflammation, urinary tract infections
- low blood potassium levels, convulsions/seizures
- breathing disorders
- hallucinations
- abnormal urine tests.

Very rare (may affect up to 1 in 10,000 people):

- memory loss, coma, neuroleptic malignant syndrome (inability to move, sweating, fever, incontinence), status epilepticus (prolonged or repeated seizures)
- shortness of breath, inflammation of the lungs
- inflammations of the pancreas (severe pain in the stomach or back)
- liver problems, kidney failure
- severe rashes or skin peeling (at the same time you may feel unwell or develop a fever)
- abnormal muscle breakdown (you may feel pain or weakness in your muscles) which can lead to kidney problems
- swollen glands, blood disorders (reduction in the number of blood cells, which can make infection more likely and can make you look pale, feel tired and feverish, and bruise more easily)
- decreased sweating, overheating
- problems with your urine
- increased blood levels of creatine phosphokinase or urea which can be seen in a blood test
- abnormal results from liver function tests
- glaucoma, which is a blockage of fluid in the eye causing increased pressure in the eye. Eye pain, blurred vision or decreased vision may occur and can be signs of glaucoma.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Zonisamide Cross Healthcare

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after EXP.

The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use Zonisamide Cross Healthcare if the capsules look damaged in any way. If your capsules become discoloured or show any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Zonisamide Cross Healthcare contains:

Zonisamide Cross Healthcare 100 mg hard capsules:
The active substance is zonisamide. Each capsule contains 100 mg of zonisamide.

The other ingredients are:

- capsule contents: microcrystalline cellulose, hydrogenated vegetable oil and sodium laurilsulfate
- capsule shell: gelatin and titanium dioxide (E171)
- printing ink: shellac, black iron oxide (E172) and potassium hydroxide.

What Zonisamide Cross Healthcare looks like and contents of the pack

Zonisamide Cross Healthcare 100 mg hard capsules have a white body and white cap, marked 'Z 100' in black and contain a white/almost white powder.

Zonisamide Cross Healthcare 100 mg are available in blister packs of 56 capsules.

Manufacturer:

J. Uriach y Compania S.A.,
Av Cami Reial 51-57,
08184 Palau-Solita i Plegamans – Barcelona,
Spain.

Procured from within the EU.

Repackaged by the Product Licence Holder:
Cross Healthcare Ltd., Unit 2a Bandeath Industrial Estate, Throsk,
Stirling, FK7 7NP.

PLGB 20504/0544 Zonisamide Cross Healthcare 100 mg hard capsules

This leaflet was last revised: 17/07/2023

POM

**Blind or partially sighted? Is this leaflet hard to see or read?
Call 01786-817707 (Regulatory department), to obtain the leaflet in a format suitable for you.**