

Package leaflet: Information for the user

Voriconazole Accordpharma 200mg Powder for Solution for Infusion

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.
- The full name of this medicine is Voriconazole Accordpharma 200mg Powder for Solution for Infusion but within the leaflet it will be referred to as Voriconazole.

What is in this leaflet

1. What Voriconazole is and what it is used for
2. What you need to know before you use Voriconazole
3. How to use Voriconazole
4. Possible side effects
5. How to store Voriconazole
6. Contents of the pack and other information

1. What Voriconazole is and what it is used for

Voriconazole Accordpharma 200mg Powder for Solution for Infusion contains the active substance voriconazole. Voriconazole is an antifungal medicine. It works by killing or stopping the growth of the fungi that cause infections.

It is used for the treatment of patients (adults and children over the age of 2) with:

- invasive aspergillosis (a type of fungal infection due to *Aspergillus sp*)
- candidaemia (another type of fungal infection due to *Candida sp*) in non-neutropenic patients (patients without abnormally low white blood cells count)
- serious invasive *Candida sp.* infections when the fungus is resistant to fluconazole (another antifungal medicine)
- serious fungal infections caused by *Scedosporium sp.* or *Fusarium sp.* (two different species of fungi).

Voriconazole is intended for patients with worsening, possibly life-threatening, fungal infections.

Prevention of fungal infections in high risk bone marrow transplant recipients.

This product should only be used under the supervision of a doctor.

2. What you need to know before you use Voriconazole

Do not use Voriconazole

- if you are allergic to the active ingredient voriconazole, or to hydroxypropylbetadex or any of the other ingredients of this medicine (listed in section 6).

It is very important that you inform your doctor or pharmacist if you are taking or have taken any other medicines, even those that are obtained without a prescription, or herbal medicines.

The medicines in the following list must not be taken during your course of Voriconazole treatment:

- Terfenadine (used for allergy)
- Astemizole (used for allergy)
- Cisapride (used for stomach problems)
- Pimozide (used for treating mental illness)
- Quinidine (used for irregular heart beat)
- Ivabradine (used for symptoms of chronic heart failure)
- Rifampicin (used for treating tuberculosis)
- Efavirenz (used for treating HIV) in doses of 400mg and above once daily
- Carbamazepine (used to treat seizures)
- Phenobarbital (used for severe insomnia and seizures)
- Ergot alkaloids (e.g., ergotamine, dihydroergotamine; used for migraine)
- Sirolimus (used in transplant patients)
- Ritonavir (used for treating HIV) in doses of 400mg and more twice daily
- St. John's Wort (herbal supplement)
- Naloxegol (used to treat constipation specifically caused by pain medicines, called opioids, (e.g., morphine, oxycodone, fentanyl, tramadol, codeine))
- Tolvaptan (used to treat hyponatremia (low levels of sodium in your blood) or to slow kidney function decline in patients with polycystic kidney disease)
- Lurasidone (used to treat depression)
- Venetoclax (used to treat patients with chronic lymphocytic leukaemia-CLL).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before using Voriconazole

- if you have had an allergic reaction to other azoles.
- if you are suffering from, or have ever suffered from liver disease. If you have liver disease, your doctor may prescribe a lower dose of Voriconazole. Your doctor should also monitor your liver function while you are being treated with Voriconazole by doing blood tests.
- if you are known to have cardiomyopathy, irregular heartbeat, slow heart rate or an abnormality of electrocardiogram (ECG) called 'long QTc syndrome'.

You should avoid any sunlight and sun exposure while being treated. It is important to cover sun exposed areas of skin and use sunscreen with high sun protection factor (SPF), as an increased sensitivity of skin to the sun's UV rays can occur. This may be further increased by other medicines that sensitise the skin to sunlight, like methotrexate. These precautions are also applicable to children.

While being treated with Voriconazole:

- tell your doctor immediately if you develop
 - sunburn
 - severe skin rash or blisters
 - bone pain.

If you develop skin disorders as described above, your doctor may refer you to a dermatologist, who after consultation may decide that it is important for you to be seen on a regular basis. There is a small chance that skin cancer could develop with long-term use of Voriconazole.

If you develop signs of 'adrenal insufficiency' where the adrenal glands do not produce adequate amounts of certain steroid hormones such as cortisol which may lead to symptoms such as: chronic, or long lasting fatigue, muscle weakness, loss of appetite, weight loss, abdominal pain, please tell your doctor.

If you develop signs of 'Cushing's syndrome' where the body produces too much of the hormone cortisol which may lead to symptoms such as: weight gain, fatty hump between the shoulders, a rounded face, darkening of the skin on the stomach, thighs breasts, and arms, thinning skin, bruising easily, high blood sugar, excessive hair growth, excessive sweating, please tell your doctor.

Your doctor should monitor the function of your liver and kidney by doing blood tests.

Children and adolescents

Voriconazole should not be given to children younger than 2 years of age.

Other medicines and Voriconazole

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines, including those that are obtained without a prescription.

Some medicines, when taken at the same time as Voriconazole, may affect the way Voriconazole works or Voriconazole may affect the way they work.

Tell your doctor if you are taking the following medicine, as treatment with Voriconazole at the same time should be avoided if possible:

- Ritonavir (used for treating HIV) in doses of 100mg twice daily
- Glasdegib (used for treating cancer) – if you need to use both drugs your doctor will monitor your heart rhythm frequently.

Tell your doctor if you are taking either of the following medicines, as treatment with Voriconazole at the same time should be avoided if possible and a dose adjustment of voriconazole may be required:

- Rifabutin (used for treating tuberculosis). If you are already being treated with rifabutin your blood counts and side effects to rifabutin will need to be monitored.
- Phenytoin (used to treat epilepsy). If you are already being treated with phenytoin your blood concentration of phenytoin will need to be monitored during your treatment with Voriconazole and your dose may be adjusted.

Tell your doctor if you are taking any of the following medicines, as a dose adjustment or monitoring may be required to check that the medicines and/ or Voriconazole are still having the desired effect:

- Warfarin and other anticoagulants (e.g., phenprocoumon, acenocoumarol; used to slow down clotting of the blood)
- Ciclosporin (used in transplant patients)
- Tacrolimus (used in transplant patients)
- Sulfonylureas (e.g. tolbutamide, glipizide, and glyburide) (used for diabetes)
- Statins (e.g. atorvastatin, simvastatin) (used for lowering cholesterol)
- Benzodiazepines (e.g. midazolam, triazolam) (used for severe insomnia and stress)
- Omeprazole (used for treating ulcers)
- Oral contraceptives (if you take Voriconazole whilst using oral contraceptives, you may get side effects such as nausea and menstrual disorders)
- Vinca alkaloids (e.g. vincristine and vinblastine) (used in treating cancer)
- Tyrosine kinase inhibitors (e.g., axitinib, bosutinib, cabozantinib, ceritinib, cobimetinib, dabrafenib, dasatinib, nilotinib, sunitinib, ibrutinib, ribociclib) (used for treating cancer)
- Tretinoin (used to treat leukaemia)
- Indinavir and other HIV protease inhibitors (used for treating HIV)
- Non-nucleoside reverse transcriptase inhibitors (e.g. efavirenz, delavirdine, nevirapine) (used for treating HIV) (some doses of efavirenz can NOT be taken at the same time as Voriconazole)
- Methadone (used to treat heroin addiction)
- Alfentanil and fentanyl and other short-acting opiates such as sufentanil (painkillers used for surgical procedures)
- Oxycodone and other long-acting opiates such as hydrocodone (used for moderate to severe pain)
- Non-steroidal anti-inflammatory drugs (e.g. ibuprofen, diclofenac) (used for treating pain and inflammation)
- Fluconazole (used for fungal infections)
- Everolimus (used for treating advanced kidney cancer and in transplant patients)
- Letermovir (used for preventing cytomegalovirus (CMV) disease after bone marrow transplant)

- Ivacaftor: used to treat cystic fibrosis
- Flucloxacillin (antibiotic used against bacterial infection).

Pregnancy and breast-feeding

Voriconazole must not be used during pregnancy, unless indicated by your doctor. Effective contraception must be used in women of childbearing potential. Contact your doctor immediately if you become pregnant while being treated with Voriconazole.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Voriconazole may cause blurring of vision or uncomfortable sensitivity to light. While affected, do not drive or operate any tools or machines. Tell your doctor if you experience this.

Voriconazole contains sodium

This medicinal product contains less than 1mmol sodium (23mg) per vial, that is to say essentially 'sodium-free'.

Voriconazole contains cyclodextrins

This medicine contains 2060.0 mg cyclodextrins in each vial which is equivalent to 103 mg/ml when reconstituted in 20 ml.

3. How to use Voriconazole

Always use this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Your doctor will determine your dose depending on your weight and the type of infection you have.

Your doctor may change your dose depending on your condition.

The recommended dose for adults (including elderly patients) is as follows:

	Intravenous
Dose for the first 24 hours (Loading Dose)	6mg/kg every 12 hours for the first 24 hours
Dose after the first 24 hours (Maintenance Dose)	4mg/kg twice a day

Depending on your response to treatment, your doctor may decrease the dose to 3mg/kg twice daily.

The doctor may decide to decrease the dose if you have mild to moderate cirrhosis.

Use in children and adolescents

The recommended dose for children and teenagers is as follows:

	Intravenous	
	Children aged 2 to less than 12 years and teenagers aged 12 to 14 years weighing less than 50kg	Teenagers aged 12 to 14 years weighing 50kg or more; and all teenagers older than 14
Dose for the first 24 hours (Loading Dose)	9mg/kg every 12 hours for the first 24 hours	6mg/kg every 12 hours for the first 24 hours
Dose after the first 24 hours	8mg/kg twice a day	4mg/kg twice a day

(Maintenance Dose)		
--------------------	--	--

Depending on your response to treatment, your doctor may increase or decrease the daily dose.

Voriconazole Powder for Solution for Infusion will be reconstituted and further diluted to the correct concentration by your hospital pharmacist or nurse. (Please refer to the end of this leaflet for further information).

This will be given to you by intravenous infusion (into a vein) at a maximum rate of 3mg/kg per hour over 1 to 3 hours.

If you or your child are taking Voriconazole for prevention of fungal infections, your doctor may stop giving Voriconazole if you or your child develop treatment related side effects.

If a dose of Voriconazole has been forgotten

As you will be given this medicine under close medical supervision, it is unlikely that a dose would be missed. However, tell your doctor or pharmacist if you think that a dose has been forgotten.

If you stop using Voriconazole

Voriconazole treatment will continue for as long as your doctor advises; however, duration of treatment with Voriconazole Powder for Solution for Infusion should be no more than 6 months.

Patients with a weakened immune system or those with difficult infections may require long-term treatment to prevent the infection from returning. You may be switched from the intravenous infusion to tablets once your condition improves.

When Voriconazole treatment is stopped by your doctor, you should not experience any effects.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If any side effects occur, most are likely to be minor and temporary. However, some may be serious and need medical attention.

Serious side effects – Stop using Voriconazole and see a doctor immediately

- Rash
- Jaundice; Changes in blood tests of liver function
- Pancreatitis.

Other side effects

Very common (may affect more than 1 in 10 people):

- Visual impairment (change in vision including blurred vision, visual colour alterations, abnormal intolerance to visual perception of light, colour blindness, eye disorder, halo vision, night blindness, swinging vision, seeing sparks, visual aura, visual acuity reduced, visual brightness, loss of part of the usual field of vision, spots before the eyes)
- Fever
- Rash
- Nausea, vomiting, diarrhoea
- Headache
- Swelling of the extremities
- Stomach pains
- Breathing difficulties

- Elevated liver enzymes.

Common (may affect up to 1 in 10 people):

- Inflammation of the sinuses, inflammation of the gums, chills, weakness
- Low numbers of some types, including severe, of red (sometimes immune-related) and/or white blood cells (sometimes with fever), low numbers of cells called platelets that help the blood to clot
- Low blood sugar, low blood potassium, low sodium in the blood
- Anxiety, depression, confusion, agitation, inability to sleep, hallucinations
- Seizures, tremors or uncontrolled muscle movements, tingling or abnormal skin sensations, increase in muscle tone, sleepiness, dizziness
- Bleeding in the eye
- Heart rhythm problems including very fast heartbeat, very slow heartbeat, fainting
- Low blood pressure, inflammation of a vein (which may be associated with the formation of a blood clot)
- Acute breathing difficulty, chest pain, swelling of the face (mouth, lips and around eyes), fluid accumulation in the lungs
- Constipation, indigestion, inflammation of the lips
- Jaundice, inflammation of the liver and liver injury
- Skin rashes which may lead to severe blistering and peeling of the skin characterised by a flat, red area on the skin that is covered with small confluent bumps, redness of the skin
- Itchiness
- Hair loss
- Back pain
- Kidney failure, blood in the urine, changes in kidney function tests
- Sunburn or severe skin reaction following exposure to light or sun
- Skin cancer.

Uncommon (may affect up to 1 in 100 people):

- Flu-like symptoms, irritation and inflammation of the gastrointestinal tract, inflammation of the gastrointestinal tract causing antibiotic associated diarrhoea, inflammation of the lymphatic vessels
- Inflammation of the thin tissue that lines the inner wall of the abdomen and covers the abdominal organ
- Enlarged lymph glands (sometimes painful), failure of blood marrow, increased eosinophil
- Depressed function of the adrenal gland, underactive thyroid gland
- Abnormal brain function, Parkinson-like symptoms, nerve injury resulting in numbness, pain, tingling or burning in the hands or feet
- Problems with balance or coordination
- Swelling of the brain
- Double vision, serious conditions of the eye including: pain and inflammation of the eyes and eyelids, abnormal eye movement, damage to the optic nerve resulting in vision impairment, optic disc swelling
- Decreased sensitivity to touch
- Abnormal sense of taste
- Hearing difficulties, ringing in the ears, vertigo
- Inflammation of certain internal organs- pancreas and duodenum, swelling and inflammation of the tongue
- Enlarged liver, liver failure, gallbladder disease, gallstones
- Joint inflammation, inflammation of the veins under the skin (which may be associated with the formation of a blood clot)
- Inflammation of the kidney, proteins in the urine, damage to the kidney
- Very fast heart rate or skipped heartbeats, sometimes with erratic electrical impulse
- Abnormal electrocardiogram (ECG)
- Blood cholesterol increased, blood urea increased

- Allergic skin reactions (sometimes severe), including life-threatening skin condition that causes painful blisters and sores of the skin and mucous membranes, especially in the mouth, inflammation of the skin, hives, skin redness and irritation, red or purple discoloration of the skin which may be caused by low platelet count, eczema
- Infusion site reaction
- Allergic reaction or exaggerated immune response
- Inflammation of the tissue surrounding the bone.

Rare (may affect up to 1 in 1,000 people):

- Overactive thyroid gland
- Deterioration of brain function that is a serious complication of liver disease
- Loss of most fibers in the optic nerve, clouding of the cornea, involuntary movement of the eye
- Bullous photosensitivity
- A disorder in which the body's immune system attacks part of the peripheral nervous system
- Heart rhythm or conduction problems (sometimes life threatening)
- Life threatening allergic reaction
- Disorder of blood clotting system
- Allergic skin reactions (sometimes severe), including rapid swelling (oedema) of the dermis, subcutaneous tissue, mucosa and submucosal tissues, itchy or sore patches of thick, red skin with silvery scales of skin, irritation of the skin and mucous membranes, life-threatening skin condition that causes large portions of the epidermis, the skin's outermost layer, to detach from the layers of skin below
- Small dry scaly skin patches, sometimes thick with spikes or 'horns'.

Side effects with frequency not known (frequency cannot be estimated from the available data):

- Freckles and pigmented spots.

Other significant side effects whose frequency is not known, but should be reported to your doctor immediately:

- Red, scaly patches or ring-shaped skin lesions that may be a symptom of an autoimmune disease called cutaneous lupus erythematosus.

Reactions during the infusion have occurred uncommonly with voriconazole (including flushing, fever, sweating, increased heart rate and shortness of breath). Your doctor may stop the infusion if this occurs.

As voriconazole has been known to affect the liver and the kidneys your doctor should monitor the function of your liver and kidney by doing blood tests. Please advise your doctor if you have any stomach pains or if your stools have a different consistency.

There have been reports of skin cancer in patients treated with voriconazole for long periods of time.

Sunburn or severe skin reaction following exposure to light or sun was experienced more frequently in children. If you or your child develops skin disorders, your doctor may refer you to a dermatologist, who after consultation may decide that it is important for you or your child to be seen on a regular basis. Elevated liver enzymes were also observed more frequently in children.

If any of these side effects persist or are troublesome, please tell your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Voriconazole

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the immediate and outer label after EXP. The expiry date refers to the last day of that month.

Do not store above 30°C. Store in the original package in order to protect from light.

Once reconstituted, Voriconazole should be used immediately, but if necessary may be stored for up to 24 hours at 2°C - 8°C (in a refrigerator) or in room temperature (15°C-25°C). Reconstituted Voriconazole needs to be further diluted with a compatible infusion solution first before it is infused. (Please refer to the end of this leaflet for further information).

Once diluted, the solutions can be stored for up to 48 hours at 2°C - 8°C (in a refrigerator) or up to 24 hours at 15-25°C (room temperature).

From a microbiological point of view, the product must be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C to 8°C, unless reconstitution/dilution has taken place in controlled and validated aseptic conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer require. These measures will help protect the environment.

6. Contents of the pack and other information

What Voriconazole contains

- The active substance is voriconazole.
- The other ingredients are hydroxypropylbetadex, arginine hydrochloride, sodium hydroxide (for pH adjustment), hydrochloric acid concentrated (for pH adjustment) (see section 2, Voriconazole contains sodium and cyclodextrins).

Each vial contains 200mg voriconazole, equivalent to a 10mg/ml solution when reconstituted as directed by your hospital pharmacist or nurse (see the information at the end of this leaflet).

What Voriconazole looks like and contents of the pack

Voriconazole is a white or almost white powder for solution for infusion presented in 30ml single use glass vials.

Marketing Authorisation Holder

Accord Healthcare Limited
Sage House, 319 Pinner Road
North Harrow, Middlesex, HA1 4HF
United Kingdom

Manufacturer

Actavis Italy S.p.A., Nerviano Plant
Via Pasteur 10
20014 Nerviano (Milan)
Italy

S.C. SINDAN-PHARMA S.R.L.
11th Ion Mihalache Blvd.

011171, Bucharest 1
Romania

Accord Healthcare Polska Sp.z o.o.
ul. Lutomska 50,
95-200 Pabianice
Poland

This leaflet was last revised in December 2024.

The following information is intended for healthcare professionals only:
Voriconazole Accordpharma 200mg Powder for Solution for Infusion

Reconstitution and Dilution information

- Voriconazole Powder for Solution for Infusion needs to first be reconstituted with either 19ml of Water for Injections or 19ml of 9mg/ml (0.9%) Sodium Chloride for Infusion to obtain an extractable volume of 20ml of clear concentrate containing 10mg/ml voriconazole.
- Discard the Voriconazole vial if the vacuum does not pull the diluent into the vial.
- It is recommended that a standard 20ml (non-automated) syringe be used to ensure that the exact amount (19.0ml) of Water for Injections or of 9mg/ml (0.9%) Sodium Chloride for Infusion is dispensed.
- The required volume of the reconstituted concentrate is then added to a recommended compatible infusion solution listed below to obtain a final Voriconazole solution containing 0.5 to 5mg/ml of voriconazole.
- This medicinal product is for single use only and any unused solution should be discarded and only clear solutions without particles should be used.
- For intravenous infusion only.
- For storage information, please refer to Section 5 ‘How to store Voriconazole’.

Required Volumes of 10mg/ml Voriconazole Concentrate

Body Weight (kg)	Volume of Voriconazole Concentrate (10mg/ml) required for:				
	3mg/kg dose (number of vials)	4mg/kg dose (number of vials)	6mg/kg dose (number of vials)	8mg/kg dose (number of vials)	9mg/kg dose (number of vials)
10	-	4.0ml (1)	-	8.0ml (1)	9.0ml (1)
15	-	6.0ml (1)	-	12.0ml (1)	13.5ml (1)
20	-	8.0ml (1)	-	16.0ml (1)	18.0ml (1)
25	-	10.0ml (1)	-	20.0ml (1)	22.5ml (2)
30	9.0ml (1)	12.0ml (1)	18.0ml (1)	24.0ml (2)	27.0ml (2)
35	10.5ml (1)	14.0ml (1)	21.0ml (2)	28.0ml (2)	31.5ml (2)
40	12.0ml (1)	16.0ml (1)	24.0ml (2)	32.0ml (2)	36.0ml (2)
45	13.5ml (1)	18.0ml (1)	27.0ml (2)	36.0ml (2)	40.5ml (3)
50	15.0ml (1)	20.0ml (1)	30.0ml (2)	40.0ml (2)	45.0ml (3)
55	16.5ml (1)	22.0ml (2)	33.0ml (2)	44.0ml (3)	49.5ml (3)
60	18.0ml (1)	24.0ml (2)	36.0ml (2)	48.0ml (3)	54.0ml (3)
65	19.5ml (1)	26.0ml (2)	39.0ml (2)	52.0ml (3)	58.5ml (3)
70	21.0ml (2)	28.0ml (2)	42.0ml (3)	-	-
75	22.5ml (2)	30.0ml (2)	45.0ml (3)	-	-
80	24.0ml (2)	32.0ml (2)	48.0ml (3)	-	-
85	25.5ml (2)	34.0ml (2)	51.0ml (3)	-	-
90	27.0ml (2)	36.0ml (2)	54.0ml (3)	-	-
95	28.5ml (2)	38.0ml (2)	57.0ml (3)	-	-

100	30.0ml (2)	40.0ml (2)	60.0ml (3)	-	-
-----	------------	------------	------------	---	---

Voriconazole is a single dose unpreserved sterile lyophilisate. Therefore, from a microbiological point of view, the reconstituted solution must be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C (in a refrigerator), unless reconstitution has taken place in controlled and validated aseptic conditions.

Compatible Infusion Solutions:

The reconstituted solution can be diluted with:

- Sodium Chloride 9mg/ml (0.9%) Solution for Injection
- 5% (50mg/ml) Glucose and Lactated Ringer's Intravenous Infusion
- 5% (50mg/ml) Glucose and 0.45% (4.5mg/ml) Sodium Chloride Intravenous Infusion
- 5% (50mg/ml) Glucose Intravenous Infusion
- 5% (50mg/ml) Glucose in 20 mEq Potassium Chloride Intravenous Infusion
- 5% (50mg/ml) Glucose and 0.9% (9mg/ml) Sodium Chloride Intravenous Infusion

The compatibility of Voriconazole with diluents other than listed above (or listed below under 'Incompatibilities') is unknown.

Incompatibilities:

Voriconazole must not be infused into the same line or cannula concomitantly with other drug infusions, including parenteral nutrition.

Infusions of blood products must not occur simultaneously with Voriconazole.

Infusion of total parenteral nutrition can occur simultaneously with Voriconazole but not in the same line or cannula.

Voriconazole must not be diluted with 4.2% (42mg/ml) Sodium Bicarbonate Infusion.

When reconstituted with Water for Injections, Voriconazole cannot be used in combination with Sodium Chloride 0.45% (4.5mg/ml) Solution for Injection and Compound Sodium Lactate Intravenous Infusion, due to low osmolality.

VORICONAZOLE PATIENT ALERT CARD

Please carry this card with you at all times

This card contains important safety information you need to be aware of before you are given Voriconazole for prevention of fungal infections and / or during treatment with Voriconazole for your fungal infection.

If you do not understand this information, please ask your doctor or healthcare professional involved in your care to explain it to you.

Show this card to any doctor or healthcare professional involved in your care.

See Voriconazole patient information leaflet for more information

Please report any suspect adverse drug reaction using the MHRA Yellow Card Scheme, Website, www.mhra.gov.uk/yellowcard

You should avoid exposure to direct sunlight during Voriconazole treatment. It is important to cover sun-exposed areas of skin and use sufficient sunscreen with high sun protection factor (SPF) as an increased sensitivity to the sun's UV rays can occur. There is a small chance that skin cancer could develop over time.

You should contact your doctor if you experience sunburn or severe skin reaction following exposure to light or sun.

Please ensure that you undergo all follow-up visits for blood test or skin evaluations arranged by your doctor. Please have a list of all your other medicines and medical conditions available at each visit to a healthcare professional.

Please complete.

Your Name:

Treating doctor's name:

Treatment Centre

(name / phone number)

Date Voriconazole first prescribed:

____/____/____