

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1 NAME OF THE MEDICINAL PRODUCT

Tramadol Krka 50 mg hard capsules

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains 50 mg tramadol hydrochloride.

For the full list of excipients, see section 6.1.

### 3 PHARMACEUTICAL FORM

Hard capsule (capsule)

The body of the capsule is white and the cap is blue. Capsules are filled with white or almost white powder. The size of the capsule is No: 4. The length of filled capsules is 14-15 mm.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Treatment of moderate to severe pain in adults and adolescents aged 12 years and over.

#### 4.2 Posology and method of administration

##### Posology

The dose should be adjusted to the intensity of the pain and the sensitivity of the individual patient. The lowest effective dose for analgesia should generally be selected. The total daily dose of 400 mg active substance should not be exceeded, except in special circumstances.

Unless otherwise prescribed, Tramadol should be administered as follows:

*Adults and adolescents aged 12 years and over*

Dosage form	Single dose	Total daily dose
Tramadol	50-100 mg Every 4 to 6 hours	400 mg

	(1 to 2 hard capsules)  (see section 5.1)	(Up to 8 hard capsules)
--	---	-------------------------

If there is no sufficient pain relief after administration of a single dose of 50 mg tramadol hydrochloride within 30 to 60 minutes, a second single dose of 50 mg can be administered.

If in severe pain the demand is likely to be higher, the higher single dose of Tramadol (100 mg tramadol hydrochloride) may be given as the initial dose.

**Acute Pain:** An initial dose of 100 mg is usually necessary. This can be followed by doses of 50 or 100 mg at 4 - 6 hourly intervals, and duration of treatment should be matched to clinical need.

**Pain Associated with Chronic Conditions:** An initial dose of 50 mg is advised and then titration according to pain severity. The need for continued treatment should be assessed at regular intervals as withdrawal symptoms and dependence have been reported (see section 4.4).

#### *Paediatric population*

Tramadol capsules are not suitable for children below the age of 12 years.

#### *Elderly*

A dose adjustment is not usually necessary in patients up to 75 years without clinically manifest hepatic or renal insufficiency. In elderly patients over 75 years elimination may be prolonged. Therefore, if necessary, the dosage interval is to be extended according to the patient's requirements.

#### *Renal insufficiency/dialysis and hepatic insufficiency*

In patients with renal and/or hepatic insufficiency the elimination of tramadol is delayed. In these patients prolongation of the dosage intervals should be carefully considered according to the patient's requirements.

#### Method of administration

For oral administration.

The capsules are to be taken whole, not divided or chewed, with sufficient liquid, with or without food.

#### *Duration of administration*

Tramadol should under no circumstances be administered for longer than absolutely necessary. If long-term pain treatment with Tramadol is necessary in view of the nature and severity of the illness, then careful and regular monitoring should be carried out (if necessary, with breaks in treatment) to establish whether and to what extent further treatment is necessary.

#### *Treatment goals and discontinuation*

Before initiating treatment with Tramadol, a treatment strategy including treatment duration and treatment goals, and a plan for end of the treatment, should be agreed together with the patient, in accordance with pain management guidelines. During treatment, there should be frequent contact between the physician and the patient to evaluate the need for continued treatment, consider discontinuation and to adjust dosages if needed. When a patient no longer requires therapy with tramadol, it may be

advisable to taper the dose gradually to prevent symptoms of withdrawal. In absence of adequate pain control, the possibility of hyperalgesia, tolerance and progression of underlying disease should be considered (see section 4.4).

### **4.3 Contraindications**

Tramadol is contraindicated:

- In hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- In acute intoxication with alcohol, hypnotics, analgesics, opioids, or other psychotropic medicinal products.
- In patients who are receiving MAO inhibitors or who have taken them within the last 14 days (see section 4.5).
- In patients with epilepsy not adequately controlled by treatment.
- For use in narcotic withdrawal treatment.

### **4.4 Special warnings and precautions for use**

Tramadol may only be used with particular caution in opioid-dependent patients, patients with head injury, shock, in an altered state of consciousness of uncertain origin, disorders of the respiratory centre or function, increased intracranial pressure.

In patients sensitive to opiates the product should only be used with caution.

Convulsions have been reported in patients receiving tramadol at the recommended dose levels. The risk may be increased when doses of tramadol exceed the recommended upper daily dose limit (400 mg). In addition, tramadol may increase the seizure risk in patients taking other medicinal products that lowers the seizure threshold (see section 4.5). Patients with epilepsy or those susceptible to seizures should be only treated with tramadol if there are compelling circumstances.

Caution should be used when treating patients with respiratory depression, or if concomitant CNS depressant drugs are being administered (see section 4.5), or if the recommended dosage is significantly exceeded (see section 4.9) as the possibility of respiratory depression cannot be excluded in these situations.

#### Serotonin syndrome

Serotonin syndrome, a potentially life-threatening condition, has been reported in patients receiving tramadol in combination with other serotonergic agents or tramadol alone (see sections 4.5, 4.8 and 4.9).

If concomitant treatment with other serotonergic agents is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose escalations.

Symptoms of serotonin syndrome may include mental status changes, autonomic instability, neuromuscular abnormalities and/or gastrointestinal symptoms.

If serotonin syndrome is suspected, a dose reduction or discontinuation of therapy should be considered depending on the severity of the symptoms. Withdrawal of the serotonergic drugs usually brings about a rapid improvement.

### Sleep-related breathing disorders

Opioids can cause sleep-related breathing disorders including central sleep apnea (CSA) and sleep-related hypoxemia. Opioid use increases the risk of CSA in a dose-dependent fashion. In patients who present with CSA, consider decreasing the total opioid dosage.

### Adrenal insufficiency

Opioid analgesics may occasionally cause reversible adrenal insufficiency requiring monitoring and glucocorticoid replacement therapy. Symptoms of acute or chronic adrenal insufficiency may include e.g. severe abdominal pain, nausea and vomiting, low blood pressure, extreme fatigue, decreased appetite, and weight loss.

### Tolerance and opioid use disorder (abuse and dependence)

Tolerance, physical and psychological dependence, and opioid use disorder (OUD) may develop upon repeated administration of opioids such as Tramadol. Repeated use of Tramadol can lead to opioid use disorder (OUD). A higher dose and longer duration of opioid treatment can increase the risk of developing OUD. Abuse or intentional misuse of Tramadol may result in overdose and/or death. The risk of developing OUD is increased in patients with a personal or a family history (parents or siblings) of substance use disorders (including alcohol use disorder), in current tobacco users or in patients with a personal history of other mental health disorders (e.g. major depression, anxiety and personality disorders).

Before initiating treatment with Tramadol and during the treatment, treatment goals and a discontinuation plan should be agreed with the patient (see section 4.2). Before and during treatment the patient should also be informed about the risks and signs of OUD. If these signs occur, patients should be advised to contact their physician.

Patients will require monitoring for signs of drug-seeking behaviour (e.g. too early requests for refills). This includes the review of concomitant opioids and psychoactive drugs (like benzodiazepines). For patients with signs and symptoms of OUD, consultation with an addiction specialist should be considered.

Tramadol is not suitable for use as a substitute in opioid-dependent patients. Although it is an opioid agonist, tramadol cannot suppress morphine withdrawal symptoms.

Concomitant use of tramadol and sedative medicines such as benzodiazepines or related drugs may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing with these sedative medicines should be reserved for patients for whom alternative treatment options are not possible. If a decision is made to prescribe tramadol concomitantly with sedative medicines, the lowest effective dose should be used, and the duration of treatment should be as short as possible.

The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and their caregivers to be aware of these symptoms (see section 4.5).

When a patient no longer requires therapy with tramadol, it may be advisable to taper the dose gradually to prevent symptoms of withdrawal.

### *CYP2D6 metabolism*

Tramadol is metabolised by the liver enzyme CYP2D6. If a patient has a deficiency or is completely lacking this enzyme an adequate analgesic effect may not be

obtained. Estimates indicate that up to 7% of the Caucasian population may have this deficiency. However, if the patient is an ultra-rapid metaboliser there is a risk of developing <side effects> of opioid toxicity even at commonly prescribed doses. General symptoms of opioid toxicity include confusion, somnolence, shallow breathing, small pupils, nausea, vomiting, constipation and lack of appetite. In severe cases this may include symptoms of circulatory and respiratory depression, which may be life threatening and very rarely fatal. Estimates of prevalence of ultra-rapid metabolisers in different populations are summarised below:

Population	Prevalence %
African/Ethiopian	29%
African American	3.4% to 6.5%
Asian	1.2% to 2%
Caucasian	3.6% to 6.5%
Greek	6.0%
Hungarian	1.9%
Northern European	1% to 2%

#### *Post-operative use in children*

There have been reports in the published literature that tramadol given post-operatively in children after tonsillectomy and/or adenoidectomy for obstructive sleep apnoea, led to rare, but life threatening adverse events. Extreme caution should be exercised when tramadol is administered to children for post-operative pain relief and should be accompanied by close monitoring for symptoms of opioid toxicity including respiratory depression.

#### *Children with compromised respiratory function*

Tramadol is not recommended for use in children in whom respiratory function might be compromised including neuromuscular disorders, severe cardiac or respiratory conditions, upper respiratory or lung infections, multiple trauma or extensive surgical procedures. <These factors may worsen symptoms of opioid toxicity>.

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Tramadol should not be combined with MAO inhibitors (see section 4.3).

In patients treated with MAO inhibitors in the 14 days prior to the use of the opioid pethidine, life-threatening interactions on the central nervous system, respiratory and cardiovascular function have been observed. The same interactions with MAO inhibitors cannot be ruled out during treatment with Tramadol.

Concomitant administration of Tramadol with other centrally depressant medicinal products including alcohol may potentiate the CNS effects (see section 4.8).

The concomitant use of opioids with sedative medicines such as benzodiazepines or related drugs increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. The dose and duration of concomitant use

should be limited (see section 4.4).

The concomitant use of Tramadol with gabapentinoids (gabapentin and pregabalin) may result in respiratory depression, hypotension, profound sedation, coma or death.

The results of pharmacokinetic studies have so far shown that on the concomitant or previous administration of cimetidine (enzyme inhibitor) clinically relevant interactions are unlikely to occur.

Simultaneous or previous administration of carbamazepine (enzyme inducer) may reduce the analgesic effect and shorten the duration of action.

Tramadol can induce convulsions and increase the potential for selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, antipsychotics and other seizure threshold-lowering medicinal product (such as bupropion, mirtazapine, tetrahydrocannabinol) to cause convulsions.

Concomitant therapeutic use of tramadol and serotonergic drugs, such as selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), MAO inhibitors (see section 4.3), tricyclic antidepressants and mirtazapine may cause serotonin syndrome, a potentially life-threatening condition (see sections 4.4 and 4.8).

Caution should be exercised during concomitant treatment with tramadol and coumarin derivatives (e.g. warfarin) due to reports of increased INR with major bleeding and ecchymoses in some patients.

Other active substances known to inhibit CYP3A4, such as ketoconazole and erythromycin, might inhibit the metabolism of tramadol (N-demethylation), probably also the metabolism of the active O-demethylated metabolite. The clinical importance of this interaction is unknown (see section 4.8).

In a limited number of studies the pre- or postoperative application of the antiemetic 5-HT<sub>3</sub> antagonist ondansetron increased the requirement of tramadol in patients with postoperative pain.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

Animal studies with tramadol revealed at very high doses effects on organ development, ossification and neonatal mortality. Tramadol crosses the placenta. There is inadequate evidence available on the safety of tramadol in human pregnancy. Therefore Tramadol should not be used in pregnant women.

Tramadol - administered before or during birth - does not affect uterine contractility. In neonates it may induce changes in the respiratory rate which are usually not clinically relevant. Chronic use during pregnancy may lead to neonatal withdrawal symptoms.

### Breast-feeding

Approximately 0.1% of the maternal dose of tramadol is excreted in breast milk. In the immediate post-partum period, for maternal oral daily dosage up to 400 mg, this

corresponds to a mean amount of tramadol ingested by breast-fed infants of 3% of the maternal weight-adjusted dosage. For this reason tramadol should not be used during lactation or alternatively, breast-feeding should be discontinued during treatment with tramadol. Discontinuation of breast-feeding is generally not necessary following a single dose of tramadol.

#### Fertility

Post marketing surveillance does not suggest an effect of tramadol on fertility. Animal studies did not show an effect of tramadol on fertility.

### **4.7 Effects on ability to drive and use machines**

Even when taken according to instructions, Tramadol may cause effects such as somnolence and dizziness and therefore may impair the reactions of drivers and machine operators. This applies particularly in conjunction and other psychotropic substances, particularly alcohol.

This medicine can impair cognitive function and can affect a patient's ability to drive safely. This class of medicine is in the list of drugs included in regulations under 5a of the Road Traffic Act 1988. When prescribing this medicine, patients should be told:

The medicine is likely to affect your ability to drive

- Do not drive until you know how the medicine affects you
- It is an offence to drive while under the influence of this medicine
- However, you would not be committing an offence (called 'statutory defence') if:
  - The medicine has been prescribed to treat a medical or dental problem and
  - You have taken it according to the instructions given by the prescriber and in the information provided with the medicine and
  - It was not affecting your ability to drive safely

### **4.8 Undesirable effects**

In evaluating undesirable effects, the frequencies are defined as follows:

- Very common ( $\geq 1/10$ )
- Common ( $\geq 1/100$  to  $< 1/10$ )
- Uncommon ( $\geq 1/1,000$  to  $< 1/100$ )
- Rare ( $\geq 1/10,000$  to  $< 1/1,000$ )
- Very rare ( $< 1/10,000$ )
- Not known (cannot be estimated from the available data)

The most commonly reported adverse reactions are nausea and dizziness, both occurring in more than 10% of patients.

	Very common	Common	Uncommon	Rare	Not known
Immune				Allergic reactions	

system disorders				(e.g. dyspnoea, bronchospasm, wheezing, angioneurotic oedema) and anaphylaxis.	
Metabolism and nutrition disorders				Changes in appetite.	Hypoglycaemia.
Psychiatric disorders				<p>Hallucinations, confusion, sleep disturbance, delirium, anxiety and nightmares. Psychic adverse reactions may occur following administration of tramadol which vary individually in intensity and nature (depending on personality and duration of treatment). These include changes in mood (usually elation, occasionally dysphoria), changes in activity (usually suppression, occasionally increase) and changes in cognitive and sensorial capacity (e.g. decision behaviour, perception disorders). Drug dependence may occur.<sup>1</sup></p> <p>Symptoms of drug withdrawal syndrome, similar to those occurring during opiate withdrawal, may occur as follows: agitation, anxiety, nervousness, insomnia,</p>	

				hyperkinesia, tremor and gastrointestinal symptoms. Other symptoms that have very rarely been seen with tramadol discontinuation include: panic attacks, severe anxiety, hallucinations, paraesthesia, tinnitus and unusual CNS symptoms (e.g. confusion, delusions, depersonalisation, derealisation, paranoia).	
Nervous system disorders	Dizziness	Headache, somnolence		Paraesthesia, tremor, involuntary muscle contractions, abnormal coordination, syncope, speech disorders. Convulsions occurred mainly after administration of high doses of tramadol or after concomitant treatment with medicinal products which can lower the seizure threshold (see sections 4.4 and 4.5).	Serotonin syndrome
Eye disorders				Miosis, mydriasis, blurred vision.	
Cardiac disorders			Cardiovascular regulation (palpitation, tachycardia). These adverse reactions may occur	Bradycardia.	

			especially on intravenous administration and in patients who are physically stressed.		
Vascular disorders			Cardiovascular regulation (postural hypotension or cardiovascular collapse). These adverse reactions may occur especially on intravenous administration and in patients who are physically stressed.		
Respiratory, thoracic and mediastinal disorders				Respiratory depression, dyspnoea. If the recommended doses are considerably exceeded and other centrally depressant substances are administered concomitantly (see section 4.5), respiratory depression may occur. Worsening of asthma has been reported, though a causal relationship has not been established.	Hiccups.
Gastrointestinal disorders	Nausea.	Vomiting, constipation, dry mouth.	Retching; gastrointestinal discomfort (a feeling of pressure in the stomach, bloating),		

			diarrhoea.		
Hepatobiliary disorders					In a few isolated cases an increase in liver enzyme values has been reported in a temporal connection with the therapeutic use of tramadol.
Skin and subcutaneous tissue disorders		Hyperhidrosis.	Dermal reactions (e.g. pruritus, redness of the skin, urticaria)		
Musculoskeletal and connective tissue disorders				Motorial weakness.	
Renal and urinary disorders				Micturition disorders (dysuria and urinary retention).	
General disorders and administration site conditions		Fatigue			
Investigations				Increase in blood pressure.	

<sup>1</sup>Repeated use of Tramadol can lead to drug dependence, even at therapeutic doses. The risk of drug dependence may vary depending on a patient's individual risk factors, dosage, and duration of opioid treatment (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme, Website: [yellowcard.mhra.gov.uk](http://yellowcard.mhra.gov.uk) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## 4.9 Overdose

### Symptoms

In principle, on intoxication with tramadol symptoms similar to those of other centrally acting analgesics (opioids) are to be expected. These include in particular miosis, vomiting, cardiovascular collapse, consciousness disorders up to coma, convulsions and respiratory depression up to respiratory arrest. Serotonin syndrome has also been reported.

### Management

The general emergency measures apply. Keep open the respiratory tract (aspiration!), maintain respiration and circulation depending on the symptoms. The antidote for respiratory depression is naloxone. In animal experiments naloxone had no effect on convulsions. In such cases diazepam should be given intravenously.

In case of intoxication orally, gastrointestinal decontamination with activated charcoal or by gastric lavage is only recommended within 2 hours after tramadol intake. Gastrointestinal decontamination at a later time point may be useful in case of intoxication with exceptionally large quantities.

Tramadol is minimally eliminated from the serum by haemodialysis or haemofiltration. Therefore treatment of acute intoxication with Tramadol with haemodialysis or haemofiltration alone is not suitable for detoxification.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: analgesics, other opioids, ATC code: N02AX02.

#### *Mechanism of action*

Tramadol is a centrally acting opioid analgesic. It is a non-selective pure agonist at  $\mu$ ,  $\delta$  and  $\kappa$  opioid receptors with a higher affinity for the  $\mu$  receptor. Other mechanisms which may contribute to its analgesic effect are inhibition of neuronal reuptake of noradrenaline and enhancement of serotonin release.

#### *Clinical efficacy and safety*

Tramadol has an antitussive effect. In contrast to morphine, analgesic doses of tramadol over a wide range have no respiratory depressant effect. Also gastrointestinal motility is less affected. Effects on the cardiovascular system tend to be slight. The potency of tramadol is reported to be 1/10 (one tenth) to 1/6 (one sixth) that of morphine.

#### *Paediatric population*

Effects of enteral and parenteral administration of tramadol have been investigated in clinical trials involving more than 2000 paediatric patients ranging in age from neonate to 17 years of age. The indications for pain treatment studied in those trials included pain after surgery (mainly abdominal), after surgical tooth extractions, due to fractures, burns and traumas as well as other painful conditions likely to require analgesic treatment for at least 7 days.

At single doses of up to 2 mg/kg or multiple doses of up to 8 mg/kg per day (to a maximum of 400 mg per day) efficacy of tramadol was found to be superior

to placebo, and superior or equal to paracetamol, nalbuphine, pethidine or low dose morphine. The conducted trials confirmed the efficacy of tramadol. The safety profile of tramadol was similar in adult and paediatric patients older than 1 year (see section 4.2).

## 5.2 Pharmacokinetic properties

### *Absorption*

More than 90% of tramadol is absorbed after oral administration. The mean absolute bioavailability is approximately 70%, irrespective of the concomitant intake of food. The difference between absorbed and non-metabolised available tramadol is probably due to the low first-pass effect. The first-pass effect after oral administration is a maximum of 30%.

### *Distribution*

Following oral use of 100 mg tramadol in liquid form, peak plasma concentrations after 1.2 hours are calculated to be  $C_{max} = 309 \pm 90$  ng/ml. Following the same dose in solid oral form peak plasma concentrations after 2 hours are  $C_{max} = 280 \pm 49$  ng/ml.

Tramadol has a high tissue affinity ( $V_{d,\beta} = 203 + 40$  l). It has a plasma protein binding of about 20%.

Tramadol passes the blood-brain and placental barriers. Very small amounts of the substance and its O-desmethyl derivative are found in the breast-milk (0.1% and 0.02% respectively of the applied dose).

### *Biotransformation*

In humans tramadol is mainly metabolised by means of N- and O-demethylation and conjugation of the O-demethylation products with glucuronic acid. Only O-desmethyltramadol is pharmacologically active. There are considerable interindividual quantitative differences between the other metabolites. So far, eleven metabolites have been found in the urine. Animal experiments have shown that O-desmethyltramadol is more potent than the parent substance by the factor 2 - 4. Its half-life  $t_{1/2,\beta}$  (6 healthy volunteers) is 7.9 h (range 5.4 - 9.6 h) and is approximately that of tramadol.

The inhibition of one or both types of the isoenzymes CYP3A4 and CYP2D6 involved in the biotransformation of tramadol may affect the plasma concentration of tramadol or its active metabolite.

### *Elimination*

Tramadol and its metabolites are almost completely excreted via the kidneys. Cumulative urinary excretion is 90% of the total radioactivity of the administered dose. Elimination half-life  $t_{1/2,\beta}$  is approximately 6 h, irrespective of the mode of administration. In patients above 75 years of age it may be prolonged by a factor of approximately 1.4. In cases of impaired hepatic and renal function the half-life may be slightly prolonged. In patients with cirrhosis of the liver, elimination half-lives of  $13.3 \pm 4.9$  hours (tramadol) and  $18.5 \pm 9.4$  hours (O-desmethyltramadol), in an extreme case 22.3 hours and 36 hours respectively, have been determined. In patients with renal insufficiency (creatinine clearance  $< 5$  ml/min) the values were  $11 \pm 3.2$  hours and  $16.9 \pm 3$  hours, in an extreme case 19.5 hours and 43.2 hours respectively.

### *Linearity/non-linearity*

Tramadol has a linear pharmacokinetic profile within the therapeutic dosage range.

### *Pharmacokinetic/pharmacodynamics relationship(s)*

The relationship between serum concentrations and the analgesic effect is dose-dependent, but

varies considerably in isolated cases. A serum concentration of 100 - 300 ng/ml is usually effective.

#### *Paediatric population*

The pharmacokinetics of tramadol and O-desmethyltramadol after single-dose and multiple-dose oral administration to subjects aged 1 year to 16 years were found to be generally similar to those in adults when adjusting for dose by body weight, but with a higher between-subject variability in children aged 8 years and below.

In children below 1 year of age, the pharmacokinetics of tramadol and O-desmethyltramadol have been investigated, but have not been fully characterized. Information from studies including this age group indicates that the formation rate of O-desmethyltramadol via CYP2D6 increases continuously in neonates, and adult levels of CYP2D6 activity are assumed to be reached at about 1 year of age. In addition, immature glucuronidation systems and immature renal function may result in slow elimination and accumulation of O-desmethyltramadol in children under 1 year of age.

### **5.3 Preclinical safety data**

On repeated oral and parenteral administration of tramadol for 6 - 26 weeks in rats and dogs and oral administration for 12 months in dogs haematological, clinico-chemical and histological investigations showed no evidence of any substance-related changes. Central nervous manifestations only occurred after high doses considerably above the therapeutic range: restlessness, salivation, convulsions, and reduced weight gain. Rats and dogs tolerated oral doses of 20 mg/kg and 10 mg/kg body weight respectively, and dogs rectal doses of 20 mg/kg body weight without any reactions.

In rats tramadol dosages from 50 mg/kg/day upwards caused toxic effects in dams and raised neonate mortality. In the offspring retardation occurred in the form of ossification disorders and delayed vaginal and eye opening. Male fertility was not affected. After higher doses (from 50 mg/kg/day upwards) females exhibited a reduced pregnancy rate. In rabbits there were toxic effects in dams from 125 mg/kg upwards and skeletal anomalies in the offspring.

In some in-vitro test systems there was evidence of mutagenic effects. In-vivo studies showed no such effects. According to knowledge gained so far, tramadol can be classified as non-mutagenic.

Studies on the tumorigenic potential of tramadol hydrochloride have been carried out in rats and mice. The study in rats showed no evidence of any substance-related increase in the incidence of tumours. In the study in mice there was an increased incidence of liver cell adenomas in male animals (a dose-dependent, non-significant increase from 15 mg/kg upwards) and an increase in pulmonary tumours in females of all dosage groups (significant, but not dose-dependent).

## **6 PHARMACEUTICAL PARTICULARS**

## **6.1 List of excipients**

### Capsule contents

Cellulose, microcrystalline  
Sodium starch glycolate (type A)  
Talc  
Magnesium stearate (E470b)

### Capsule shell

Titanium dioxide (E171)  
Indigo carmine (E132)  
Gelatin

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

5 years

## **6.4 Special precautions for storage**

Do not store above 30°C.  
Store in the original package in order to protect from moisture.

## **6.5 Nature and contents of container**

Blisters (PVC film/Alu foil): 10, 20, 30, 50, 60, 90 and 100 hard capsules, in a box.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

No special requirements for disposal.  
Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

KRKA, d.d, Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

**8      MARKETING AUTHORISATION NUMBER(S)**

PL 01656/0248

**9      DATE OF FIRST AUTHORISATION/RENEWAL OF THE  
AUTHORISATION**

28/03/2024

**10     DATE OF REVISION OF THE TEXT**

**19/07/2024**