



Information for the User Ibuprofen and Codeine 200 mg/12.8 mg Tablets Ibuprofen, Codeine Phosphate Hemihydrate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Ibuprofen and Codeine 200 mg/12.8 mg Tablets may be referred to as this medicine throughout this leaflet

- Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.
- Keep this leaflet. You may need to read it again.
 - Ask your pharmacist if you need more information or advice.
 - If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
 - You must talk to a doctor if you do not feel better or if you feel worse within 3 days.
 - This medicine can only be used for the short-term treatment of acute moderate pain such as headaches, rheumatic and muscular pain, backache, migraine, period pain, dental pain and neuralgia that is not relieved by aspirin, ibuprofen or paracetamol alone.
 - This medicine should only be taken when necessary.
 - You should only take this medicine for a maximum of 3 days** at a time. If you need to take it for longer than 3 days you should see your doctor or pharmacist for advice.
 - This medicine contains codeine which can cause addiction** if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it.
 - If you take this medicine for headaches for more than 3 days** it can make them worse.

- What Ibuprofen and Codeine 200 mg/12.8 mg Tablets are and what they are used for
- What you need to know before you take Ibuprofen and Codeine 200 mg/12.8 mg Tablets
- How to take Ibuprofen and Codeine 200 mg/12.8 mg Tablets
- Possible side effects
- How to store Ibuprofen and Codeine 200 mg/12.8 mg Tablets
- Contents of the pack and other information

1. What Ibuprofen and Codeine 200 mg/12.8 mg Tablets are and what they are used for

The active ingredients (which make this medicine work) are Ibuprofen and Codeine phosphate.

This medicine contains codeine. Codeine belongs to a group of medicines called opioid analgesics which act to relieve pain.

This medicine also contains ibuprofen, Ibuprofen, a non-steroidal anti-inflammatory medicine (NSAID) which also acts to reduce swelling (inflammation).

This medicine can be used in adults and children over 12 years of age, for the short-term treatment of acute moderate pain such as headaches, rheumatic and muscular pain, backache, migraine, period pain, dental pain and neuralgia that is not relieved by aspirin, ibuprofen or paracetamol alone.

2. What you need to know before you take Ibuprofen and Codeine 200 mg/12.8 mg Tablets

This medicine contains codeine which can cause addiction if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it. **If you take a painkiller for headaches for more than 3 days** it can make them worse.

- ✗ Do not take Ibuprofen and Codeine Tablets:**
- If you have (or have had two or more episodes of) a stomach ulcer, perforation, or bleeding
 - If you have breathing difficulties (including respiratory depression) or raised intracranial pressure
 - If you are allergic to ibuprofen, codeine, or any other ingredient of this medicine (listed in section 6), or to aspirin or other painkillers
 - If you suffer from severe liver, kidney or heart problems
 - If you suffer from chronic constipation
 - If you have had gastrointestinal bleeding or perforation when previously taking NSAIDs
 - If you have had a worsening of asthma, skin rash, itchy runny nose or facial swelling when previously taking ibuprofen, aspirin or similar medicines
 - If you are taking other NSAID painkillers or aspirin with a daily dose above 75mg
 - If you are pregnant, and in the last 3 months of pregnancy
 - If you are breastfeeding (see "Other important information")
 - If you are under 12 years old
 - If you are taking monoamine oxidase inhibitors or if you have taken them in the last 14 days
 - For pain relief in children and adolescents (0-18 years of age) after removal of their tonsils or adenoids due to obstructive sleep apnoea syndrome
 - If you know that you metabolise very rapidly codeine into morphine

Warnings and precautions

Tolerance, dependence, and addiction

This medicine contains codeine, which is an opioid medicine. It can cause dependence and/or addiction. Repeated use of opioids can result in the drug being less effective (you become accustomed to it, known as tolerance). Repeated use of Ibuprofen and Codeine Tablets can also lead to dependence, abuse and addiction, which may result in life-threatening overdose. If you are taking Ibuprofen and Codeine Tablets for longer than the recommended time or at higher than recommended doses the risk of these side effects can increase and you are also at risk of serious harms to the stomach/gut and kidneys, as well as very low levels of potassium in your blood. These can be fatal (see section 4).

Dependence or addiction can make you feel that you are no longer in control of how much medicine you need to take or how often you need to take it.

The risk of becoming dependent or addicted varies from person to person. You may have a greater risk of becoming dependent on or addicted to Ibuprofen and Codeine Tablets if:

- You or anyone in your family have ever abused or been dependent on alcohol, prescription medicines or illegal drugs ("addiction").
- You are a smoker.
- You have ever had problems with your mood (depression, anxiety, or a personality disorder) or have been treated by a psychiatrist for other mental illnesses.

If you notice any of the following signs whilst taking Ibuprofen and Codeine Tablets it could be a sign that you have become dependent or addicted:

- You need to take this medicine for longer than advised by your doctor
- You need to take more than the recommended dose
- You might feel that you need to carry on taking your medicine, even when it doesn't help to relieve your pain.
- You are using the medicine for reasons other than prescribed, for instance, 'to stay calm' or 'help you sleep'
- You have made repeated, unsuccessful attempts to quit or control the use of the medicine
- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again ('withdrawal effects')

If you notice any of these signs, speak to your doctor to discuss the best treatment pathway for you, including when it is appropriate to stop and how to stop safely (See section 3, When you stop taking Ibuprofen and Codeine Tablets).

Signs of an allergic reaction to this medicine, including breathing problems, swelling of the face and neck region (angioedema), chest pain have been reported with ibuprofen.

Stop taking this medicine immediately and immediately contact your doctor or medical emergency department if you notice any of these signs.

Rarely, increasing the dose of this medicine can make you more sensitive to pain. If this happens, you need to speak to your prescriber about your treatment.

Contact your doctor if you experience severe upper abdominal pain possibly radiating to the back, nausea, vomiting or fever as this could be symptoms associated with inflammation of the pancreas (pancreatitis) and the biliary tract system.

This medicine belongs to a group of medicines which **may impair fertility in women**. This is reversible on stopping the medicine. It is unlikely that this medicine, used occasionally will affect your chances of becoming pregnant. However, tell your doctor before taking this medicine if you have problems becoming pregnant.

Anti-inflammatory/painkiller medicines like ibuprofen may be associated with a **small increased risk of heart attack or stroke**. Any risk is more likely with high doses or prolonged treatment. **Do not exceed the recommended dose or take for longer than necessary to control your symptoms (3 days).**

If you have heart problems, have had a stroke or think that you might be at risk of these conditions (for example if you have high blood pressure, diabetes, high cholesterol, or are a smoker), you should discuss this treatment with your doctor or pharmacist.

Do not drink alcohol (wine, beer, spirits) whilst taking this medicine. Alcohol may make you feel more drowsy.

Children and adolescents

Use in children and adolescents after surgery: Codeine should not be used for pain relief in children and adolescents after removal of the tonsils or adenoids due to obstructive sleep apnoea syndrome.

Use in children with breathing problems: Codeine is not recommended in children with breathing problems, since the symptoms of morphine toxicity may be worse in these children.

Pregnancy and breast-feeding

Pregnancy: Do not take this medicine if you are in the last 3 months of pregnancy as it could harm your unborn child or cause problems at delivery. It can cause kidney and heart problems in your unborn baby. It may affect your and your baby's tendency to bleed and cause labour to be later or longer than expected.

You should not take this medicine during the first 6 months of pregnancy unless absolutely necessary and advised by your doctor. If you need treatment during this period or while you are trying to get pregnant, the lowest dose for the shortest time possible should be used. If taken for more than a few days from 20 weeks of pregnancy onward, this medicine can cause kidney problems in your unborn baby that may lead to low levels of amniotic fluid that surrounds the baby (oligohydramnios) or narrowing of a blood vessel (ductus arteriosus) in the heart of the baby. If you need treatment for longer than a few days, your doctor may recommend additional monitoring.

Breast-feeding: Do not take codeine while you are breast-feeding. Codeine and morphine pass into breast milk.

- Driving and using machines**
- This medicine can affect your ability to drive or use machines as it may make you feel sleepy or dizzy or cause drowsiness.
- Do not drive or use machines while taking this medicine until you are sure you are not affected
 - It is an offence to drive if this medicine affects your ability to drive
 - However, you would not be committing an offence if:
 - The medicine has been prescribed to treat a medical or dental problem and
 - You have taken it according to the instructions given by the prescriber or in the information provided with the medicine and
 - It was not affecting your ability to drive safely.

Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking this medicine.

Ibuprofen and Codeine Tablets contain sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Ibuprofen and Codeine 200 mg/12.8 mg Tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Adults, the elderly and children aged 12 years and older: Swallow 1 or 2 tablets with water, up to three times a day as required. Leave at least four hours between doses.

Do not take more than 6 tablets in any 24-hour period.

This medicine should not be taken by children below the age of 12 years, due to the risk of severe breathing problems. Do not take for more than 3 days. Ibuprofen and Codeine Tablets should be used for 3 days only to relieve symptoms. If no effective pain relief is achieved while taking the medicine, you should seek the advice of a healthcare professional.

If symptoms persist, or pain or fever worsen, or if any new symptoms occur, consult your doctor or pharmacist.

This medicine is for **short-term use only**. The lowest effective dose should be used for the shortest duration necessary to relieve symptoms. If you have an infection, consult a doctor without delay if symptoms (such as fever and pain) persist or worsen (see section 2).

! If you take more Ibuprofen and Codeine Tablets than you should

If you have taken more of this medicine than you should, or if children have taken this medicine by accident always contact a doctor or nearest hospital to get an opinion on the risk and advice on action to be taken.

The symptoms can include nausea, stomach pain, vomiting (may be blood streaked), headache, ringing in the ears, confusion and shaky eye movement. At high doses, drowsiness, chest pain, palpitations, loss of consciousness, convulsions (mainly in children), weakness and dizziness, blood in urine, cold body feeling, and breathing problems have been reported.

When you stop taking Ibuprofen and Codeine Tablets

This medicine contains codeine and can cause addiction if you take it continuously for more than 3 days. When you stop taking it you may get withdrawal symptoms. You should talk to your doctor or pharmacist if you think you are suffering from withdrawal symptoms.

If Ibuprofen and Codeine Tablets have been prescribed to you, before starting treatment and regularly during treatment, your doctor will discuss with you what you may expect from using Ibuprofen and Codeine Tablets, when and how long you need to take it, when to contact your doctor, and when you need to stop it.

4. Possible side effects

Some people may have side-effects when taking this medicine. If you have any unwanted side effects you should seek advice

from your doctor, pharmacist or healthcare professional.

STOP TAKING this medicine and **seek immediate medical help** if you develop:

- signs of **intestinal bleeding** such as: bright red faeces (stools/motions), black tarry stools, vomiting blood or dark particles that look like coffee grounds.
- signs of rare but serious **allergic reaction** such as severe skin rashes, peeling, flaking or blistering skin, facial swelling, unexplained wheezing, shortness of breath, easy bruising.
- chest pain, which can be a sign of a potentially serious allergic reaction called Kounis syndrome.
- redish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (exfoliative dermatitis, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis).
- widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome).
- a red, scaly widespread rash with bumps under the skin and blisters accompanied by fever. The symptoms usually appear at the initiation of treatment (acute generalised exanthematous pustulosis). See also section 2.
- meningitis (e.g. stiff neck, headache, fever, disorientation).
- kidney problems, which may lead to kidney failure (You may pass more or less urine, have blood in the urine or cloudy urine, or feel breathless, very tired or weak, have no appetite, or have swollen ankles).

Ibuprofen and Codeine Tablets, especially when taken at higher than recommended doses or for a prolonged period of time, can cause damage to your kidneys and affect them removing acids properly from your blood into the urine (renal tubular acidosis). It can also cause very low levels of potassium in your blood (see section 2). This is a very serious condition and will require immediate treatment. Signs and symptoms include muscle weakness and light-headedness.

Tell your doctor if you experience the below other possible side effects:

- Uncommon:** may affect up to 1 in 100 people
- indigestion, heartburn, nausea
 - stomach pain or other abnormal stomach symptoms
 - rashes, itching
 - headache.

Rare: may affect up to 1 in 1,000 people

- constipation.
- Very rare:** may affect up to 1 in 10,000 people

Not known: frequency cannot be estimated from the available data

- worsening of asthma
- sweating, thirst, muscle weakness or tremors, sleeplessness
- fast, irregular heart beat
- dizziness or drowsiness
- increased sensitivity to pain
- skin becomes sensitive to light
- symptoms associated with inflammation of the pancreas (pancreatitis) and the biliary tract system (a problem affecting a valve in the intestines known as sphincter of Oddi dysfunction), e.g. severe upper abdominal pain possibly radiating to the back, nausea, vomiting or fever.

Medicines such as this medicine may be associated with a small increased risk of heart attack (myocardial infarction) or stroke.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

! How do I know if I am addicted?

If you take the medicine according to the instructions on the pack it is unlikely that you will become addicted to the medicine. **However, if the following apply to you** it is important that you talk to your doctor:

- You need to take the medicine for longer periods of time
- You need to take more than the recommended dose
- When you stop taking the medicine you feel very unwell but you feel better if you start taking the medicine again.

5. How to store Ibuprofen and Codeine 200 mg/12.8 mg Tablets

These film-coated tablets do not require any special storage conditions.

Store this medicine in a safe and secure storage space, where other people cannot access it. It can cause serious harm and be fatal to people when it has not been intended for them. Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the foil or flap of the carton.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Ibuprofen and Codeine Tablets contain

Each film-coated tablet contains Codeine Phosphate Hemihydrate 12.8 mg and Ibuprofen 200 mg, which are the active ingredients.

As well as the active ingredients, the film-coated tablets also contain microcrystalline cellulose, sodium starch glycolate, pregelatinised maize starch, hypromellose. The tablet coating contains hypromellose, titanium dioxide (E171), talc.

What Ibuprofen and Codeine Tablets look like and contents of the pack

The pack contains 16, 18, 24 or 32 white capsule-shaped tablets embossed with 1+. Not all pack sizes may be marketed.

Marketing Authorisation Holder

The Boots Company PLC, Nottingham, NG2 3AA, UK.

Manufacturer

Custom Pharmaceuticals Ltd, Conway Street, Hove, East Sussex, BN3 3LW, UK.

This leaflet was last revised in August 2026.

Other sources of information

To request a copy of this leaflet in Braille, large print or audio please call free of charge:

0800 198 5000 (UK only)

Please be ready to give the following information:

Product name: Ibuprofen and Codeine 200 mg/12.8 mg Tablets

Reference Number: 00014/0662

This is a service provided by the Royal National Institute of Blind People.

AL0313/13
AL0392/13
LFLT00103