

# Crestor® 5mg, 10mg, 20mg and 40mg Film-coated Tablets

## Rosuvastatin 5mg, 10mg, 20mg and 40mg Film-coated Tablets

(rosuvastatin calcium)

Your medicine is known by one of the above names, but will be referred to as Crestor throughout this:

### Patient Information Leaflet

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- **This medicine has been prescribed for you only.** Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

- 1) What Crestor is and what it is used for
- 2) What you need to know before you take Crestor
- 3) How to take Crestor
- 4) Possible side effects
- 5) How to store Crestor
- 6) Contents of the pack and other information

#### 1) What Crestor is and what it is used for

Crestor belongs to a group of medicines called statins.

##### You have been prescribed Crestor because:

- You have a high cholesterol level. This means you are at risk from a heart attack or stroke. Crestor is used in adults, adolescents and children 6 years or older to treat high cholesterol.
- You have been advised to take a statin, because changing your diet and doing more exercise were not enough to correct your cholesterol levels. You should continue with your cholesterol-lowering diet and exercise while you are taking Crestor.

Or

- You have other factors that increase your risk of having a heart attack, stroke or related health problems.

Heart attack, stroke and other problems can be caused by a disease called atherosclerosis. Atherosclerosis is due to build-up of fatty deposits in your arteries.

#### Why it is important to keep taking Crestor

Crestor is used to correct the levels of fatty substances in the blood called lipids, the most common of which is cholesterol.

There are different types of cholesterol found in the blood – ‘bad’ cholesterol (LDL-C) and ‘good’ cholesterol (HDL-C).

- Crestor can reduce the ‘bad’ cholesterol and increase the ‘good’ cholesterol.
- It works by helping to block your body’s production of ‘bad’ cholesterol. It also improves your body’s ability to remove it from your blood.

For most people, high cholesterol does not affect the way they feel because it does not produce any symptoms. However, if it is left untreated, fatty deposits can build up in the walls of your blood vessels causing them to narrow.

Sometimes, these narrowed blood vessels can get blocked which can cut off the blood supply to the heart or brain leading to a heart attack or a stroke. By lowering your cholesterol levels, you can reduce your risk of having a heart attack, a stroke or related health problems.

You need to **keep taking Crestor**, even if it has got your cholesterol to the right level, because **it prevents your cholesterol levels from creeping up again** and causing build-up of fatty deposits. However, you should stop if your doctor tells you to do so, or you have become pregnant.

#### 2) What you need to know before you take Crestor

Do not take Crestor:

- **If you have ever had an allergic reaction to Crestor**, or to any of its ingredients.
- **If you are pregnant** or breast-feeding. If you become pregnant while taking Crestor **stop taking it immediately and tell your doctor**. Women should avoid becoming pregnant while taking Crestor by using suitable contraception.
- **If you have liver disease.**
- **If you have severe kidney problems.**
- **If you have repeated or unexplained muscle aches or pains.**
- **If you take a drug combination of sofosbuvir/velpatasvir/voxilaprevir** (used for viral infection of the liver called hepatitis C).
- **If you take a drug called ciclosporin** (used, for example, after organ transplants).

If any of the above applies to you (or you are in doubt), please go back and see your doctor.

#### In addition, do not take Crestor 40 mg (the highest dose):

- **If you have moderate kidney problems** (if in doubt, please ask your doctor).
- **If your thyroid gland** is not working properly.
- **If you have had any repeated or unexplained muscle aches or pains**, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines.
- **If you regularly drink large amounts of alcohol.**
- **If you are of Asian origin** (Japanese, Chinese, Filipino, Vietnamese, Korean and Indian).
- **If you take other medicines called fibrates** to lower your cholesterol.

If any of the above applies to you (or you are in doubt), **please go back and see your doctor.**

#### Warnings and precautions

Talk to your doctor or pharmacist before taking Crestor.

- **If you have problems with your kidneys.**
- **If you have problems with your liver.**
- **If you have had repeated or unexplained muscle aches or pains**, a personal or family history of muscle problems, or a previous history of muscle problems when taking other cholesterol-lowering medicines. Tell your doctor immediately if you have unexplained muscle aches or pains, especially if you feel unwell or have a fever. Also, tell your doctor or pharmacist if you have a muscle weakness that is constant.
- **If you have or have had myasthenia** (a disease with general muscle weakness including in some cases muscles used when breathing), or ocular myasthenia (a disease causing eye muscle weakness) as statins may sometimes aggravate the condition or lead to the occurrence of myasthenia (see section 4).
- **If you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking Crestor or other related medicines.**
- **If you regularly drink large amounts of alcohol.**

- **If your thyroid gland** is not working properly.
- **If you take other medicines called fibrates** to lower your cholesterol. Please read this leaflet carefully, even if you have taken other medicines for high cholesterol before.
- **If you take medicines used to treat the HIV infection** e.g. ritonavir with lopinavir and/or atazanavir, please see “**Other medicines and Crestor**”.
- **If you are taking or have taken in the last 7 days a medicine called fusidic acid** (a medicine for bacterial infection), orally or by injection. The combination of fusidic acid and Crestor can lead to serious muscle problems (rhabdomyolysis), please see “**Other medicines and Crestor**”.
- **If you are over 70** (as your doctor needs to choose the right start dose of Crestor to suit you)
- **If you have severe respiratory failure.**
- **If you are of Asian origin** – that is Japanese, Chinese, Filipino, Vietnamese, Korean and Indian. Your doctor needs to choose the right start dose of Crestor to suit you.

If any of the above applies to you (or if you are not sure):

- **Do not take Crestor 40 mg (the highest dose) and check with your doctor or pharmacist before you actually start taking any dose of Crestor.**

Serious skin reactions including Stevens-Johnson syndrome and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with Crestor treatment. Stop using Crestor and seek medical attention immediately if you notice any of the symptoms described in section 4.

In a small number of people, statins can affect the liver. This is identified by a simple test which looks for increased levels of liver enzymes in the blood. For this reason, your doctor will usually carry out this blood test (liver function test) before and during treatment with Crestor.

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

#### Children and adolescents

- **If the patient is under 6 years old:** Crestor should not be given to children younger than 6 years.
- **If the patient is below 18 years of age:** The Crestor 40 mg tablet is not suitable for use in children and adolescents below 18 years of age.

#### Other medicines and Crestor

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines

Tell your doctor if you are taking any of the following:

- ciclosporin (used for example, after organ transplants),
- warfarin, clopidogrel **or ticagrelor** (or any other drug used for thinning the blood),
- fibrates (such as gemfibrozil, fenofibrate) or any other medicine used to lower cholesterol (such as ezetimibe),
- indigestion remedies (used to neutralise acid in your stomach),
- erythromycin (an antibiotic), fusidic acid (an antibiotic – please see below and Warnings and precautions),
- an oral contraceptive (the pill),
- regorafenib (used to treat cancer),
- darolutamide (used to treat cancer),
- capmatinib (used to treat cancer),
- hormone replacement therapy,
- fostamatinib (used to treat low platelet counts),
- febuxostat (used to treat and prevent high blood levels of uric acid),
- teriflunomide (used to treat multiple sclerosis),
- any of the following drugs used to treat viral infections, including HIV or hepatitis C infection, alone or in combination (please see Warnings and precautions): ritonavir, lopinavir, atazanavir, sofosbuvir, voxilaprevir, ombitasvir, paritaprevir, dasabuvir, velpatasvir, grazoprevir, elbasvir, glecaprevir, pibrentasvir.

The effects of these medicines could be changed by Crestor or they could change the effect of Crestor.

**If you need to take oral fusidic acid to treat a bacterial infection you will need to temporarily stop using this medicine. Your doctor will tell you when it is safe to restart Crestor. Taking Crestor with fusidic acid may rarely lead to muscle weakness, tenderness or pain (rhabdomyolysis). See more information regarding rhabdomyolysis in Section 4.**

#### Pregnancy and breast-feeding

**Do not take Crestor** if you are pregnant or breast-feeding. If you become pregnant while taking Crestor **stop taking it immediately** and tell your doctor. Women should avoid becoming pregnant while taking Crestor by using suitable contraception.

Ask your doctor or pharmacist for advice before taking any medicine.

#### Driving and using machines

Most people can drive a car and operate machinery while using Crestor – it will not affect their ability. However, some people feel dizzy during treatment with Crestor. If you feel dizzy, consult your doctor before attempting to drive or use machines.

Crestor contains lactose.

If you have been told by your doctor that you have an intolerance to some sugars (lactose or milk sugar), contact your doctor before taking Crestor.

For a full list of ingredients, please see **Contents of the pack and other information.**

#### 3) How to take Crestor

Always take this medicine as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### Usual doses in adults

##### If you are taking Crestor for high cholesterol:

##### Starting dose

Your treatment with Crestor must start with the **5 mg or the 10 mg dose**, even if you have taken a higher dose of a different statin before. The choice of your start dose will depend upon:

- Your cholesterol levels.
- The level of risk you have of experiencing a heart attack or stroke.
- Whether you have a factor that may make you more sensitive to possible side effects.

Please check with your doctor or pharmacist which start dose of Crestor will best suit you.

Your doctor may decide to give you the lowest dose (5 mg) if:

- You are of **Asian origin** (Japanese, Chinese, Filipino, Vietnamese, Korean and Indian).
- You are **over 70 years** of age.
- You have moderate kidney problems.
- You are at risk of muscle aches and pains (myopathy).

#### **Increasing the dose and maximum daily dose**

Your doctor may decide to increase your dose. This is so that you are taking the amount of Crestor that is right for you. If you started with a 5 mg dose, your doctor may decide to double this to 10 mg, then 20 mg and then 40 mg if necessary. If you started on 10 mg, your doctor may decide to double this to 20 mg and then 40 mg if necessary. There will be a gap of four weeks between every dose adjustment.

The maximum daily dose of Crestor is 40 mg. It is only for patients with high cholesterol levels and a high risk of heart attacks or stroke whose cholesterol levels are not lowered enough with 20 mg.

#### **If you are taking Crestor to reduce your risk of having a heart attack, stroke or related health problems:**

The recommended dose is 20 mg daily. However, your doctor may decide to use a lower dose if you have any of the factors mentioned above.

#### **Use in children and adolescents aged 6-17 years**

The dose range in children and adolescents aged 6 to 17 years is 5 to 20 mg once daily. The usual start dose is 5 mg per day, and your doctor may gradually increase your dose to find the right amount of Crestor for you. The maximum daily dose of Crestor is 10 or 20 mg for children aged 6 to 17 years depending on your underlying condition being treated. Take your dose once a day. Crestor **40 mg** tablet should **not** be used by children.

#### **Taking your tablets**

Swallow each tablet whole with a drink of water.

**Take Crestor once daily.** You can take it at any time of the day with or without food.

Try to take your tablet at the same time every day to help you to remember it.

#### **Regular cholesterol checks**

It is important to go back to your doctor for regular cholesterol checks, to make sure your cholesterol has reached and is staying at the correct level. Your doctor may decide to increase your dose so that you are taking the amount of Crestor that is right for you.

#### **If you take more Crestor than you should**

Contact your doctor or nearest hospital for advice.

If you go into hospital or receive treatment for another condition, tell the medical staff that you're taking Crestor.

#### **If you forget to take Crestor**

Don't worry, just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

#### **If you stop taking Crestor**

Talk to your doctor if you want to stop taking Crestor. Your cholesterol levels might increase again if you stop taking Crestor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### **4) Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

It is important that you are aware of what these side effects may be. They are usually mild and disappear after a short time.

**Stop taking Crestor and seek medical help immediately** if you have any of the following allergic reactions:

- Difficulty in breathing, with or without swelling of the face, lips, tongue and/or throat.
- Swelling of the face, lips, tongue and/or throat, which may cause difficulty in swallowing.
- Severe itching of the skin (with raised lumps).
- Reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome).
- Widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome or drug hypersensitivity syndrome).

**Also, stop taking Crestor and talk to your doctor immediately:**

- **If you have any unusual aches or pains** in your muscles which go on for longer than you might expect. Muscle symptoms are more common in children and adolescents than in adults. As with other statins, a very small number of people have experienced unpleasant muscle effects and rarely these have gone on to become a potentially life threatening muscle damage known as *rhabdomyolysis*.
- **If you experience muscle rupture.**
- **If you have lupus-like disease syndrome** (including rash, joint disorders and effects on blood cells).

**Common possible side effects (these may affect between 1 in 10 and 1 in 100 patients):**

- Headache, stomach pain, constipation, feeling sick, muscle pain, feeling weak, dizziness.
- An increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking your Crestor tablets (only Crestor 40 mg).
- Diabetes. This is more likely if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine.

**Uncommon possible side effects (these may affect between 1 in 100 and 1 in 1,000 patients):**

- Rash, itching or other skin reactions.
- An increase in the amount of protein in the urine - this usually returns to normal on its own without having to stop taking your Crestor tablets (only Crestor 5 mg, 10 mg and 20 mg).

**Rare possible side effects (these may affect between 1 in 1,000 and 1 in 10,000 patients):**

- Severe allergic reaction – signs include swelling of the face, lips, tongue and/or throat, difficulty in swallowing and breathing, a severe itching of the skin (with raised lumps). **If you think you are having an allergic reaction, then stop taking Crestor** and seek medical help immediately.
- Muscle damage in adults – as a precaution, stop taking Crestor and talk to your doctor immediately if you have any unusual aches or pains in your muscles which go on for longer than expected.
- A severe stomach pain (inflamed pancreas).

- Increase in liver enzymes in the blood.
- Bleeding or bruising more easily than normal due to low level of blood platelets.
- Lupus-like disease syndrome (including rash, joint disorders and effects on blood cells).

**Very rare possible side effects (these may affect less than 1 in 10,000 patients):**

- Jaundice (yellowing of the skin and eyes), hepatitis (an inflamed liver), traces of blood in your urine, damage to the nerves of your legs and arms (such as numbness), joint pain, memory loss and breast enlargement in men (gynaecomastia).

**Side effects of unknown frequency may include:**

- Diarrhoea (loose stools), cough, shortness of breath, oedema (swelling), sleep disturbances, including insomnia and nightmares, sexual difficulties, depression, breathing problems, including persistent cough and/or shortness of breath or fever, tendon injury and muscle weakness that is constant.
- Myasthenia gravis (a disease causing general muscle weakness including in some cases muscles used when breathing), Ocular myasthenia (a disease causing eye muscle weakness). Talk to your doctor if you experience weakness in your arms or legs that worsens after periods of activity, double vision or drooping of your eyelids, difficulty swallowing, or shortness of breath.

**Reporting of side effects:**

If you get any side effects talk to your doctor or pharmacist. This includes any side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

### **5) How to store Crestor**

- Keep all medicines out of the sight and reach of children.
- Do not take Crestor after the expiry date printed on the carton, label or blister strip. The expiry date refers to the last day of the month.
- Do not store above 30°C. Store in the original package in order to protect from moisture.
- If the tablets become discoloured or show any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.
- Medicines should not be thrown in your bin or put down the drain. If they are out of date, or no longer suitable for you, return them to your pharmacist for safe disposal.

### **6) Contents of the pack and other information**

**What Crestor contains:**

Your medicine is called Crestor. The active substance in Crestor tablets is rosuvastatin. Crestor film-coated tablets contain rosuvastatin calcium equivalent to 5mg, 10 mg, 20mg or 40 mg of rosuvastatin.

The other ingredients are: lactose monohydrate, microcrystalline cellulose, calcium phosphate, crospovidone, magnesium stearate, hypromellose, triacetin, titanium dioxide (E171). Crestor 10mg, 20mg and 40mg film-coated tablets also contain red iron oxide (E172). Crestor 5mg film-coated tablets also contain yellow iron oxide (E172).

**What Crestor looks like and contents of the pack**

Crestor comes in calendar blister packs containing 28 tablets.

Crestor comes in four tablet strengths:

- **Crestor 5 mg film-coated tablets** are round, yellow and marked 'ZD4522 5' on one face, plain on the other.
- **Crestor 10 mg film-coated tablets** are round, pink and marked 'ZD4522 10' on one face, plain on the other.
- **Crestor 20 mg film coated tablets** are round, pink and marked 'ZD4522 20' on one face, plain on the other.
- **Crestor 40 mg film-coated tablets** are oval, pink, oval and marked 'ZD4522' on one face and '40' on the other.

PL 46420/0124	Crestor 5mg Film-coated Tablets	<b>POM</b>
PL 46420/0125	Crestor 10mg Film-coated Tablets	
PL 46420/0126	Crestor 20mg Film-coated Tablets	
PL 46420/0127	Crestor 40mg Film-coated Tablets	

**Who makes and repackages your medicine?**

Your medicine is manufactured by AstraZeneca UK Ltd, Silk Road Business Park, Macclesfield, Cheshire, SK102NA, UK or AstraZeneca GmbH, Tinsdaler Weg 183, 22880, Wedel, Germany or AstraZeneca Reims, Parc Industriel Pompelle, Chemin de Vrilly, Box 1050, Reims Cedex 2, France or AstraZeneca AB, S-151085, Södertälje, Sweden or AndersonBrecon (UK) Ltd, 2-7 Wye Valley Business Park, Brecon Road, Hay-on-Wye, Hereford, HR3 5PG, UK. Procured from within the EU and repackaged by the Product Licence holder: Suerte Pharma Ltd, 4/5 Northolt Trading Estate, Belvue Road, Northolt, Middlesex, UB5 5QS.

**Leaflet date: 19.12.2023**

Crestor® is a registered trademark holder of IPR Pharmaceuticals Inc., Puerto Rico.

HEART UK

For more information about Cholesterol and health contact HEART UK The Cholesterol Charity

HEART UK Cholesterol Helpline

T: 0345 450 5988 Monday-Friday 10am-3pm

E: [ask@heartuk.org.uk](mailto:ask@heartuk.org.uk)

HEART UK Website

W: [www.heartuk.org.uk](http://www.heartuk.org.uk)

Patient Community

W: [www.healthunlocked.com/heartuk](http://www.healthunlocked.com/heartuk)

**For any information about this medicine, or to request a copy of this leaflet in Braille, large print or audio please call the PL holder: Suerte Pharma Limited on 020 8839 3000**