

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Tildiem Retard 120mg Prolonged-Release Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 120 mg of the active substance diltiazem hydrochloride.
Also contains: 69.4 mg of sucrose.

For full list of excipients, see section 6.1

3 PHARMACEUTICAL FORM

Prolonged-release tablet.

White to off-white, round convex coated tablet.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Mild to moderate hypertension and angina pectoris.

4.2 Posology and method of administration

Tildiem Retard tablets should be swallowed whole with a little water and not crushed or chewed.

Patients should be advised that the tablet membrane may pass through the gastrointestinal tract unchanged.

Tildiem (diltiazem hydrochloride) is available in a range of presentations to enable dosage to be adjusted to meet the individual requirements of the patient. Careful titration of the dose should be considered where appropriate, as individual patient response may vary. When changing from one type of Tildiem formulation to another it may be necessary to adjust the dosage until a satisfactory response is obtained. To ensure consistency of response once established, particularly in the sustained release formulations, Tildiem Retard 90 mg and 120 mg should continue to be prescribed by

brand name.

Adults

Angina and hypertension:

The usual starting dose is one tablet (90 mg or 120 mg) twice daily. Patient responses may vary, and dosage requirements can differ significantly between individual patients. Higher divided doses up to 480 mg/day have been used with benefit in some angina patients especially in unstable angina. Doses of 360 mg/day may be required to provide adequate BP control in hypertensive patients.

Elderly and patients with impaired hepatic or renal function

Heart rate should be monitored in these patients and if it falls below 50 beats per minute the dose should not be increased.

Angina:

The recommended starting dose is one Tildiem 60 mg tablet twice daily. This dose may be increased to one 90 mg or 120 mg Tildiem Retard tablet twice daily.

Hypertension:

The starting dose should be one 120 mg Tildiem Retard tablet daily. Dose adjustment to one 90 mg or one 120 mg Tildiem Retard tablet twice daily may be required.

Paediatric population

Safety and efficacy in children have not been established. Therefore, diltiazem is not recommended for use in children.

4.3 Contraindications

- Hypersensitivity to diltiazem or to any of the excipients listed in section 6.1.
- Sick sinus syndrome, 2nd or 3rd degree AV block in patients without a functioning pacemaker.
- Severe bradycardia (less than 50 beats per minute).
- Left ventricular failure with pulmonary stasis.
- Lactation.
- Concurrent use with dantrolene infusion (see section 4.5).
- Combination with ivabradine (see section 4.5).
- Concurrent use with lomitapide (see section 4.5).
- Concurrent use with asunaprevir (see section 4.5).

4.4 Special warnings and precautions for use

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

Close observation is necessary in patients with reduced left ventricular function, bradycardia (risk of exacerbation) or with a 1st degree AV block or prolonged PR interval detected on the electrocardiogram (risk of exacerbation and rarely, of complete block).

Increase of plasma concentrations of diltiazem may be observed in the elderly and

patients with renal or hepatic insufficiency. The contraindications and precautions should be carefully observed and close monitoring, particularly of heart rate, should be carried out at the beginning of treatment.

Cases of acute renal failure secondary to decreased renal perfusion have been reported in patients with existing cardiac disease especially reduced left ventricular function, severe bradycardia or severe hypotension. Careful monitoring of renal function is advised.

In the case of general anaesthesia, the anaesthetist must be informed that the patient is taking diltiazem. The depression of cardiac contractility, conductivity and automaticity as well as the vascular dilatation associated with anaesthetics may be potentiated by calcium channel blockers.

Treatment with diltiazem may be associated with mood changes, including depression (see section 4.5 and 4.8). Early recognition of relevant symptoms is important, especially in predisposed patients. In such cases, drug discontinuation should be considered.

Diltiazem has an inhibitory effect on intestinal motility. Therefore, it should be used with caution in patients at risk of developing an intestinal obstruction.

Careful monitoring is necessary in patients with latent or manifest diabetes mellitus due to a possible increase in blood glucose.

The use of diltiazem may induce bronchospasm, including asthma aggravation, especially in patients with preexisting bronchial hyper-reactivity. Cases have also been reported after dose increase. Patients should be monitored for signs and symptoms of respiratory impairment during diltiazem therapy.

This medicine contains less than 1 mmol sodium (23 mg) per tablet that is to say essentially 'sodium-free'.

Caution should be exercised when direct oral anticoagulants (DOACs) are co-administered with Diltiazem which is a moderate CYP3A4 and a weak P-gp inhibitor, particularly in patients at high risk of bleeding (see Interactions).

4.5 Interaction with other medicinal products and other forms of interaction

Combination Contraindicated for Safety Reasons:

Dantrolene (infusion)

Lethal ventricular fibrillation is regularly observed in animals when intravenous verapamil and dantrolene are administered concomitantly.

The combination of a calcium antagonist and dantrolene is therefore potentially dangerous (see section 4.3).

Ivabradine

Concomitant use with ivabradine is contraindicated due to the additional heart rate lowering effect of diltiazem to ivabradine (see section 4.3)

Lomitapide

Diltiazem (a moderate CYP3A4 inhibitor) may increase lomitapide plasma concentrations through CYP3A4 inhibition leading to increased risk of elevations in liver enzymes (see section 4.3).

Asunaprevir

Diltiazem (a moderate CYP3A4 inhibitor) may increase asunaprevir plasma concentrations through CYP3A4 inhibition (see section 4.3).

Combinations Requiring Caution:

Alpha-antagonists

Increased anti-hypertensive effects. Concomitant treatment with alpha- antagonists may produce or aggravate hypotension. The combination of diltiazem with an alpha antagonist should be considered only with strict monitoring of blood pressure.

Beta-blockers

Possibility of rhythm disturbances (pronounced bradycardia, sinus arrest), sino-atrial and atrio-ventricular conduction disturbances and heart failure (synergistic effect).

Such a combination must only be used under close clinical and ECG monitoring, particularly at the beginning of treatment.

An increased risk of depression has been reported when diltiazem is co-administered with beta-blockers (see section 4.8).

Amiodarone, Digoxin

Increased risk of bradycardia; caution is required when these are combined with diltiazem, particularly in elderly subjects and when high doses are used.

Antiarrhythmic agents

Since diltiazem has antiarrhythmic properties, its concomitant prescription with other antiarrhythmic agents is not recommended due to the risk of increased cardiac adverse effects due to an additive effect. This combination should only be used under close clinical and ECG monitoring.

Nitrate derivatives

Increased hypotensive effects and faintness (additive vasodilating effects). In all patients treated with calcium antagonists, the prescription of nitrate derivatives should only be carried out at gradually increasing doses.

Ciclosporin

Increase in circulating ciclosporin levels. It is recommended that the ciclosporin dose be reduced, renal function be monitored, circulating ciclosporin levels be assayed and that the dose should be adjusted during combined therapy and after its discontinuation.

Phenytoin

When co-administered with phenytoin, diltiazem may increase phenytoin plasma concentration.

It is recommended that the phenytoin plasma concentrations be monitored

X-Ray Contrast Media

Cardiovascular effects of an intravenous bolus of an ionic X-ray contrast media, such as hypotension, may be increased in patients treated with diltiazem.

Special caution is required in patients who concomitantly receive diltiazem and X-ray contrast media

Carbamazepine

Increase in circulating carbamazepine levels. It is recommended that the plasma carbamazepine concentrations be assayed and that the dose should be adjusted if necessary.

Theophylline

Increase in circulating theophylline levels.

Anti-H2 agents (cimetidine and ranitidine)

Increase in plasma diltiazem concentrations. Patients currently receiving diltiazem therapy should be carefully monitored when initiating or discontinuing therapy with anti-H2 agents. An adjustment in diltiazem daily dose may be necessary.

Rifampicin

Risk of decrease of diltiazem plasma levels after initiating therapy with rifampicin. The patient should be carefully monitored when initiating or discontinuing rifampicin treatment.

Lithium

Risk of increase in lithium-induced neurotoxicity.

Antiplatelet drugs

In a pharmacodynamic study, diltiazem was shown to inhibit platelet aggregation. Although the clinical significance of this finding is unknown, potential additive effects when used with antiplatelet drugs should be considered.

Combinations to be Taken into Account:

Diltiazem is metabolised by CYP3A4. A moderate (less than 2-fold) increase of diltiazem plasma concentration in cases of co-administration with a stronger CYP3A4 inhibitor has been documented. Grapefruit juice may increase diltiazem exposure (1.2-fold). Patients who consume grapefruit juice should be monitored for increased adverse effects of diltiazem. Grapefruit juice should be avoided if an interaction is suspected. Diltiazem is also a CYP3A4 isoform inhibitor. Co-administration with other CYP3A4 substrates may result in an increase in plasma concentration of either co-administered drug. Co-administration of diltiazem with a CYP3A4 inducer may result in a decrease of diltiazem plasma concentrations.

Statins

Diltiazem is an inhibitor of CYP3A4 and has been shown to significantly increase the AUC of some statins. The risk of myopathy and rhabdomyolysis is increased by concomitant administration of diltiazem with statins metabolised by CYP3A4 (e.g. atorvastatin, fluvastatin, and simvastatin). An adjustment of the dose of statin may be necessary (see also product information of the relevant statin). When possible, it is recommended to use a statin not metabolised by CYP3A4 (e.g. pravastatin) with diltiazem.

Cilostazol

Inhibition of cilostazol metabolism (CYP3A4). Diltiazem has been shown to increase cilostazol exposure and to enhance its pharmacological activity.

Benzodiazepines (midazolam, triazolam)

Diltiazem significantly increases plasma concentrations of midazolam and triazolam and prolongs their half-life. Special care should be taken when prescribing short-acting benzodiazepines metabolised by the CYP3A4 pathway in patients using diltiazem.

Corticosteroids (methylprednisolone):

Diltiazem can increase methylprednisolone levels (through inhibition of CYP3A4 and possible inhibition of P-glycoprotein). The patient should be monitored when initiating methylprednisolone treatment. An adjustment to the dose of methylprednisolone may be necessary.

Colchicine

Colchicine is a substrate for both CYP3A and the efflux transporter P-glycoprotein (P-gp). Diltiazem is known to inhibit CYP3A and P-gp. When Diltiazem and colchicine are administered together, inhibition of P-gp and/or CYP3A by Diltiazem may lead to increased exposure to colchicine. Combined use is not recommended.

Diltiazem which is a moderate CYP3A4 and weak P-gp inhibitor may increase the plasma concentration of DOACs when co-administered with Diltiazem.

Diltiazem may lead to QT prolongation, when administered with drugs with potential known for prolonging the QT interval. Co-administration of diltiazem with drugs known to prolong the QT interval must be based on a careful assessment of the potential risks and benefits of the treatment.

General Information to be Taken into Account:

Due to the potential for additive effects, caution and careful titration are necessary in patients receiving diltiazem concomitantly with other agents known to affect cardiac contractility and/or conduction.

4.6 Fertility, pregnancy and lactation

Pregnancy: There is very limited data from the use of diltiazem in pregnant patients. Diltiazem has been shown to have reproductive toxicity (see section

5.3) in certain animal species (rat, mice, rabbit). Diltiazem is therefore not recommended during pregnancy, as well as in women of child-bearing potential not using effective contraception.

Breast feeding: as this drug is excreted in breast milk, breast feeding whilst taking diltiazem is contraindicated.

4.7 Effects on ability to drive and use machines

On the basis of reported adverse drug reactions, i.e. dizziness (common), malaise (common), the ability to drive and use machines could be altered. However, no studies have been performed.

4.8 Undesirable effects

The following CIOMS frequency rating is used, when applicable: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $\leq 1/100$); rare ($\geq 1/10,000$ to $\leq 1/1,000$); very rare ($\leq 1/10,000$); not known (cannot be estimated from the available data).

Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

	<i>Very common</i>	<i>Common</i>	<i>Uncommon</i>	<i>Rare</i>	<i>Not known</i>
<i>Blood and lymphatic system disorders</i>					Thrombocytopenia
<i>Psychiatric disorders</i>			Nervousness, insomnia		Mood changes including depression
<i>Nervous system disorders</i>		Headache, dizziness			Extrapyramidal syndrome
<i>Respiratory, thoracic and mediastinal disorders</i>					Bronchospasm (including asthma aggravation)
<i>Cardiac disorders</i>		Atrioventricular block (may be of first, second or third degree; bundle branch block may occur), palpitations	Bradycardia		Sinoatrial block, congestive heart failure, sinus arrest, cardiac arrest (asystole)
<i>Vascular disorders</i>		Flushing	Orthostatic hypotension		Vasculitis (including leukocytoclastic vasculitis)

<i>Gastrointestinal disorders</i>		Constipation, dyspepsia, gastric pain, nausea	Vomiting, diarrhea	Dry mouth	Gingival hyperplasia
<i>Metabolism and nutrition disorders</i>					Hyperglycemia
<i>Hepatobiliary disorders</i>			Hepatic enzymes increase (AST, ALT, LDH, ALP increase)		Hepatitis
<i>Skin and subcutaneous tissue disorders</i>		Erythema		Urticaria	Photosensitivity (including lichenoid keratosis at sun exposed skin areas), angioneurotic oedema, rash, erythema multiforme (including Steven-Johnson's syndrome and toxic epidermal necrolysis), sweating, exfoliative dermatitis, acute generalized exanthematous pustulosis, occasionally desquamative erythema with or without fever, Lupus-like syndrome, Lichenoid drug eruption
<i>Reproductive system and breast disorders</i>					Gynecomastia
<i>General disorders and administration site conditions</i>	Peripheral oedema	Malaise			

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

The clinical effects of acute overdose can involve pronounced hypotension leading to collapse and acute kidney injury, sinus bradycardia with or without isorhythmic dissociation, sinus arrest, atrioventricular conduction disturbances and cardiac arrest.

Non-cardiogenic pulmonary oedema has rarely been reported as a consequence of diltiazem overdose that may manifest with a delayed onset (24-48 hours post-ingestion) and required ventilatory support. Early resuscitative measures (including fluid overload) to maintain perfusion and cardiac output may be precipitating factors.

Treatment, under hospital supervision, will include gastric lavage, osmotic diuresis. Conduction disturbances may be managed by temporary cardiac pacing.

Proposed corrective treatments: atropine, vasopressors, inotropic agents, glucagon and calcium gluconate infusion.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Calcium Channel Blockers; Benzothiazepine derivatives, ATC code: C08DB01

Tildiem is a calcium antagonist. It restricts the slow channel entry of calcium into the cell and so reduces the liberation of calcium from stores in the sarcoplasmic reticulum. This results in a reduction of the amount of available intracellular calcium reducing myocardial oxygen consumption. It increases exercise capacity and improves all indices of myocardial ischaemia in the angina patient. Tildiem relaxes large and small coronary arteries and relieves the spasm of vasospastic (Prinzmetal's) angina and the response to catecholamines but has little effect on the peripheral vasculature. There is therefore no possibility of reflex tachycardia. A small reduction in heart rate occurs which is accompanied by an increase in cardiac output, improved myocardial perfusion and reduction of ventricular work. In animal studies, Tildiem protects the myocardium against the effects of ischaemia and reduces the damage produced by excessive entry of calcium into the myocardial cell during reperfusion.

5.2 Pharmacokinetic properties

Diltiazem is well absorbed (90%) in healthy volunteers following oral administration.

These formulations of diltiazem hydrochloride provide prolonged absorption of the active ingredient. Peak plasma concentrations occur between 4 – 8 hours post-dose.

Bioavailability of this formulation of diltiazem is approximately 90% of that of the conventional tablet. The mean apparent plasma half-life is 7 - 8 hours.

Diltiazem is 80 – 85% bound to plasma proteins. It is extensively metabolised by the liver.

The major circulating metabolite, N-monodesmethyl diltiazem accounts for approximately 35% of the circulating diltiazem.

Less than 5% of diltiazem is excreted unchanged in the urine.

During long term administration to any one patient, plasma concentrations of diltiazem remain constant.

Mean plasma concentrations in elderly subjects and patients with renal and hepatic insufficiency are higher than in young subjects.

Diltiazem and its metabolites are poorly dialysed.

Twice daily formulations of diltiazem have been shown to have different pharmacokinetic profiles and therefore it is not advised to substitute different brands for one another.

5.3 Preclinical safety data

Pregnancy

Reproduction studies have been conducted in mice, rats, and rabbits. Administration of doses ranging from 4 – 6 times (depending on species) the upper limit of the optimum dosage range in clinical trials (480 mg q.d. or 8 mg/kg q.d. for a 60-kg patient) resulted in embryo and fetal lethality. These studies revealed, in one species or another, a propensity to cause fetal abnormalities of the skeleton, heart, retina, and tongue. Also observed were reductions in early individual pup weights, pup survival, as well as prolonged delivery times and an increased incidence of stillbirths.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core:
Sodium dihydrogen citrate
Sucrose
Povidone
Magnesium stearate
Macrogol 6000

Coating:
Sucrose
Coating polymer
Tributyl acetylcitrate
Sodium hydrogen carbonate
Ethyl vanillin
Titanium dioxide (E171)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

Tildiem Retard tablets are coated with a porous polymer membrane which enables the diltiazem to diffuse out of the tablet at a gradual rate. This membrane may pass through the gastro-intestinal tract unchanged. This has no bearing on the efficacy of the product.

6.5 Nature and contents of container

14 or 56 tablets in PVC/foil strips.

14 or 56 tablets in aluminium/(oPA/aluminium/PVC) blisters

56 or 100 tablets in securitainers

56 or 100 tablets in glass bottle

Not all pack sizes may be marketed

6.6 Special precautions for disposal

No special requirements

7 MARKETING AUTHORISATION HOLDER

Aventis Pharma Limited
410 Thames Valley Park Drive
Reading
Berkshire
RG6 1PT
UK

Trading as:

Sanofi
410 Thames Valley Park Drive
Reading
Berkshire
RG6 1PT
UK

8 MARKETING AUTHORISATION NUMBER(S)

PL 04425/0642

**9 DATE OF FIRST AUTHORISATION / RENEWAL OF THE
AUTHORISATION**

Date of first authorisation: 18 April 1991

Date of latest renewal: 28 August 2003

10 DATE OF REVISION OF THE TEXT

12/11/2024