

# Ongentys® 50 mg hard capsules

Ref: 2441/250425/1/F

(opicapone)

## Patient Information Leaflet

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- \* Keep this leaflet. You may need to read it again.
- \* If you have any further questions, ask your doctor or pharmacist.
- \* This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- \* If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Ongentys 50 mg hard capsules but will be referred to as Ongentys throughout the leaflet.

### What is in this leaflet

- 1 What Ongentys is and what it is used for
- 2 What you need to know before you take Ongentys
- 3 How to take Ongentys
- 4 Possible side effects
- 5 How to store Ongentys
- 6 Contents of the pack and other information

## 1 What Ongentys is and what it is used for

Ongentys contains the active substance opicapone. It is used to treat Parkinson's disease and associated movement problems. Parkinson's disease is a progressive disease of the nervous system that causes shaking and affects your movement.

Ongentys is for use in adults who are already taking medicines containing levodopa and DOPA decarboxylase inhibitors. It increases the effects of levodopa and helps to relieve the symptoms of Parkinson's disease and movement problems.

## 2 What you need to know before you take Ongentys

### Do not take Ongentys:

- \* if you are allergic to opicapone or any of the other ingredients of this medicine (listed in section 6);
- \* if you have a tumour of the adrenal gland (known as pheochromocytoma), or of the nervous system (known as paraganglioma), or any other tumour which increase the risk of severe high blood pressure;
- \* if you have ever suffered from neuroleptic malignant syndrome which is a rare reaction to antipsychotic medicines;
- \* if you have ever suffered from a rare muscle disorder called rhabdomyolysis which was not caused by injury;
- \* if you are taking certain antidepressants called monoamine-oxidase (MAO) inhibitors (e.g. phenelzine, tranylcypromine or moclobemide). Ask your doctor or pharmacist if you can take your antidepressant together with Ongentys.

### Warnings and precautions

Talk to your doctor or pharmacist before taking Ongentys:

- \* if you have severe liver problems and suffered from loss of appetite, weight loss, weakness, or exhaustion within a short period of time. Your doctor may need to reconsider your treatment.

**Talk to your doctor or pharmacist** if you or your family/carer notices you are **developing urges or cravings to behave in ways that are unusual for you** or you cannot resist the impulse, drive or temptation to carry out certain activities that could harm you or others. These behaviours are called 'impulse control disorders' and can include: addictive gambling, an abnormally high sex drive or an increased preoccupation with sexual thoughts or feelings. Behaviours such as these have been reported in patients using other medicines for Parkinson's disease. Your doctor may need to review your treatments.

### Children and adolescents

Children and adolescents under the age of 18 years must not take this medicine. It has not been studied in these age groups since treatment of Parkinson's disease is not relevant in children and adolescents.

### Other medicines and Ongentys

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Tell your doctor if you are taking:

- \* medicines for depression or anxiety such as venlafaxine, maprotiline and desipramine. Taking Ongentys with these medicines may increase the risk of side effects. Your doctor may need to adjust your treatment;
- \* safinamide used for Parkinson's disease. There is no experience taking Ongentys and safinamide together. Your doctor may need to adjust your treatment;
- \* medicines to treat asthma such as rimiterole or isoprenaline. Ongentys may increase their effect;
- \* medicines used to treat allergic reactions such as adrenaline. Ongentys may increase their effect;
- \* medicines used to treat heart failure such as dobutamine, dopamine or dopexamine. Ongentys may increase their effects;
- \* medicines for high cholesterol such as rosuvastatin, simvastatin, atorvastatin or pravastatin Ongentys may increase their effects;
- \* medicines that affect the **immune system** such as methotrexate. Ongentys may increase its effect
- \* medicines containing **quinidine**, a medicine used to treat abnormal heart rhythms or malaria. Taking Ongentys and quinidine together, i.e. at the same time, may decrease the effect of Ongentys.

### Pregnancy and breast-feeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Ongentys is not recommended if you are pregnant. You should use effective contraception if you might become pregnant.

It is not known if Ongentys passes into breast milk in humans. Since the risk to the baby/infant cannot be excluded, you should stop breast-feeding during treatment with Ongentys.

### Driving and using machines

Ongentys taken with levodopa may make you feel **light-headed**, dizzy, or sleepy.

**Do not** drive or operate machinery if you experience any of these side effects.

### Ongentys contains lactose and sodium

- \* Lactose: If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicine.
- \* Sodium: This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

## 3 How to take Ongentys

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 50 mg once daily.

Ongentys should be taken preferably at bedtime.

Take Ongentys at least one hour before or after taking your levodopa medicine.

### Doses of other medicines to treat Parkinson's disease

The dose of other medicines to treat Parkinson's disease may need to be adjusted when you start taking Ongentys. Follow the instructions that your doctor has given you.

### Method of administration

Ongentys is for oral use.

Swallow the capsule whole with a glass of water.

### If you take more Ongentys than you should

If you take more Ongentys than you should, tell your doctor or pharmacist, or go to a hospital immediately. Take the medicine package and this leaflet with you. This will help the doctor identify what you have taken.

### If you forget to take Ongentys

If you forget to take one dose, you should continue the treatment and take the next dose as scheduled.

**Do not** take a double dose to make up for a forgotten dose.

# Ongentys® 50 mg hard capsules

(opicapone)

## Patient Information Leaflet (continued)

### If you stop taking Ongentys

**Do not** stop taking Ongentys unless your doctor tells you to as your symptoms may get worse.

If you stop taking Ongentys your doctor may need to adjust the dose of other medicines that you are taking to treat Parkinson's disease.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects caused by Ongentys are usually mild to moderate and occur mostly within the first weeks of treatment. Some side effects may be caused by the increased effects of using Ongentys together with levodopa.

**Contact your doctor straight away** if you experience any side effects at the start of treatment. Many of the side effects can be managed by your doctor adjusting your levodopa medicine.

**Tell your doctor as soon as possible** if you notice any of the following side effects: Very common: may affect more than 1 in 10 people  
\* involuntary and uncontrollable, or difficult or painful body movements

Common: may affect up to 1 in 10 people

- \* constipation
- \* dry mouth
- \* feeling sick (nausea)
- \* vomiting (being sick)
- \* increased levels of the enzyme (creatinase) in your blood
- \* muscle spasm
- \* dizziness
- \* headache
- \* sleepiness
- \* difficulty falling or staying asleep
- \* strange dreams
- \* experiencing or seeing things which do not exist (hallucinations)
- \* a fall in blood pressure on standing up which causes dizziness, light-headedness or fainting

Uncommon: may affect up to 1 in 100 people

- \* palpitations or irregular heartbeat
- \* blocked ear
- \* dry eye
- \* pain or swelling of the abdomen
- \* indigestion
- \* weight loss
- \* loss of appetite
- \* increased levels of triglycerides (fats) in your blood
- \* muscle twitching, stiffness or pain
- \* pain in arms or legs
- \* altered sense of taste
- \* excessive body movements
- \* fainting
- \* anxiety
- \* depression
- \* hearing things which do not exist
- \* nightmares
- \* sleep disorder
- \* abnormal colour of urine
- \* need to wake and pass urine at night
- \* shortness of breath
- \* high or low blood pressure
- \* having falls
- \* feeling low in energy or tired

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5 How to store Ongentys

### Keep out of the sight and reach of children.

This medicine does not require any special temperature storage conditions. Blisters: Store in the original blister in order to protect from moisture. Do not use this medicine after the expiry date which is stated on the blister label and carton, after 'EXP'. The expiry date refers to the last day of that month.

If your doctor tells you to stop taking this medicine, take any remaining medicine back to the pharmacist for safe disposal. Only keep this medicine if your doctor advises you to.

If your medicine becomes discoloured or shows any other signs of deterioration, consult your pharmacist who will tell you what to do.

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6 Contents of the pack and other information

### What Ongentys contains

- \* The active substance is opicapone. Each hard capsule contains 50 mg of opicapone.
- \* The other ingredients are:
  - \* capsule content: lactose monohydrate, sodium starch glycolate (Type A), pregelatinized maize starch and magnesium stearate
  - \* capsule shell: gelatine, indigo carmine aluminium lake (E 132), erythrosine (E 127) and titanium dioxide (E 171)
  - \* printing ink: shellac, titanium dioxide (E 171), propylene glycol, ammonia solution, concentrated, simeticone

### What Ongentys looks like and contents of the pack

Ongentys 50 mg hard capsules are dark blue, approximately 19 mm length, with "OPC 50" and "Bial" printed on the capsules.

Ongentys is available in packs of 30 hard capsules.

### Manufacturer and Licence Holder

Manufactured by Bial - Portela & C<sup>a</sup>, S.A., À Av. da Siderurgia Nacional, 4745-457 S. Mamede do Coronado, Portugal and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

For any information about this medicine, please contact the Product Licence Holder: Lexon UK limited, Tel: 01527 505414.

**POM**

PL: 15184/2441 Ongentys 50 mg hard capsules

This leaflet was last revised in 25/04/25

Ongentys is a registered trademark of BIAL Portela & C<sup>a</sup>, S.A.

Blind or partially sighted?  
Is this leaflet hard to see or read?  
Phone Lexon (UK) Limited,  
Tel: 01527 505414 to obtain the leaflet  
in a format suitable for you