

Betmiga® 50 mg prolonged-release tablets (mirabegron)

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of this product is Betmiga® mg prolonged release tablets but will be referred to as Betmiga throughout the rest of the leaflet

What is in this leaflet:

1. What Betmiga is and what it is used for
2. What you need to know before you take Betmiga
3. How to take Betmiga
4. Possible side effects
5. How to store Betmiga
6. Contents of the pack and other information

1. What Betmiga is and what it is used for

Betmiga contains the active substance mirabegron. It is a bladder muscle relaxant (a so called beta 3-adrenoceptor agonist), which reduces the activity of an overactive bladder and treats the related symptoms.

Betmiga is used to treat the symptoms of an overactive bladder in adults such as:

- suddenly needing to empty your bladder (called urgency)
- having to empty your bladder more than usual (called increased urinary frequency)
- not being able to control when to empty your bladder (called urgency incontinence)

2. What you need to know before you take Betmiga

Do not take Betmiga:

- if you are allergic to mirabegron or any of the other ingredients of this medicine (listed in section 6)
- if you have very high uncontrolled blood pressure.

Warnings and precautions

Talk to your doctor or pharmacist before taking Betmiga:

- if you have trouble emptying your bladder or you have a weak urine stream or if you take other medicines for the treatment of overactive bladder such as anticholinergic medicines.
- if you have kidney or liver problems. Your doctor may need to reduce your dose or may tell you not to take Betmiga, especially if you are taking other medicines such as itraconazole, ketoconazole (fungal infections), ritonavir (HIV/AIDS) or clarithromycin (bacterial infections). Tell your doctor about the medicines that you take.
- if you have an ECG (heart tracing) abnormality known as QT prolongation or you are taking any medicine known to cause this such as
 - o medicines used for abnormal heart rhythm such as quinidine, sotalol, procainamide, ibutilide, flecainide, dofetilide, and amiodarone;
 - o medicines used for allergic rhinitis;
 - o antipsychotic medicines (medicines for mental illness) such as thioridazine, mesoridazine, haloperidol, and chlorpromazine;
 - o anti-infectives such as pentamidine, moxifloxacin, erythromycin, and clarithromycin.

Mirabegron may cause your blood pressure to increase or make your blood pressure worse if you have a history of high blood pressure. It is recommended that your doctor check your blood pressure while you are taking Mirabegron

Children and adolescents

Do not give this medicine to children and adolescents under the age of 18 years because the safety and efficacy of Betmiga in this age group has not been established.

Other medicines and Betmiga

Tell your doctor or pharmacist if you are taking, have recently used or might use any other medicines.

Betmiga may affect the way other medicines work, and other medicines may affect how this medicine works.

- Tell your doctor if you use thioridazine (a medicine for mental illness), propafenone or flecainide (medicines for abnormal heart rhythm), imipramine or desipramine (medicines used for depression). These specific medicines may require dose adjustment by your doctor.
- Tell your doctor if you use digoxin (a medicine for heart failure or abnormal heart rhythm). Blood levels of this medicine are measured by your doctor. If the blood level is out of range, your doctor may adjust the dose of digoxin.
- Tell your doctor if you use dabigatran etexilate (a medicine which is used to reduce the risk of brain or body vessel obstruction by blood clot formation in adult patients with an abnormal heart beat (atrial fibrillation) and additional risk factors). This medicine may require dose adjustment by your doctor.

Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant or are planning to have a baby you should not take Betmiga.

If you are breast feeding, ask your doctor or pharmacist for advice before taking this medicine. It is likely that this medicine passes into your breast milk. You and your doctor should decide if you should take Betmiga or breastfeed. You should not do both.

Driving and using machines

There is no information to suggest that this medicine affects your ability to drive or use machines.

3. How to take Betmiga

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is one 50 mg tablet by mouth once daily. If you have kidney or liver problems, your doctor may need to reduce your dose to one 25 mg tablet by mouth once daily. You should take this medicine with liquids and swallow the tablet whole. Do not crush or chew the tablet. Betmiga can be taken with or without food.

If you take more Betmiga than you should

If you have taken more tablets than you have been told to take, or if someone else accidentally takes your tablets, contact your doctor, pharmacist or hospital for advice immediately.

Symptoms of overdose may include a forceful beating of the heart, an increased pulse rate or an increased blood pressure.

If you forget to take Betmiga

If you forget to take your medicine, take the missed dose as soon as you remember. If it is less than 6 hours before your next scheduled dose, skip the dose and continue to take your medicine at the usual time.

Do not take a double dose to make up for a forgotten dose. If you miss several doses, tell your doctor and follow the advice given to you.

If you stop taking Betmiga

Do not stop treatment with Betmiga early if you do not see an immediate effect. Your bladder might need some time to adapt. You should continue taking your tablets. Do not stop taking them when your bladder condition improves. Stopping treatment may result in recurrence of symptoms of overactive bladder.

Do not stop taking Betmiga without talking to your doctor first, as your overactive bladder symptoms may come back.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects may include irregular heart beat (atrial fibrillation). This is an uncommon side effect (may affect up to 1 in 100 people), but if this side effect occurs, immediately stop taking the medicine and seek urgent medical advice.

If you get headaches, especially sudden, migraine-like (throbbing) headaches, tell your doctor. These may be signs of severely elevated blood pressure.

Other side effects include:

Common side effects (may affect up to 1 in 10 people)

- Increased heart rate (tachycardia)
- Infection of the structures that carry urine (urinary tract infections)
- Nausea
- Constipation
- Headache
- Diarrhoea
- Dizziness

Uncommon side effects (may affect up to 1 in 100 people)

- Bladder infection (cystitis)
- Feeling your heartbeat (palpitations)
- Vaginal infection
- Indigestion (dyspepsia)
- Infection of the stomach (gastritis)
- Swelling of the joints
- Itching of the vulva or vagina (vulvovaginal pruritus)
- Increased blood pressure
- Increase in liver enzymes (GGT, AST and ALT)
- Itching, rash or hives (urticaria, rash, rash macular, rash papular, pruritus)

Rare side effects (may affect up to 1 in 1,000 people)

- Swelling of the eyelid (eyelid oedema)
- Swelling of the lip (lip oedema)
- Swelling of the deeper layers of the skin caused by a build-up of fluid, which can affect any part of the body including the face, tongue or throat and may cause difficulty in breathing (angioedema)
- Small purple spots on the skin (purpura)
- Inflammation of small blood vessels mainly affecting the skin (leukocytoclastic vasculitis)
- Inability to completely empty the bladder (urinary retention)

Very rare side effects (may affect up to 1 in 10,000 people)

- Hypertensive crisis

Not known side effects (frequency cannot be estimated from the available data)

- Insomnia
- Confusion

Betmiga may increase your chances of not being able to empty your bladder if you have bladder outlet obstruction or if you are taking other medicines to treat overactive bladder. Tell your doctor right away if you are unable to empty your bladder.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed on this leaflet.

You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**.

By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Betmiga

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister or bottle after EXP.

The expiry date refers to the last day of that month.

After first opening of the bottle, the tablets can be stored for 6 months. This medicine does not require any special storage conditions.

If the tablets become discoloured or show any signs of deterioration, seek the advice of your pharmacist who will advise you what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Betmiga contains

- The active substance is mirabegron.
Each tablet contains 50 mg of mirabegron.
- The other ingredients are:

Tablet core: Macrogols, hydroxypropylcellulose, butylhydroxytoluene, magnesium stearate

Film-coating: Hypromellose, macrogol, iron oxide yellow (E172)

What Betmiga looks like and contents of the pack

Betmiga 50 mg prolonged release film-coated tablets are oval, yellow film-coated tablets, debossed with the company logo and "355" on the same side.

Betmiga is available in aluminium-aluminium blister in packs containing 30 tablets. Other strengths are available.

Manufacturer

Astellas Pharma Europe B.V. Sylviusweg 62, 2333 BE Leiden
The Netherlands

Delpharm Meppel B.V., Hogemaat 2, 7942 JG Meppel,
The Netherlands

Procured within the EU

Product Licence Holder: Ecosse Pharmaceuticals Limited,
3 Young Place, East Kilbride, G75 0TD, United Kingdom.

Re-packaged by Munro Wholesale Medical Supplies,
3 Young Place, East Kilbride, G75 0TD, United Kingdom.

PL 19065/0655

This leaflet was last revised: 12.08.2025

E0655/3 04.19

POM

If you are blind or partially sighted and require this leaflet in a different format, call 01355 574450 and ask for the Regulatory Department.

Betmiga® is a registered trademark of Astellas Pharma inc.

**Mirabegron Astellas® 50 mg
prolonged-release tablets**
(mirabegron)

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of this product is Mirabegron Astellas 50mg prolonged release tablets but will be referred to as Mirabegron throughout the rest of the leaflet

What is in this leaflet:

1. What Mirabegron is and what it is used for
2. What you need to know before you take Mirabegron
3. How to take Mirabegron
4. Possible side effects
5. How to store Mirabegron
6. Contents of the pack and other information

1. What Mirabegron is and what it is used for

Mirabegron contains the active substance mirabegron. It is a bladder muscle relaxant (a so called beta 3-adrenoceptor agonist), which reduces the activity of an overactive bladder and treats the related symptoms.

Mirabegron is used to treat the symptoms of an overactive bladder in adults such as:

- suddenly needing to empty your bladder (called urgency)
- having to empty your bladder more than usual (called increased urinary frequency)
- not being able to control when to empty your bladder (called urgency incontinence)

2. What you need to know before you take Mirabegron

Do not take Mirabegron:

- if you are allergic to mirabegron or any of the other ingredients of this medicine (listed in section 6)
- if you have very high uncontrolled blood pressure.

Warnings and precautions

Talk to your doctor or pharmacist before taking Mirabegron:

- if you have trouble emptying your bladder or you have a weak urine stream or if you take other medicines for the treatment of overactive bladder such as anticholinergic medicines.
- if you have kidney or liver problems. Your doctor may need to reduce your dose or may tell you not to take Mirabegron, especially if you are taking other medicines such as itraconazole, ketoconazole (fungal infections), ritonavir (HIV/AIDS) or clarithromycin (bacterial infections). Tell your doctor about the medicines that you take.
- if you have an ECG (heart tracing) abnormality known as QT prolongation or you are taking any medicine known to cause this such as
 - o medicines used for abnormal heart rhythm such as quinidine, sotalol, procainamide, ibutilide, flecainide, dofetilide, and amiodarone;
 - o medicines used for allergic rhinitis;
 - o antipsychotic medicines (medicines for mental illness) such as thioridazine, mesoridazine, haloperidol, and chlorpromazine;
 - o anti-infectives such as pentamidine, moxifloxacin, erythromycin, and clarithromycin.

Mirabegron may cause your blood pressure to increase or make your blood pressure worse if you have a history of high blood pressure. It is recommended that your doctor check your blood pressure while you are taking Mirabegron

Children and adolescents

Do not give this medicine to children and adolescents under the age of 18 years because the safety and efficacy of Mirabegron in this age group has not been established.

Other medicines and Mirabegron

Tell your doctor or pharmacist if you are taking, have recently used or might use any other medicines.

Mirabegron may affect the way other medicines work, and other medicines may affect how this medicine works.

- Tell your doctor if you use thioridazine (a medicine for mental illness), propafenone or flecainide (medicines for abnormal heart rhythm), imipramine or desipramine (medicines used for depression). These specific medicines may require dose adjustment by your doctor.
- Tell your doctor if you use digoxin (a medicine for heart failure or abnormal heart rhythm). Blood levels of this medicine are measured by your doctor. If the blood level is out of range, your doctor may adjust the dose of digoxin.
- Tell your doctor if you use dabigatran etexilate (a medicine which is used to reduce the risk of brain or body vessel obstruction by blood clot formation in adult patients with an abnormal heart beat (atrial fibrillation) and additional risk factors). This medicine may require dose adjustment by your doctor.

Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant or are planning to have a baby you should not take Mirabegron.

If you are breast feeding, ask your doctor or pharmacist for advice before taking this medicine. It is likely that this medicine passes into your breast milk. You and your doctor should decide if you should take Mirabegron or breastfeed. You should not do both.

Driving and using machines

There is no information to suggest that this medicine affects your ability to drive or use machines.

3. How to take Mirabegron

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is one 50 mg tablet by mouth once daily. If you have kidney or liver problems, your doctor may need to reduce your dose to one 25 mg tablet by mouth once daily. You should take this medicine with liquids and swallow the tablet whole. Do not crush or chew the tablet. Mirabegron can be taken with or without food.

If you take more Mirabegron than you should

If you have taken more tablets than you have been told to take, or if someone else accidentally takes your tablets, contact your doctor, pharmacist or hospital for advice immediately.

Symptoms of overdose may include a forceful beating of the heart, an increased pulse rate or an increased blood pressure.

If you forget to take Mirabegron

If you forget to take your medicine, take the missed dose as soon as you remember. If it is less than 6 hours before your next scheduled dose, skip the dose and continue to take your medicine at the usual time.

Do not take a double dose to make up for a forgotten dose. If you miss several doses, tell your doctor and follow the advice given to you.

If you stop taking Mirabegron

Do not stop treatment with Mirabegron early if you do not see an immediate effect. Your bladder might need some time to adapt. You should continue taking your tablets. Do not stop taking them when your bladder condition improves. Stopping treatment may result in recurrence of symptoms of overactive bladder.

Do not stop taking Mirabegron without talking to your doctor first, as your overactive bladder symptoms may come back.
If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects may include irregular heart beat (atrial fibrillation). This is an uncommon side effect (may affect up to 1 in 100 people), but if this side effect occurs, immediately stop taking the medicine and seek urgent medical advice.

If you get headaches, especially sudden, migraine-like (throbbing) headaches, tell your doctor. These may be signs of severely elevated blood pressure.

Other side effects include:

Common side effects (may affect up to 1 in 10 people)

- Increased heart rate (tachycardia)
- Infection of the structures that carry urine (urinary tract infections)
- Nausea
- Constipation
- Headache
- Diarrhoea
- Dizziness

Uncommon side effects (may affect up to 1 in 100 people)

- Bladder infection (cystitis)
- Feeling your heartbeat (palpitations)
- Vaginal infection
- Indigestion (dyspepsia)
- Infection of the stomach (gastritis)
- Swelling of the joints
- Itching of the vulva or vagina (vulvovaginal pruritus)
- Increased blood pressure
- Increase in liver enzymes (GGT, AST and ALT)
- Itching, rash or hives (urticaria, rash, rash macular, rash papular, pruritus)

Rare side effects (may affect up to 1 in 1,000 people)

- Swelling of the eyelid (eyelid oedema)
- Swelling of the lip (lip oedema)
- Swelling of the deeper layers of the skin caused by a build-up of fluid, which can affect any part of the body including the face, tongue or throat and may cause difficulty in breathing (angioedema)
- Small purple spots on the skin (purpura)
- Inflammation of small blood vessels mainly affecting the skin (leukocytoclastic vasculitis)
- Inability to completely empty the bladder (urinary retention)

Very rare side effects (may affect up to 1 in 10,000 people)

- Hypertensive crisis

Not known side effects (frequency cannot be estimated from the available data)

- Insomnia
- Confusion

Mirabegron may increase your chances of not being able to empty your bladder if you have bladder outlet obstruction or if you are taking other medicines to treat overactive bladder. Tell your doctor right away if you are unable to empty your bladder.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed on this leaflet.

You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**.

By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Mirabegron

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister or bottle after EXP.

The expiry date refers to the last day of that month.

After first opening of the bottle, the tablets can be stored for 6 months. This medicine does not require any special storage conditions.

If the tablets become discoloured or show any signs of deterioration, seek the advice of your pharmacist who will advise you what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Mirabegron contains

- The active substance is mirabegron.
Each tablet contains 50 mg of mirabegron.
- The other ingredients are:

Tablet core: Macrogols, hydroxypropylcellulose, butylhydroxytoluene, magnesium stearate

Film-coating: Hypromellose, macrogol, iron oxide yellow (E172)

What Mirabegron looks like and contents of the pack

Mirabegron Astellas 50 mg prolonged release film-coated tablets are oval, yellow film-coated tablets, debossed with the company logo and "355" on the same side.

Mirabegron is available in aluminium-aluminium blister in packs containing 30 tablets. Other strengths are available.

Manufacturer

Astellas Pharma Europe B.V. Sylviusweg 62, 2333 BE Leiden
The Netherlands

Delpharm Meppel B.V., Hogemaat 2, 7942 JG Meppel,
The Netherlands

Procured within the EU

Product Licence Holder: Ecosse Pharmaceuticals Limited,
3 Young Place, East Kilbride, G75 0TD, United Kingdom.

Re-packaged by Munro Wholesale Medical Supplies,
3 Young Place, East Kilbride, G75 0TD, United Kingdom.

PL 19065/0655

This leaflet was last revised: 12.08.2025

E0655/3(G) 04.19

POM

If you are blind or partially sighted and require this leaflet in a different format, call 01355 574450 and ask for the Regulatory Department.

Astellas® is the registered trademark of Astellas Pharma Inc