

20 mm 160 mm

290 mm

40 mm 40 mm 40 mm 40 mm 40 mm 40 mm 10 mm

**PACKAGE LEAFLET: INFORMATION FOR THE USER**

# histahive

**Histahive 180 mg Film-Coated Tablets**  
Fexofenadine hydrochloride

**Histahive 180 mg Film-Coated Tablets will also be referred to in this leaflet as Histahive.**  
**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after taking this medicine.

**What is in this leaflet:**

1. What Histahive is and what they are used for
2. What you need to know before you take Histahive
3. How to take Histahive
4. Possible side effects
5. How to store Histahive
6. Contents of the pack and other information

**1. WHAT HISTAHIVE IS AND WHAT THEY ARE USED FOR**

The active ingredient in your tablets is fexofenadine hydrochloride, which belongs to a group of medicines called antihistamines. It is used in adults and adolescents of 12 years and older to relieve the symptoms of urticaria (red, raised itchy skin) which is often known as hives or nettle rash. Histahive is not suitable for children under 12 years of age.

You must talk to a doctor if you do not feel better or if you feel worse after taking this medicine.

**2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE HISTAHIVE**

**Do not take Histahive**

- If you are allergic to fexofenadine or any of the other ingredients of this medicine listed in Section 6.

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Histahive if you:

- have problems with your liver or kidneys
- have or ever had heart problems, as fexofenadine, like other antihistamines may cause your heart to beat faster (tachycardia) or you feel your heart beating (palpitations).
- are elderly.

If any of these apply to you, or if you are not sure, tell your doctor or pharmacist before taking Histahive.

**Other medicines and Histahive**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is especially important for the following medicines:

- Erythromycin (an antibiotic)
- Ketoconazole (a treatment for fungal infections)

If you are taking apalutamide (a medicine to treat prostate cancer), as the effect of fexofenadine may be decreased.

Indigestion remedies containing aluminium and magnesium may affect the action of Histahive, by lowering the amount of medicinal product absorbed.

It is recommended that you leave about 2 hours between the time that you take Histahive and your indigestion remedy.

**Histahive with food and drink**

Histahive should be taken before meals with water.

**Pregnancy and breastfeeding**

If you are pregnant or breastfeeding, think you may be pregnant or planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Histahive should not be taken during pregnancy and whilst breast-feeding unless advised by a doctor.

**Driving and using machines**

Histahive is unlikely to affect your ability to drive or operate machinery. However, you should check that these tablets do not make you feel sleepy or dizzy before driving or operating machinery.

**Histahive contains Allura Red AC Lake, which may cause allergic reactions.**

**Histahive contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

**3. HOW TO TAKE HISTAHIVE**


Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

**For adults and adolescents aged 12 years and over**

The recommended dose is one tablet (180mg) daily. Take your tablet with water before a meal. This medicine starts to relieve your symptoms within 1 hour and lasts for 24 hours.

If the symptoms of allergic skin reactions persist, or gets worse or other symptoms appear, you should stop the treatment and consult a doctor.

**Do not exceed the stated dose. Do not give to adolescents below 12 years of age.**

**Artwork** Dr.Reddy's 

**Version No:** v1.5

**Submission**

**Brand:** Dr. Reddy's  
**Country:** UK  
**Product Name:** Histahive  
**Strength:** 180 mg  
**Form:** Film Coated Tablets  
**Component:** Leaflet  
**Pack Size:** 30s  
**Date Created:** 02 APR 2024  
**Date Modified:** 14 AUG 2024

**Project:** NPL

**Previous DR No:** DRXXXXXX  
**Previous Material Number:** 1500XXX

**Commercial**

**DR No:** XXXXXX  
**Material Code:** 1500xxx  
**Third Party Material Code:** xxxxxx  
**Pharmacode No:** xxxxx

**Implementation Date:** xx xxx xxxx


**Technical Information**

**Die Cut** **Guides**

**Min Font Size:** 9 pt  
**Printer:** India  
**Dimensions:** 160 x 290 mm


**Colours**

**Process Black**

**Dr.Reddy's**   
**Good Health Can't Wait**  
Dr. Reddy's Laboratories (UK) Ltd,  
410 Cambridge Science Park, Milton Road, Cambridge,  
CB4 0PE, United Kingdom

**UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM DR.REDDY'S ARTWORK EU.**

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.



**If you take more Histahive than you should**

If you have accidentally taken more than this leaflet recommended, contact the nearest hospital emergency department or tell your doctor or pharmacist immediately. Remember to take the pack and any remaining tablets with you.

The most common signs of overdose are dizziness, drowsiness, fatigue and dry mouth.

**If you forget to take Histahive**

Do not take a double dose to make up for a forgotten tablet. Take the next dose at the usual time.

**If you stop taking Histahive**

Tell your doctor if you want to stop taking Histahive before you have finished your course of treatment. If you stop taking Histahive earlier than planned, your symptoms may return.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If you get any of the following symptoms after taking these tablets you should contact your doctor or pharmacist immediately as these may be signs of a serious allergic reaction:**

- swelling of the face, lips, tongue or throat and difficulty breathing

The following side effects have also been reported:

**Common side effects** (may affect up to 1 in 10 people):

- headache
- drowsiness
- feeling sick (nausea)
- dizziness.

**Uncommon side effects** (may affect up to 1 in 100 people):

- tiredness
- sleepiness.

**Not known** (frequency not known: cannot be estimated from the available data) which may occur are:

- difficulty sleeping (insomnia)
- sleeping disorders
- bad dreams
- nervousness
- fast or irregular heart beat
- diarrhoea
- skin rash and itching
- hives.
- serious allergic reaction which can cause swelling of the face, lips, tongue or throat, flushing, chest tightness
- difficulty breathing
- blurred vision

**Reporting of side effects**

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

**5. HOW TO STORE HISTAHIVE**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage condition.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. CONTENTS OF THE PACK AND OTHER INFORMATION**

**What Histahive contains**

The active substance is fexofenadine hydrochloride. Each film-coated tablet contains 180mg of fexofenadine hydrochloride; equivalent to 168mg of fexofenadine.

The other ingredients (excipients) are:

Tablet core: powdered cellulose, mannitol (E421), maize starch, croscarmellose sodium, colloidal anhydrous silica, magnesium stearate.

Tablet coating: opadry pink film-coating mixture containing hypromellose (E464), titanium dioxide (E171), macrogol 400, allura red AC lake (FD&C Red #40) (E129), and iron oxide, black (E172)

**What Histahive looks like and contents of the pack**

Histahive Tablets are pink coloured, oval, tablets marked with "EXF" on one side and "180" on the other side.

Histahive Tablets are packed in blister strips packaged in cartons.

The cartons contain 10, 20 or 30 Tablets. Not all packs may be marketed.

**Marketing Authorisation Holder and Manufacturer**

Dr. Reddy's Laboratories (UK) Ltd, 410 Cambridge Science Park, Milton Road, Cambridge, CB4 0PE, UK.

This leaflet was last revised in 07/2024.