

**Package leaflet: Information for the patient
Lamotrigine Medreich 25 mg Tablets
Lamotrigine Medreich 50 mg Tablets
Lamotrigine Medreich 100 mg Tablets
Lamotrigine Medreich 200 mg Tablets**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any of the side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Lamotrigine Medreich tablets are and what they are used for
2. What you need to know before you take Lamotrigine Medreich tablets
3. How to take Lamotrigine Medreich tablets
4. Possible side effects
5. How to store Lamotrigine Medreich tablets
6. Contents of the pack and other information

1. What Lamotrigine Medreich tablets are and what they are used for

Lamotrigine Medreich tablets belongs to a group of medicines called *anti-epileptics*. It is used to treat two conditions - **epilepsy** and **bipolar disorder**.

Lamotrigine Medreich tablets treat epilepsy

by blocking the signals in the brain that trigger epileptic seizures (fits)

- For adults and children aged 13 years and over, Lamotrigine Medreich tablets can be used on its own or with other medicines, to treat epilepsy. Lamotrigine Medreich tablets can also be used with other medicines to treat the seizures that occur with a condition called Lennox-Gastaut syndrome.
- For children aged between 2 and 12 years, Lamotrigine Medreich tablets can be used with other medicines, to treat those conditions. It can be used on its own to treat a type of epilepsy called typical absence seizures.

Lamotrigine Medreich tablets also treat bipolar disorder

People with bipolar disorder (sometimes called *manic depression*) have extreme mood swings, with periods of mania (excitement or euphoria) alternating with periods of depression (deep sadness or despair). For adults aged 18 years and over, Lamotrigine Medreich tablets can be used on its own or with other medicines, to prevent the periods of depression that occur in bipolar disorder. It is not yet known how Lamotrigine Medreich tablets works in the brain to have this effect.

2. What you need to know before you take Lamotrigine Medreich tablets

Do not take Lamotrigine Medreich tablets:

- **If you are allergic (*hypersensitive*)** to lamotrigine or any of the other ingredients of this medicine (listed in Section 6).

If this applies to you:

→ **Tell your doctor** and don't take Lamotrigine Medreich tablets.

Warnings and precautions

Take special care with Lamotrigine Medreich tablets.

Talk to your doctor or pharmacist before taking Lamotrigine Medreich tablets:

- **if you have any kidney problems**
- **if you have ever developed a rash** after taking lamotrigine or other medicines for bipolar disorder or epilepsy
- **if you experience a rash or sunburn after taking lamotrigine and having been exposed to sun or artificial light (e.g. solarium).** Your doctor will check your treatment and may advise you to avoid sunlight or protect yourself against the sun (e.g. use of a sunscreen and/or to wear protective clothing)
- **if you have ever developed meningitis after taking lamotrigine** (*read the description of these symptoms in Section 4 of this leaflet: Rare side effects*)
- **if you are already taking medicine that contains lamotrigine.**
- **if you have a condition called Brugada syndrome or other heart problems.** Brugada syndrome is a genetic disease that results in abnormal electrical activity within the heart. ECG abnormalities which may lead to arrhythmias (abnormal heart rhythm) can be triggered by lamotrigine.

If any of these applies to you:

→ **Tell your doctor**, who may decide to lower the dose or that Lamotrigine Medreich tablets are not suitable for you.

Important information about potentially life-threatening reactions

A small number of people taking Lamotrigine Medreich tablets get an allergic reaction or potentially life-threatening skin reaction, which may develop into more serious problems if they are not treated. These can include Stevens-Johnson Syndrome (SJS), toxic epidermal necrolysis (TEN) and Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). You need to know the symptoms to look out for while you are taking Lamotrigine Medreich tablets. This risk may be associated with a variant in genes in people from Asian origin (mainly Han Chinese and Thai). If you are of such origin and have been tested previously carrying this genetic variant (HLA B* 1502), discuss this with your doctor before taking Lamotrigine Medreich tablets.

→ **Read the description of these symptoms in Section 4 of this leaflet** under '*Potentially life-threatening reactions: get a doctor's help straight away*'.

Haemophagocytic lymphohistiocytosis (HLH)

There have been reports of a rare but very serious immune system reaction, in patients taking lamotrigine.

→ **Contact your doctor or pharmacist immediately** if you experience any of the following symptoms while taking lamotrigine: fever, rash, neurological symptoms (e.g. shaking or tremor, confusional state, disturbances of brain function).

Thoughts of harming yourself or suicide

Anti-epileptic medicines are used to treat several conditions, including epilepsy and bipolar disorder. People with bipolar disorder can sometimes have thoughts of harming themselves or committing suicide. If you have bipolar disorder, you may be more likely to think like this:

- when you first start treatment
- if you have previously had thoughts about harming yourself or about suicide
- if you are under 25 years old.

If you have distressing thoughts or experiences, or if you notice that you feel worse or develop new symptoms while you're taking Lamotrigine Medreich tablets:

→ **See a doctor as soon as possible or go to the nearest hospital for help.**

You may find it helpful to tell a family member, caregiver or close friend that you can become depressed or have significant changes in mood, and ask them to read this leaflet. You might ask them to tell you if they are worried about your depression or other changes in your behaviour.

A small number of people being treated with anti-epileptics such as Lamotrigine Medreich tablets have also had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

If you're taking Lamotrigine Medreich tablets for epilepsy

The seizures in some types of epilepsy may occasionally become worse or happen more often while you're taking Lamotrigine Medreich tablets. Some patients may experience severe seizures, which may cause serious health problems. If your seizures happen more often or if you experience a severe seizure while you're taking Lamotrigine Medreich tablets:

→ **See a doctor as soon as possible.**

Lamotrigine Medreich tablets should not be given to people aged under 18 years to treat bipolar disorder.

Children and adolescents

Medicines to treat depression and other mental health problems increase the risk of suicidal thoughts and behaviour in children and adolescents aged under 18 years.

Other medicines and Lamotrigine Medreich tablets:

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines including herbal medicines or other medicines bought without a prescription.

It is important to tell your doctor or pharmacist if you are taking any other medicines for epilepsy at the same time as lamotrigine if you are pregnant, think you may be pregnant, or are planning to have a baby (see also the information in Section 2 on pregnancy and breast-feeding).

Your doctor needs to know if you are taking other medicines to treat epilepsy or mental health problems. This is to make sure you take the correct dose of Lamotrigine Medreich tablets. These medicines include:

- **oxcarbazepine, felbamate, gabapentin, levetiracetam, pregabalin, topiramate or zonisamide**, used to treat epilepsy
- **lithium, olanzapine or aripiprazole** used to treat **mental health problems**
- **bupropion**, used to treat **mental health problems** or to **stop smoking**
- **paracetamol**, used to treat **pain** and **fever**

→ **Tell your doctor** if you are taking any of these.

Some medicines interact with Lamotrigine Medreich tablets or make it more likely that people will have side effects. These include:

- **valproate**, used to treat **epilepsy** and **mental health problems**
- **carbamazepine**, used to treat **epilepsy** and **mental health problems**
- **phenytoin, primidone or phenobarbitone**, used to treat **epilepsy**
- **risperidone**, used to treat **mental health problems**
- **rifampicin**, which is an **antibiotic**
- medicines used to treat **Human Immunodeficiency Virus (HIV) infection** (a combination of lopinavir and ritonavir or atazanavir and ritonavir)
- **oestrogen-containing products, including hormonal contraceptives**, such as the **Pill** (*see below*) and **hormonal replacement therapy (HRT)**.

→ **Tell your doctor** if you are taking any of these or if you start or stop taking any.

Hormonal contraceptives (such as the Pill) can affect the way Lamotrigine Medreich tablets works

Your doctor may recommend that you use a particular type of hormonal contraceptive or another method of contraception, such as condoms, a cap or coil. If you are using a hormonal contraceptive like the Pill, your doctor may take samples of your blood to check the level of Lamotrigine Medreich tablets. If you are using a hormonal contraceptive or if you plan to start using one:

→ **Talk to your doctor**, who will discuss suitable methods of contraception with you.

Lamotrigine Medreich tablets can also affect the way hormonal contraceptives work, although it's unlikely to make them less effective. If you are using a hormonal contraceptive and you notice any changes in your menstrual pattern, such as breakthrough bleeding or spotting between periods:

→ **Tell your doctor**. These may be signs that Lamotrigine Medreich tablet is affecting the way your contraceptive is working.

HRT can affect the way Lamotrigine Medreich tablets works

If you are using oestrogen-containing hormone replacement therapy (HRT), your doctor may take samples of your blood to check the level of Lamotrigine Medreich tablets.

Pregnancy and breast-feeding:

→ **If you are pregnant, or think you may be pregnant, you must tell your doctor straight away and discuss possible risks the epilepsy medicine you are taking might pose to your unborn baby.**

→ **If you are planning to become pregnant you should discuss your epilepsy treatment with your doctor as early as possible before you become pregnant.**

- **You should not stop treatment without discussing this with your doctor.** This is particularly important if you have epilepsy.
- Pregnancy may alter the effectiveness of Lamotrigine Medreich tablets, so you may need blood tests and your dose of Lamotrigine Medreich tablets may be adjusted.
- Studies of women who took lamotrigine at usual doses while they were pregnant have not shown an increased risk of physical birth abnormalities.
- The risk of physical birth abnormalities may increase when higher doses of epilepsy medicines are taken or more than one epilepsy medicine is used at the same time. Where possible, your doctor will consider using one epilepsy medicine at the lowest effective dose to control your epilepsy.
- Your doctor may advise you to take **follic acid supplements** if you're planning to become pregnant and while you're pregnant.

→ **If you are breast-feeding or planning to breast-feed ask your doctor or pharmacist for advice before taking this medicine.** The active ingredient of Lamotrigine Medreich tablets passes into breast milk and may affect your baby. Your doctor will discuss the risks and benefits of breast-feeding while you're taking Lamotrigine Medreich tablets and will check your baby from time to time, whether drowsiness, rash or poor weight gain occurs, if you decide to breast-feed. Inform your doctor if you observe any of these symptoms in your baby.

Driving and using machines:

Lamotrigine Medreich tablets can cause dizziness and double vision.

→ **Don't drive or use machines unless you are sure you're not affected.**

If you have epilepsy, talk to your doctor about driving and using machines.

Lamotrigine Medreich tablets contains Lactose:

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Information on Sodium content:

This medicine contains less than 1mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Lamotrigine Medreich tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much Lamotrigine Medreich tablets to take

It may take a while to find the best dose of Lamotrigine Medreich tablets for you. The dose you take will depend on:

This section is continued overleaf →



16.5 mm

6 mm

16.5 mm

50 mm

39 mm

- your age
- whether you are taking Lamotrigine Medreich tablets with other medicines
- whether you have any kidney or liver problems.

Your doctor will prescribe a low dose to start and gradually increase the dose over a few weeks until you reach a dose that works for you (called the *effective dose*). **Never take more Lamotrigine Medreich tablet than your doctor tells you to.**

The recommended effective dose of Lamotrigine Medreich tablets for adults and children aged 13 years or over is between 100 mg and 400 mg each day.

For children aged 2 to 12 years, the effective dose depends on their body weight - usually, it's between 1 mg and 15 mg for each kilogram of the child's weight, up to a maximum maintenance dose of 200 mg daily.

Use in children

Lamotrigine Medreich tablets are not recommended for children aged under 2 years.

How to take your dose of Lamotrigine Medreich tablets

Take your dose of Lamotrigine Medreich tablets once or twice a day, as your doctor advises. It can be taken with or without food. Your doctor may also advise you to start or stop taking other medicines, depending on what condition you're being treated for and the way you respond to treatment.

- **Swallow your tablets whole.** Don't break, chew or crush them.
- **Always take the full dose** that your doctor has prescribed. Never take only part of a tablet.

If you take more Lamotrigine Medreich tablets than you should

→ **Contact a doctor or nearest hospital emergency department immediately.** If possible, show them the Lamotrigine Medreich tablets packet.

If you take too much **Lamotrigine Medreich tablets you may be more likely to have serious side effects which may be fatal.**

Someone who has taken too much Lamotrigine Medreich tablets may have any of these symptoms:

- rapid, uncontrollable eye movements (*nystagmus*)
- clumsiness and lack of co-ordination, affecting their balance (*ataxia*)
- heart rhythm changes (detected usually on ECG)
- loss of consciousness, fits (convulsions) or coma.

If you forget to take a single dose of Lamotrigine Medreich tablets

→ **Do not take a double dose to make up for a forgotten dose. Just take your next dose at the usual time.**

In case you forget to take multiple doses of Lamotrigine Medreich tablets

→ **Ask your doctor for advice on how to start taking it again.** It's important that you do this.

Don't stop taking Lamotrigine Medreich tablets without advice

Lamotrigine Medreich tablets must be taken for as long as your doctor recommends. Don't stop unless your doctor advises you to.

If you're taking Lamotrigine Medreich tablets for epilepsy

To stop taking Lamotrigine Medreich tablet, it is **important that the dose is reduced gradually**, over about 2 weeks. If you suddenly stop taking Lamotrigine Medreich tablet, your epilepsy may come back or get worse.

If you're taking Lamotrigine Medreich tablets for bipolar disorder

Lamotrigine Medreich tablets may take some time to work, so you are unlikely to feel better straight away. If you stop taking Lamotrigine Medreich tablets, your dose will not need to be reduced gradually. But you should still talk to your doctor first, if you want to stop taking Lamotrigine Medreich tablets.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, but not everyone gets them.

Potentially life-threatening reactions: get a doctor's help straight away

A small number of people taking Lamotrigine Medreich tablets get an allergic reaction or potentially life-threatening skin reaction, which may develop into more serious problems if they are not treated.

These symptoms are more likely to happen during the first few months of treatment with Lamotrigine Medreich tablets, especially if the starting dose is too high or if the dose is increased too quickly or if Lamotrigine Medreich tablets are taken with another medicine called valproate. Some of the symptoms are more common in children, so parents should be especially careful to watch out for them.

Symptoms of these reactions include:

- **skin rashes or redness**, which may develop into serious or sometimes life-threatening skin reactions including rash with target lesions (*Erythema multiforme*), widespread rash with blisters and peeling skin, particularly occurring around the mouth, nose, eyes and genitals (*Stevens-Johnson syndrome*), extensive peeling of the skin (more than 30% of the body surface - *toxic epidermal necrolysis*) or extended rashes with liver, blood and other body organs involvement (Drug Reaction with Eosinophilia and Systemic Symptoms which is also known as DRESS hypersensitivity syndrome)
- **ulcers in the mouth, throat, nose or genitals**
- **a sore mouth or red or swollen eyes** (*conjunctivitis*)
- **a high temperature** (*fever*), flu-like symptoms or drowsiness
- **swelling around your face** or **swollen glands** in your neck, armpit or groin
- **unexpected bleeding or bruising**, or the fingers turning blue
- **a sore throat** or more infections (such as colds) than usual
- increased levels of liver enzymes seen in blood tests
- an increase in a type of white blood cell (eosinophils)
- enlarged lymph nodes
- involvement of the organs of the body including liver and kidneys.

In many cases, these symptoms will be signs of less serious side effects **but you must be aware that they are potentially life-threatening and can develop into more serious problems**, such as organ failure, if they are not treated. If you notice any of these symptoms:

→ **Contact a doctor immediately.** Your doctor may decide to carry out tests on your liver, kidneys or blood and may tell you to stop taking Lamotrigine Medreich tablets.

In case you have developed Stevens-Johnson syndrome or toxic epidermal necrolysis your doctor will tell you that you must never use lamotrigine again.

Haemophagocytic lymphohistiocytosis (HLH)

(see section 2: What you need to know before you take Lamotrigine Medreich tablets).

Very common: may affect more than 1 in 10 people

- headache
- Skin rash.

Common: may affect up to 1 in 10 people

- aggression or irritability
- feeling sleepy or drowsy
- feeling dizzy
- shaking or tremors
- difficulty in sleeping (*insomnia*)
- feeling agitated
- diarrhoea
- dry mouth
- feeling sick (*nausea*) or being sick (*vomiting*)
- feeling tired
- pain in your back or joints, or elsewhere.

Uncommon: may affect up to 1 in 100 people

- clumsiness and lack of co-ordination (*ataxia*)
- double vision or blurred vision
- unusual hair loss or thinning (*alopecia*)
- skin rash or sunburn after exposure to sun or artificial light (*photosensitivity*)

Rare: may affect up to 1 in 1,000 people

- skin reaction that causes red spots or patches on the skin, that may look like a target or "bulls-eye" with a dark red centre surrounded by paler red rings (*erythema multiforme*)
- a life-threatening skin reaction (*Stevens-Johnson syndrome*): (see also the information at the beginning of Section 4)
- a group of symptoms together including: fever, nausea, vomiting, headache, stiff neck and extreme sensitivity to bright light. This may be caused by an inflammation of the membranes that cover the brain and spinal cord (*meningitis*). These symptoms usually disappear once treatment is stopped however if the symptoms continue or get worse **contact your doctor**
- rapid, uncontrollable eye movements (*nystagmus*)
- itchy eyes, with discharge and crusty eyelids (*conjunctivitis*).

Very rare: may affect up to 1 in 10,000 people

- a life-threatening skin reaction (*toxic epidermal necrolysis*): see also the information at the beginning of Section 4
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS): (see also the information at the beginning of Section 4)
- a high temperature (*fever*): (see also the information at the beginning of Section 4)
- swelling around the face (*oedema*) or swollen glands in the neck, armpit or groin (*lymphadenopathy*): (see also the information at the beginning of Section 4)
- changes in liver function, which will show up in blood tests, or liver failure: (see also the information at the beginning of Section 4)
- a serious disorder of blood clotting, which can cause unexpected bleeding or bruising (*disseminated intravascular coagulation*): (see also the information at the beginning of Section 4)
- haemophagocytic lymphohistiocytosis (HLH) (see Section 2: What you need to know before you take Lamotrigine Medreich tablets).
- changes which may show up in blood tests - including reduced numbers of red blood cells (*anaemia*), reduced numbers of white blood cells (*leucopenia*, *neutropenia*, *agranulocytosis*), reduced numbers of platelets (*thrombocytopenia*), reduced numbers of all these types of cell (*pancytopenia*) and a disorder of the bone marrow called aplastic anaemia
- hallucinations ('seeing' or 'hearing' things that aren't really there)
- confusion
- feeling 'wobbly' or unsteady when you move about
- uncontrollable repeated body movements and/or sounds or words (*tics*), uncontrollable muscle spasms affecting the eyes, head and torso (*choreoathetosis*) or other unusual body movements such as jerking, shaking or stiffness
- in people who already have epilepsy, seizures happening more often
- in people who already have Parkinson's disease, worsening of the symptoms.
- lupus-like reaction (symptoms may include: back or joint pain which sometimes may be accompanied by fever and/or general ill health).

Other side effects

Other side effects have occurred in a small number of people but their exact frequency is:

Not known: frequency cannot be estimated from the available data

- There have been reports of bone disorders including osteopenia and osteoporosis (thinning of the bone) and fractures. Check with your doctor or pharmacist if you are on long-term anti-epileptic medication, have a history of osteoporosis or take steroids.
- Inflammation of the kidney (*tubulointerstitial nephritis*), or inflammation of both the kidney and the eye (*tubulointerstitial nephritis and uveitis syndrome*).
- Nightmares.
- Lower immunity because of lower levels of antibodies called immunoglobulins in the blood which help protect against infection.
- Red nodules or patches on the skin (pseudolymphoma).

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard, or search for MHRA Yellow Card in the Google Play or Apple App Store. **By reporting side effects you can help provide more information on the safety of this medicine.**

5. How to store Lamotrigine Medreich tablets

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date shown on the blister or carton. The expiry date refers to the last day of that month.

Store below 25°C.

Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. This will help protect the environment.

6. Contents of the pack and other information

What Lamotrigine Medreich tablets contains:

- The active substance is lamotrigine. Each tablet contains 25 mg, 50 mg, 100 mg or 200 mg lamotrigine.
- The other ingredients are: lactose monohydrate, microcrystalline cellulose, sodium starch glycoate (Type A), iron oxide yellow (E172), povidone (K-30), magnesium stearate.

What Lamotrigine Medreich tablets looks like and contents of the pack:

Lamotrigine Medreich 25 mg Tablets are Pale yellowish brown, square with rounded corner uncoated tablets and marked with '25' on one side and plain on the other side. (Nominal tablet diameter is 5.00 mm).

Lamotrigine Medreich 50 mg Tablets are Pale yellowish brown, square with rounded corner uncoated tablets and marked with '50' on one side and plain on the other side. (Nominal tablet diameter is: 6.80 mm).

Lamotrigine Medreich 100 mg Tablets are Pale yellowish brown, square with rounded corner uncoated tablets and marked with '100' on one side and plain on the other side. (Nominal tablet diameter is: 8.20 mm).

Lamotrigine Medreich 200 mg Tablets are Pale yellowish brown, square with rounded corner uncoated tablets and marked with '200' on one side and plain on the other side. (Nominal tablet diameter is: 11.40 mm).

Lamotrigine Medreich tablets are available in PVC / Alu blister pack of 10, 14, 28, 30, 42, 56, 90, 98 or 100 tablets and in HDPE bottle packs of 500's and 1000's.

Not all pack sizes may be marketed

MA Holder and Manufacturer:

MEDREICH PLC

Warwick House, Plane Tree Crescent, Feltham TW13 7HF, United Kingdom

This leaflet was last revised in 12/2025.

To request a copy of this leaflet in Braille or large print, please call, 02088311580 (UK only).

Please be ready to give the following information:

Product name	License number
Lamotrigine Medreich 25 mg Tablets	PL 21880/0321
Lamotrigine Medreich 50 mg Tablets	PL 21880/0322
Lamotrigine Medreich 100 mg Tablets	PL 21880/0323
Lamotrigine Medreich 200 mg Tablets	PL 21880/0324

Colours:

 Black

 Red 032 C

Font Type & Size	Times New Roman Bold
	Heading - 10.5 pts
	Body - 9 pts
Version	

MEDREICH LIMITED				Title: Art Work Approval Form			
Product	Lamotrigine Medreich Tablets - S/L - English - PIL SAP DESCR.: LAMOTRIGINE MEDREICH TAB PIL S/L ENG-WOA			Specification:	Printed on 40.5 (±3.5) GSM news print paper		
Customer	MEDREICH PLC, UK			Colours:	2 - Red 032 C and Black		
Reason for Issue	PIF No.: 1130PI2520087 / 88 / 89 / 90 New Product			Dimensions:	135 x 620 mm (Open size) 135 x 39 mm (Folded Size) LxW		
Related FG Codes	130XXXX	Pharmacode No.	3881	No. of Folds (only for PIL)	04	Artwork made to	100%
Item Code	1230044-V1	Blister Layout No.	NA				
		Carton Layout No.	NA				
		PIL Layout No.	NA				
Verified By				Approved By			
PDC:		PDC:		Regulatory:		CQA:	