




Product:	<b>Co-tenidone 50/12.5 mg &amp; 100/25 mg FC Tablets BP</b>	Printing Colours:	Non-Printing Colours:	
MA Holder:	<b>Activase</b>	 Black	 Profile	
PL No.:	<b>28444/0101-0100</b>	Dimensions:	<b>170 x 320 mm</b>	Final Preparation Date For Submission: <b>12/09/2025</b>
Pack Size:	<b>All Packs</b>	AW Code:	<b>A0101-0100/O/PIL2/A4</b>	Packing Site Technical Approval: <b>00/00/0000</b>
Livery:	<b>Activase</b>	Bar Code:	<b>-</b>	Authority Approval Date: <b>00/00/0000</b>
Manufacturer:	<b>-</b>	Font & Size:	<b>Swiss 721 BT, 8.5 pt</b>	Latest Implementation Deadline: <b>00/00/0000</b>
Market:	<b>UK</b>	Created By:	<b>MKumbhar</b>	Print Proof Approval: <b>00/00/0000</b>
Component:	<b>Leaflet</b>	Version:	<b>0.2</b>	

**Prompt list for artwork checks**

- Are the dimensions correct?
- Are the font sizes correct?
- Are the livery and colours correct?
- Is the content (text) correct?
- Is the braille correct?
- Is the artwork code correct?

SPACE FOR  
PHARMA  
CODE  
10mmx10mm

000

PACKAGE LEAFLET: INFORMATION FOR THE PATIENT

## Co-tenidone 50/12.5 mg Film-coated Tablets BP Co-tenidone 100/25 mg Film-coated Tablets BP (atenolol and chlortalidone)

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### WHAT IS IN THIS LEAFLET:

1. What Co-tenidone Tablets are and what they are used for
2. What you need to know before you take Co-tenidone Tablets
3. How to take Co-tenidone Tablets
4. Possible side effects
5. How to store Co-tenidone Tablets
6. Contents of the pack and other information

### 1. WHAT CO-TENIDONE TABLETS ARE AND WHAT THEY ARE USED FOR

Co-tenidone Tablets are used to treat high blood pressure (hypertension). It contains two different medicines: atenolol and chlortalidone. These medicines work together to lower your blood pressure.

- atenolol belongs to a group of medicines called beta-blockers. It works by making your heart beat more slowly and with less force.
- chlortalidone belongs to a group of medicines called diuretics. It works by increasing the amount of urine produced by your kidneys.

### 2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE CO-TENIDONE TABLETS

#### Do not take Co-tenidone Tablets if:

- you are allergic to atenolol, chlortalidone or any of the other ingredients of this medicine (listed in section 6).
- you have ever had any of the following heart problems:
  - heart failure which is not under control (this usually makes you breathless and causes your ankles to swell).
  - second- or third-degree heart block (a condition which may be treated by a pacemaker)
  - very slow or very uneven heart beats, very low blood pressure or very poor circulation.
- you have problems with your kidneys.
- you have a tumour called phaeochromocytoma that is not being treated. This is usually near your kidney and can cause high blood pressure.
- you have been told that you have higher than normal levels of acid in your blood (metabolic acidosis).
- you are pregnant, are trying to become pregnant or are breast-feeding.

Do not take Co-tenidone Tablets if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Co-tenidone Tablets.

#### Warnings and precautions

Talk to your doctor or pharmacist before taking Co-tenidone Tablets if:

- **you have asthma, wheezing or any other similar breathing problems, or you get allergic reactions, for example to insect stings. If you have ever had asthma or wheezing, do not take this medicine without first checking with your doctor.**
- you have a tight, painful feeling in the chest in periods of rest (Prinzmetal's angina).

- you have poor blood circulation or controlled heart failure.
- you have a heart condition, first-degree heart block.
- you have diabetes. Your medicine may change how you respond to having low blood sugar. You may feel your heart beating faster.
- you have thyrotoxicosis (a condition caused by an overactive thyroid gland). Your medicine may hide the symptoms of thyrotoxicosis.
- you have problems with your adrenal glands.
- you have liver problems.
- you experience a decrease in vision or eye pain. These could be symptoms of fluid accumulation in the vascular layer of the eye (choroidal effusion) or an increase of pressure in your eye and can happen within hours to a week of taking Co-tenidone Tablets. This can lead to permanent vision loss, if not treated. If you earlier have had a penicillin or sulfonamide allergy, you can be at higher risk of developing this.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Co-tenidone Tablets.

#### Children

This medicine is not for use in children.

#### Other medicines and Co-tenidone Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines that you buy without a prescription and herbal medicines. This is because Co-tenidone Tablets can affect the way some other medicines work and some medicines can have an effect on Co-tenidone Tablets.

In particular, tell your doctor if you are taking any of the following medicines:

- clonidine (for high blood pressure or migraine). If you are taking clonidine and Co-tenidone Tablets together, do not stop taking clonidine unless your doctor tells you to do so. If you have to stop taking clonidine, your doctor will give you careful instructions about how to do it.
- verapamil, nifedipine and diltiazem (for high blood pressure or chest pain).
- disopyramide or amiodarone (for an uneven heart beat).
- digoxin (for heart problems).
- adrenaline, also known as epinephrine (a medicine that stimulates the heart).
- baclofen (a medicine used for muscle relaxation).
- ibuprofen or indometacin (for pain and inflammation).
- insulin or medicines that you take by mouth for diabetes.
- lithium (for certain mental illnesses).
- medicines to treat nose or sinus congestion or other cold remedies (including those you can buy in the pharmacy).
- betaxolol, carteolol, levobunolol and timolol (eye drops for glaucoma).

#### Operations

If you go into hospital to have an operation, tell the anaesthetist or medical staff that you are taking Co-tenidone Tablets. This is because you can get low blood pressure (hypotension) if you are given certain anaesthetics while you are taking Co-tenidone Tablets.

#### Pregnancy and breast-feeding

Do not take Co-tenidone Tablets if you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Driving and using machines

Your medicine is not likely to affect you being able to drive or use any tools or machines. However, it is best to wait to see how your medicine affects you before trying these activities.

If you feel dizzy or tired when taking this medicine, ask your doctor for advice.

#### Sodium content

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

### 3. HOW TO TAKE CO-TENIDONE TABLETS

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how many tablets to take each day and when to take them. Read the label on the carton to remind you what the doctor said.



The tablets must be swallowed with a drink of water.

The recommended dose is:

#### Adults

One 50/12.5 mg tablet or one 100/25 mg tablet a day.

#### People with kidney problems

If you have problems with your kidneys, your doctor may give you a lower dose or ask you to take this medicine less often.

#### Children

Co-tenidone Tablets are not recommended for children.

#### If you take more Co-tenidone Tablets than you should

If you take too many of these tablets than prescribed by your doctor, contact your doctor or nearest accident and emergency department immediately (take any remaining medicine in its original package with you if you can).

Symptoms of overdose include a slow heart rate, low blood pressure, nausea (feeling sick), weakness, dizziness and imbalance of electrolytes.

#### If you forget to take Co-tenidone Tablets

If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the forgotten dose and just carry on as before.

Do not take a double dose to make up for a forgotten dose.

#### If you stop taking Co-tenidone Tablets

Do not stop taking Co-tenidone Tablets without talking to your doctor. In some cases, you may need to stop taking it gradually.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

#### Allergic reactions:

If you have an allergic reaction, see a doctor straight away. The signs may include raised lumps on your skin (weals) or swelling of your face, lips, mouth, tongue or throat.

#### Other possible side effects:

##### Common (may affect up to 1 in 10 people)

- Cold hands and feet.
- You may notice that your pulse rate becomes slower while you are taking the tablets. This is normal, but if you are concerned please tell your doctor about it.
- Diarrhoea.
- Feeling sick (nausea).
- Feeling tired.
- Changes in the amount of certain substances in your blood. It can cause:
  - higher levels of sugar (glucose) than normal.
  - low levels of sodium, which may cause weakness, being sick (vomiting) and cramps.
  - low levels of potassium.
  - an increase in the amount of uric acid.

Your doctor may take blood samples every so often to check on these levels.

##### Uncommon (may affect up to 1 in 100 people)

- Disturbed sleep.

##### Rare (may affect up to 1 in 1,000 people)

- Heart block (which can cause an abnormal heart beat, dizziness, tiredness or fainting).
- Numbness and spasm in your fingers which is followed by warmth and pain (Raynaud's disease).
- Dizziness (particularly when standing up).
- Headache.
- Feeling confused.
- Changes in personality (psychoses) or hallucinations.
- Mood changes.
- Nightmares.
- Dry mouth.
- Thinning of your hair.
- Dry eyes.
- Skin rash.
- Tingling of your hands.
- Disturbances of vision.
- Bruising more easily or purplish marks on your skin.

- Being unable to get an erection (impotence).
- Jaundice (causing yellowing of your skin or the whites of your eyes).
- Pancreatitis (inflammation of a large gland behind the stomach).
- A reduced number of white blood cells.

#### Very rare (may affect up to 1 in 10,000 people)

- Changes to some of the cells or other parts of your blood. Your doctor may take blood samples every so often to check whether Co-tenidone Tablets has had any effect on your blood.

#### Not known (frequency cannot be estimated from the available data)

- Constipation.
- Lupus-like syndrome (a disease where the immune system produces antibodies that attacks mainly skin and joints).
- Decrease in vision or pain in your eyes due to high pressure (possible signs of fluid accumulation in the vascular layer of the eye (choroidal effusion) or acute angle-closure glaucoma).
- Depression

#### Conditions that may get worse

If you have any of the following conditions, they may get worse when you start to take your medicine. This happens rarely affecting less than 1 in 1,000 people:

- Psoriasis (a skin condition).
- Being short of breath or having swollen ankles (if you have heart failure).
- Asthma or breathing problems.
- Poor circulation.

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. HOW TO STORE CO-TENIDONE TABLETS

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original package to protect from light and moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. CONTENTS OF THE PACK AND OTHER INFORMATION

#### What Co-tenidone Tablets contain

The active substances are atenolol and chlortalidone.

Each Co-tenidone 50/12.5 mg tablet contains 50 mg atenolol and 12.5 mg chlortalidone.

Each Co-tenidone 100/25 mg tablet contains 100 mg atenolol and 25 mg chlortalidone.

The other ingredients are maize starch, calcium hydrogen phosphate dihydrate, microcrystalline cellulose, povidone K30, sodium starch glycolate (type A), and magnesium stearate. The film coating consists of hypromellose (E464), titanium dioxide (E171), macrogol, iron oxide red (E172), iron oxide yellow (E172) and iron oxide black (E172).

#### What Co-tenidone Tablets look like and contents of the pack

The 50/12.5 mg and 100/25 mg tablets are brownish pink, round, biconvex and film-coated, marked CT/50 or CT/100 on one side.

They are available in blister packs containing 28, 30, 56 or 60 tablets. Not all pack sizes may be marketed.

#### Manufacturer

DDSA Pharmaceuticals Limited,  
84 Pembroke Road,  
London, W8 6NX, United Kingdom

#### Marketing Authorisation Holder

Activase Pharmaceuticals Limited,  
11 Boumpoulinas, 3rd floor,  
P.C. 1060, Nicosia, Cyprus

This leaflet was last revised in 09/2025.

A0101-0100/O/PIL2/A4

**Co-tenidone 50/12.5 mg Film-coated Tablets BP**  
**Co-tenidone 100/25 mg Film-coated Tablets BP**

<b>AW Code</b>	<b>Version</b>	<b>Date</b>	<b>Req by</b>	<b>Change Reasons</b>
A0101-0100/O/PIL2/A4	0.1	24/03/2025	VH	PIL update in line with brand leader.
A0101-0100/O/PIL2/A4	0.2	12/09/2025	VH	RFI update