

Propofol 20 mg/ml

Propoven 2%

Emulsion for injection/infusion in pre-filled syringe

For intravenous use.

Each ml contains 20 mg propofol

V004
M0xxxxx/02 UK/IE

Each 50 ml syringe contains 1000 mg propofol.

For single use in one patient.

Pre-filled syringes should be shaken before use. Use immediately after opening. Read the package leaflet for the shelf life of the diluted product. Keep out of the sight and reach of children. Read the package leaflet before use. Do not store above 25 °C. Do not freeze.

Contains Soya-bean oil, refined, medium chain triglycerides, purified egg phosphatides, glycerol, oleic acid, sodium hydroxide, water for injections. See leaflet for further information.

MAH for UK
Fresenius Kabi Ltd
Cestrian Court, Eastgate Way
Manor Park, Runcorn
Cheshire, WA7 1NT, UK

MAH for IE
Fresenius Kabi Deutschland GmbH
Eise-Kroener Strasse 1
Bad Homburg v.d.H
613252, Germany

 **FRESENIUS
KABI**

KabiFill



1 x 50 ml

POM

PL 08828/0239
PA2059/017/005

Batch
EXP



M0xxxx/03 UK/IE
V004

Propofol 20 mg/ml


Propoven 2%
Emulsion for injection/infusion in pre-filled syringe



1 x 50 ml



1 x 50 ml



Propofol 20 mg/ml
Propoven 2%
Emulsion for injection/infusion in pre-filled syringe

For intravenous use.
Each ml contains 20 mg propofol.

Each 50 ml syringe contains 1000 mg propofol.

For single use only.

Pre-filled syringes should be shaken before use.

Use only if the emulsion is homogeneous and the pre-filled syringe undamaged.

Use immediately after opening.

Keep out of the sight and reach of children.

Do not store above 25 °C. Do not freeze.

Contains Soyabean oil, refined, medium chain triglycerides, purified egg phosphatides, glycerol, oleic acid, sodium hydroxide, water for injections.
See leaflet for further information.



1 x 50 ml

MAH for UK
FRESENIUS KABI Ltd
Easton Court, Eastgate Way
Macclesfield, Cheshire, Warrington, UK
6835Z, Germany



PL 08828/0239
PA209/07/005

Propofol 20 mg/ml


Propoven 2%
Emulsion for injection/infusion in pre-filled syringe



1 x 50 ml



1 x 50 ml




Propofol 20 mg/ml

Propoven 2%
Emulsion for injection/infusion in pre-filled syringe

Sterility has to be ensured.
The outer surface of the syringe and the plunger rod are not sterile.

- 1) Take out the syringe from the packaging and shake it.
- 2) Insert the plunger rod by screwing it clock-wise into the syringe.
- 3) Remove the tip cap from the syringe and connect the infusion line, needle or cannula to the syringe. Get rid of the air bubble (a small bubble can remain) and the ready-to-use syringe will be installed in the pump or administered manually. Read the package leaflet before use.



For single use in one patient. Risk of sepsis in multiple use.

Any unused emulsion must be discarded.
Use only under aseptic conditions.



1 x 50 ml



PC: 0xxxxxxxxxxxxx

Batch:

EXP:

SN: