

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

Yaltormin SR 750mg Prolonged Release Tablets

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

One prolonged release tablet contains 750 mg metformin hydrochloride corresponding to 585 mg metformin base.

For the full list of excipients, see section 6.1.

### **3 PHARMACEUTICAL FORM**

Prolonged Release Tablets

White to off-white, capsule shaped tablet debossed with 'SR 750' on one side and plain on other side. The tablets are approximately 19.6 mm in length and 9.3 mm in breadth.

#### **4.1 Therapeutic indications**

- Reduction in the risk or delay of the onset of type 2 diabetes mellitus in adult, overweight patients with IGT\* and/or IFG\*, and/or increased HbA<sub>1C</sub> who are:
  - at high risk for developing overt type 2 diabetes mellitus (see section 5.1) and
  - still progressing towards type 2 diabetes mellitus despite implementation of intensive lifestyle change for 3 to 6 months.

Treatment with Yaltormin SR must be based on a risk score incorporating appropriate measures of glycaemic control and including evidence of high cardiovascular risk (see section 5.1).

Lifestyle modifications should be continued when metformin is initiated, unless the patient is unable to do so because of medical reasons.

\*IGT: Impaired Glucose Tolerance; IFG: Impaired Fasting Glucose

- Treatment of type 2 diabetes mellitus in adults, particularly in overweight patients, when dietary management and exercise alone does not result in adequate glycaemic control. Yaltormin SR may be used as monotherapy or in combination with other oral antidiabetic agents, or with insulin.
- Polycystic ovary syndrome (PCOS)  
Ovulation stimulation to support conception in women with PCOS. Yaltormin SR can be used as monotherapy or in combination with other treatment options.

## 4.2 Posology and method of administration

### Posology

Adults with normal renal function ( $\text{GFR} \geq 90 \text{ mL/min}$ )

#### *Reduction in the risk or delay of the onset of type 2 diabetes*

- Metformin should only be considered where intensive lifestyle modifications for 3 to 6 months have not resulted in adequate glycaemic control.
- The therapy should be initiated with one tablet Yaltormin SR 500 mg once daily with the evening meal.
- After 10 to 15 days dose adjustment on the basis of blood glucose measurements is recommended (OGTT and/or FPG and/or  $\text{HbA}_{1\text{C}}$  values to be within the normal range). A slow increase of dose may improve gastro-intestinal tolerability. The maximum recommended dose is 4 tablets (2000 mg) once daily with the evening meal.
- It is recommended to regularly monitor (every 3-6 months) the glycaemic status (OGTT and/or FPG and/or  $\text{HbA}_{1\text{C}}$  value) as well as the risk factors to evaluate whether treatment needs to be continued, modified or discontinued.
- A decision to re-evaluate therapy is also required if the patient subsequently implements improvements to diet and/or exercise, or if changes to the medical condition will allow increased lifestyle interventions to be possible.

#### *Monotherapy and combination with other oral antidiabetic agents:*

Yaltormin SR 750 mg is intended for patients who are already treated with metformin tablets (prolonged or immediate release).

The dose of Yaltormin SR 750 mg should be equivalent to the daily dose of metformin tablets (prolonged or immediate release), up to a maximum dose of 1500 mg given with the evening meal.

After 10 to 15 days, it is recommended to check that the dose of Yaltormin SR 750 mg is adequate on the basis of blood glucose measurements.

*Combination with insulin:*

For patients already treated with metformin and insulin in combination therapy, the dose of Yaltormin SR 750 mg should be equivalent to the daily dose of metformin tablets up to a maximum of 1500 mg given with the evening meal, while insulin dosage is adjusted on the basis of blood glucose measurements.

*Ovulation stimulation to support conception in women with PCOS:*

The usual dose is 1,500-2000 mg once daily. A slow increase of dose may improve gastrointestinal tolerability. The maximum recommended dose is 2000 mg daily with the evening meal. Yaltormin SR should be considered for at least 6 months.

The decision to maintain therapy after conception and during pregnancy should be made clinically and on a case-by-case basis, taking into consideration the patient needs and an evaluation of the potential risks and benefits.

*Elderly:*

Due to the potential for decreased renal function in elderly subjects, the metformin dosage should be adjusted based on renal function. Regular assessment of renal function is necessary (see section 4.4).

Benefit in the reduction of risk or delay of the onset of type 2 diabetes mellitus has not been established in patients 75 years and older (see section 5.1) and metformin initiation is therefore not recommended in these patients (see section 4.4).

*Renal impairment*

A GFR should be assessed before initiation of treatment with metformin containing products and at least annually thereafter. In patients at an increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3 – 6 months.

| GFR mL/min) | Total maximum daily dose | Additional considerations  |
|-------------|--------------------------|--|
| 60-89       | 2000 mg                  | Dose reduction may be considered in relation to declining renal function   |
| 45-59       | 2000 mg                  | Factors that may increase the risk of lactic acidosis (see section 4.4) should be reviewed before considering initiation of metformin.<br>The starting dose is at most half of the |
| 30-44       | 1000 mg                  |  |

|    |   |                              |
|----|---|------------------------------|
|    |   | maximum dose.                |
| 30 | - | Metformin is contraindicated |

*Paediatric population:*

In the absence of available data, Yaltormin SR should not be used in children.

Method of administration

The tablets should be swallowed whole with a drink of water. They should not be chewed or crushed.

### 4.3 Contraindications

- Hypersensitivity to metformin or to any of the excipients listed in section 6.1
- Any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis)
- Diabetic pre-coma
- Severe renal failure (GFR <30mL/min)
- Acute conditions with the potential to alter renal function such as:
  - dehydration,
  - severe infection,
  - shock
- Disease which may cause tissue hypoxia (especially acute disease, or worsening of chronic disease) such as:
  - decompensated heart failure,
  - respiratory failure,
  - recent myocardial infarction,
  - shock
- Hepatic insufficiency, acute alcohol intoxication, alcoholism.

### 4.4 Special warnings and precautions for use

**Lactic acidosis**

Lactic acidosis, a very rare, but serious metabolic complication, most often occurs at acute worsening of renal function or cardiorespiratory illness or sepsis. Metformin accumulation occurs at acute worsening of renal function and increases the risk of lactic acidosis.

In case of dehydration (severe diarrhoea or vomiting, fever or reduced fluid intake), metformin should be temporarily discontinued and contact with a health care professional is recommended.

Medicinal products that can acutely impair renal function (such as antihypertensives, diuretics and NSAIDs) should be initiated with caution in metformin-treated patients. Other risk factors for lactic acidosis are excessive alcohol intake, hepatic insufficiency, inadequately controlled diabetes, ketosis, prolonged fasting and any conditions associated with hypoxia, as well as

concomitant use of medicinal products that may cause lactic acidosis (see sections 4.3 and 4.5).

Patients and/or care-givers should be informed of the risk of lactic acidosis. Lactic acidosis is characterised by acidotic dyspnoea, abdominal pain, muscle cramps, asthenia and hypothermia followed by coma. In case of suspected symptoms, the patient should stop taking metformin and seek immediate medical attention. Diagnostic laboratory findings are decreased blood pH (< 7.35), increased plasma lactate levels (>5 mmol/L) and an increased anion gap and lactate/pyruvate ratio.

#### Patients with known or suspected mitochondrial diseases:

In patients with known mitochondrial diseases such as Mitochondrial Encephalopathy with Lactic Acidosis, and Stroke-like episodes (MELAS) syndrome and Maternal inherited diabetes and deafness (MIDD), metformin is not recommended due to the risk of lactic acidosis exacerbation and neurologic complications which may lead to worsening of the disease.

In case of signs and symptoms suggestive of MELAS syndrome or MIDD after the intake of metformin, treatment with metformin should be withdrawn immediately and prompt diagnostic evaluation should be performed.

#### **Renal function:**

GFR should be assessed before treatment initiation and regularly thereafter, see section 4.2. Metformin is contraindicated in patients with GFR<30 mL/min and should be temporarily discontinued in the presence of conditions that alter renal function, see section 4.3.

#### **Cardiac function:**

Patients with heart failure are more at risk of hypoxia and renal insufficiency. In patients with stable chronic heart failure, metformin may be used with a regular monitoring of cardiac and renal function.

For patients with acute and unstable heart failure, metformin is contraindicated (see section 4.3).

#### **Elderly:**

Due to the limited therapeutic efficacy data in the reduction of risk or delay of type 2 diabetes in patients 75 years and older, metformin initiation is not recommended in these patients.

#### **Administration of iodinated contrast agents:**

Intravascular administration of iodinated contrast agents may lead to contrast induced nephropathy, resulting in metformin accumulation and an increased risk of lactic acidosis. Metformin should be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable, see sections 4.2 and 4.5.

**Surgery:**

Metformin must be discontinued at the time of surgery with general, spinal or epidural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and provided that renal function has been re-evaluated and found to be stable.

**Other precautions:**

All patients should continue their diet with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.

The usual laboratory tests for diabetes monitoring should be performed regularly.

Metformin may reduce vitamin B12 serum levels. The risk of low vitamin B12 levels increases with increasing metformin dose, treatment duration, and/or in patients with risk factors known to cause vitamin B12 deficiency. In case of suspicion of vitamin B12 deficiency (such as anaemia or neuropathy), vitamin B12 serum levels should be monitored. Periodic vitamin B12 monitoring could be necessary in patients with risk factors for vitamin B12 deficiency. Metformin therapy should be continued for as long as it is tolerated and not contra-indicated and appropriate corrective treatment for vitamin B12 deficiency provided in line with current clinical guidelines.

Metformin alone never causes hypoglycaemia, although caution is advised when it is used in combination with insulin or other oral antidiabetics (e.g. sulphonylureas or meglitinides).

The tablet shells may be present in the faeces. Patients should be advised that this is normal.

**Yaltormin SR contain sodium**

This medicine contains less than 1mmol sodium (23mg) per dosage unit, that is to say it is essentially 'sodium free'.

**4.5 Interaction with other medicinal products and other forms of interaction**Concomitant use not recommended*Alcohol*

Alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in case of fasting, malnutrition or hepatic impairment.

*Iodinated contrast agents*

Metformin must be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable, see sections 4.2 and 4.4.

### Combinations requiring precautions for use

Some medicinal products can adversely affect renal function which may increase the risk of lactic acidosis, e.g. NSAIDs, including selective cyclooxygenase (COX) II inhibitors, ACE inhibitors, angiotensin II receptor antagonists and diuretics, especially loop diuretics. When starting or using such products in combination with metformin, close monitoring of renal function is necessary.

*Medicinal products with intrinsic hyperglycaemic activity (e.g. glucocorticoids (systemic and local routes) and sympathomimetics).*

More frequent blood glucose monitoring may be required, especially at the beginning of treatment. If necessary, adjust the metformin dosage during therapy with the other drug and upon its discontinuation.

*Organic cation transporters (OCT)*

Metformin is a substrate of both transporters OCT1 and OCT 2.

Co-administration of metformin with

- Inhibitors of OCT 1 (such as verapamil) may reduce efficacy of metformin.
- Inducers of OCT 1 (such as rifampicin) may increase gastrointestinal absorption and efficacy of metformin.
- Inhibitors of OCT 2 (such as cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazole) may decrease the renal elimination of metformin and thus lead to an increase in metformin plasma concentration.
- Inhibitors of both OCT 1 and OCT 2 (such as crizotinib, olaparib) may alter efficacy and renal elimination of metformin.

Caution is therefore advised, especially in patients with renal impairment, when these drugs are co-administered with metformin, as metformin plasma concentration may increase. If needed, dose adjustment of metformin may be considered as OCT inhibitors/inducers may alter the efficacy of metformin.

## **4.6 Fertility, pregnancy and lactation**

*Pregnancy*

Uncontrolled hyperglycaemia in the periconceptional phase and during pregnancy is associated with increased risk of congenital abnormalities, pregnancy loss, pregnancy-induced hypertension, preeclampsia and perinatal mortality. It is important to maintain blood glucose levels as close to normal as possible throughout pregnancy, to reduce the risk of adverse hyperglycaemia-related outcomes to the mother and her child.

Metformin crosses the placenta with levels that can be as high as maternal concentrations.

A large amount of data on pregnant women (more than 1000 exposed outcomes) from a register-based cohort study and published data (meta-analyses, clinical studies, and registries) indicates no increased risk of congenital

abnormalities nor feto/neonatal toxicity after exposure to metformin in the periconceptual phase and/or during pregnancy.

There is limited and inconclusive evidence on the metformin effect on the long-term weight outcome of children exposed in utero. Metformin does not appear to affect motor and social development up to 4 years of age in children exposed during pregnancy although data on long term outcomes are limited.

If clinically needed, the use of metformin can be considered during pregnancy and in the periconceptual phase as an addition or an alternative to insulin.

*Women of childbearing potential:*

In premenopausal women, treatment with Yaltormin SR may result in ovulation in anovulatory women, which may lead to unintended pregnancy.

*Breast-feeding*

Metformin is excreted into human breast milk. No adverse effects were observed in breastfed newborns/infants. However, as only limited data are available, breastfeeding is not recommended during metformin treatment. A decision on whether to discontinue breast-feeding should be made, taking into account the benefit of breast-feeding and the potential risk to adverse effect on the child.

*Fertility*

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

#### **4.7 Effects on ability to drive and use machines**

Metformin monotherapy does not cause hypoglycaemia and therefore has no effect on the ability to drive or to use machines.

However, patients should be alerted to the risk of hypoglycaemia when metformin is used in combination with other antidiabetic agents (e.g. sulphonylureas, insulin, or meglinitides).

#### **4.8 Undesirable effects**

In post marketing data and in controlled clinical studies, adverse event reporting in patients treated with metformin SR was similar in nature and severity to that reported in patients treated with metformin immediate release.

During treatment initiation, the most common adverse reactions are nausea, vomiting, diarrhoea, abdominal pain and loss of appetite, which resolve spontaneously in most cases.

The following adverse reactions may occur with Yaltormin SR.

Frequencies are defined as follows: very common:  $>1/10$ ; common  $\geq 1/100$ ,  $<1/10$ ; uncommon  $\geq 1/1,000$ ,  $<1/100$ ; rare  $\geq 1/10,000$ ,  $<1/1,000$ ; very rare  $<1/10,000$ .

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

#### Metabolism and nutrition disorders

*Common:*

- Vitamin B12 decrease/deficiency (see section 4.4)

*Very rare:*

- Lactic acidosis (see 4.4. Special warnings and precautions for use).

#### Nervous system disorders

*Common:*

- Taste disturbance

#### Gastrointestinal disorders

*Very common:*

- Gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite. These undesirable effects occur most frequently during initiation of therapy and resolve spontaneously in most cases. A slow increase of the dose may also improve gastrointestinal tolerability.

#### Hepatobiliary disorders

*Very rare:*

- *Isolated* reports of liver function tests abnormalities or hepatitis resolving upon metformin discontinuation.

#### Skin and subcutaneous tissue disorders

*Very rare:*

- Skin reactions such as erythema, pruritus, urticaria

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

Hypoglycaemia has not been seen with metformin doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose or concomitant risks of metformin may lead to lactic acidosis. Lactic acidosis is a medical emergency and

must be treated in hospital. The most effective method to remove lactate and metformin is haemodialysis.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

#### ORAL ANTI-DIABETICS

(A10BA02: Gastrointestinal tract and metabolism)

Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and postprandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

#### Mechanism of action

Metformin is a biguanide with antihyperglycaemic effects, on both basal and postprandial hyperglycaemia. It does not stimulate insulin secretion and therefore does not cause hypoglycaemia.

Metformin reduces basal hyperinsulinemia, and in combination with insulin, reduces insulin requirement.

Metformin exerts its antihyperglycaemic effect via multiple mechanisms: Metformin reduces hepatic glucose production.

Metformin facilitates peripheral glucose uptake and utilisation, in part by increasing insulin action.

Metformin alters glucose turnover in the gut: Uptake from circulation is increased and absorption from food is decreased. Additional mechanisms attributed to the gut include an increase in release of glucagon-like peptide 1 (GLP-1) and a decrease of bile acid resorption. Metformin alters the gut microbiome.

Metformin can improve the lipid profile in hyperlipidaemic individuals. In clinical studies, use of metformin was associated with either a stable body weight or modest weight loss.

Metformin lowers PCOS-related hyperandrogenism.

Metformin is an adenosine monophosphate-protein-kinase (AMPK) activator and increases the transport capacity of all types of membrane glucose transporters (GLUTs).

#### Clinical efficacy

***Reduction in the risk or delay of type 2 diabetes mellitus***

The **Diabetes Prevention Program (DPP)** was a multicenter randomised controlled clinical trial in adults assessing the efficacy of an intensive lifestyle intervention or metformin to prevent or delay the development of type 2 diabetes mellitus. Inclusion criteria were age  $\geq 25$  years, BMI  $\geq 24$  kg/m<sup>2</sup> ( $\geq 22$  kg/m<sup>2</sup> for Asian-Americans), and impaired glucose tolerance plus a fasting plasma glucose of 95 – 125 mg/dl (or  $\leq 125$  mg/dl for American Indians). Patients were either treated with intensive lifestyle intervention, 2x850 mg metformin plus standard lifestyle change, or placebo plus standard lifestyle change.

The mean baseline values of the DPP participants (n=3,234 for 2.8 years) were age 50.6 $\pm$ 10.7 years, 106.5 $\pm$ 8.3 mg/dl fasted plasma glucose, 164.6 $\pm$ 17.0 mg/dl plasma glucose two hours after an oral glucose load, and 34.0 $\pm$ 6.7 kg/m<sup>2</sup> BMI. Intensive lifestyle intervention as well as metformin significantly reduced the risk of developing overt diabetes compared to placebo, 58% (95% CI 48-66%) and 31% (95% CI 17-43%), respectively.

The advantage of the lifestyle intervention over metformin was greater in older persons.

The patients who benefited most from the metformin treatment were aged below 45 years, with a BMI equal or above 35kg/m<sup>2</sup>, a baseline glucose 2 h value of 9.6-11.0 mmol/l, a baseline HbA<sub>1C</sub> equal or above 6.0% or with a history of gestational diabetes.

To prevent one case of overt diabetes during the three years in the whole population of the DPP, 6.9 patients had to participate in the intensive lifestyle group and 13.9 in the metformin group. The point of reaching a cumulative incidence of diabetes equal to 50% was delayed by about three years in the metformin group compared to placebo.

The **Diabetes Prevention Program Outcomes Study (DPPOS)** is the long-term follow-up study of the DPP including more than 87% of the original DPP population for long-term follow up.

Among the DPPOS participants (n=2776), the cumulative incidence of diabetes at year 15 is 62% in the placebo group, 56% in the metformin group, and 55% in the intensive lifestyle intervention group. Crude rates of diabetes are 7.0, 5.7 and 5.2 cases per 100 person-years among the placebo, metformin, and intensive lifestyle participants, respectively. Reductions in the diabetes risk were of 18% (hazard ratio (HR) 0.82, 95% CI 0.72–0.93; p=0.001) for the metformin group and 27% (HR 0.73, 95% CI 0.65–0.83; p<0.0001) for the intensive lifestyle intervention group, when compared with the placebo group. For an aggregate microvascular endpoint of nephropathy, retinopathy and neuropathy, the outcome was not significantly different between the treatment groups, but among the participants who had not developed diabetes during DPP/DPPOS, the prevalence of the aggregate microvascular outcome was 28% lower compared with those who had developed diabetes (Risk Ratio 0.72, 95% CI 0.63–0.83; p<0.0001). No prospective comparative data for metformin

on macrovascular outcomes in patients with IGT and/or IFG and/or increased HbA<sub>1C</sub> are available.

Published risk factors for type 2 diabetes include: Asian or black ethnic background, age above 40, dyslipidaemia, hypertension, obesity or being overweight, age, 1st degree family history of diabetes, history of gestational diabetes mellitus, and polycystic ovary syndrome (PCOS).

Consideration must be given to current national guidance on the definition of prediabetes.

Patients at high risk should be identified by a validated risk-assessment tool.

### ***Treatment of type 2 diabetes mellitus***

The prospective randomised (UKPDS) study has established the long-term benefit of intensive blood glucose control in overweight type 2 diabetic patients treated with immediate release metformin as first-line therapy after diet failure. Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:

- a significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/1000 patient-years) versus diet alone (43.3 events/ 1000 patient-years),  $p=0.0023$ , and versus the combined sulphonylurea and insulin monotherapy groups (40.1 events/ 1000 patient-years),  $p=0.0034$ .
- a significant reduction of the absolute risk of diabetes-related mortality: metformin 7.5 events/1000 patient-years, diet alone 12.7 events/ 1000 patient-years,  $p=0.017$ ;
- a significant reduction of the absolute risk of overall mortality: metformin 13.5 events/ 1000 patient-years versus diet alone 20.6 events/ 1000 patient-years ( $p=0.011$ ), and versus the combined sulphonylurea and insulin monotherapy groups 18.9 events/ 1000 patient-years ( $p=0.021$ );
- a significant reduction in the absolute risk of myocardial infarction: metformin 11 events/ 1000 patient-years, diet alone 18 events/ 1000 patient-years ( $p=0.01$ )

For metformin used as second-line therapy, in combination with a sulphonylurea, benefit regarding clinical outcome has not been shown.

In type 1 diabetes, the combination of metformin and insulin has been used in selected patients, but the clinical benefit of this combination has not been formally established.

## **5.2 Pharmacokinetic properties**

### Absorption

Following a single oral administration of 1500 mg of metformin SR 750 mg, a mean peak plasma concentration of 1193 ng/ml is achieved with a median value of 5 hours and a range of 4 to 12 hours.

Metformin SR 750 mg was shown to be bioequivalent to metformin SR 500 mg at a 1500 mg dose with respect to C<sub>max</sub> and AUC in healthy fed and fasted subjects.

*The bioequivalent product shows the following properties:*

At steady state, similar to the immediate release formulation, C<sub>max</sub> and AUC are not proportionally increased to the administered dose. The AUC after a single oral administration of 2000mg of metformin prolonged release tablets is similar to that observed after administration of 1000mg of metformin immediate release tablets b.i.d.

Intrasubject variability of C<sub>max</sub> and AUC of metformin prolonged release tablets is comparable to that observed with metformin immediate release tablets.

When the prolonged release tablet is administered in fasting conditions the AUC is decreased by 30% (both C<sub>max</sub> and T<sub>max</sub> are unaffected).

Metformin absorption from the prolonged release formulation is not altered by meal composition.

No accumulation is observed after repeated administration of up to 2000mg of metformin prolonged release tablets.

#### Distribution

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean volume of distribution (V<sub>d</sub>) ranged between 63-276 L.

#### Metabolism

Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

#### Elimination

Renal clearance of metformin is > 400 ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

Characteristics in specific groups of patients

Renal impairment

The available data in subjects with moderate renal insufficiency are scarce and no reliable estimation of the systemic exposure to metformin in this subgroup as compared to subjects with normal renal function could be made. Therefore, the dose adaptation should be made upon clinical efficacy/tolerability considerations (see section 4.2).

### **5.3 Preclinical safety data**

Preclinical data reveal no special hazard for humans based on conventional studies on safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity reproduction.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Magnesium stearate  
Silica colloidal anhydrous  
Carmellose sodium  
Hypromellose

### **6.2 Incompatibilities**

None

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

28 and 56 tablets in blister strips composed of aluminium foil and PVC.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements. Any unused product or waste material should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Wockhardt UK Ltd  
Ash Road North  
Wrexham  
LL13 9UF  
United Kingdom

## **8 MARKETING AUTHORISATION NUMBER(S)**

PL 29831/0656

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

21/12/2021

**10 DATE OF REVISION OF THE TEXT**

09/07/2025