

120.00 mm

80.00 mm

Care +
Heartburn Relief
Oral Suspension

Sodium Bicarbonate,
Sodium Alginate,
Calcium Carbonate

Provides fast effective relief from
Acid Indigestion & Heartburn

Peppermint Flavour

SUGAR FREE **500 ml**

Please read this entire label before use.
Dosage: For oral use. Shake well before use.
Adults and children over 12 years:
Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.
Children under 12 years:
Should be given only on medical advice.
See leaflet for further information.
This product is sugar free and gluten free. You can take this product if you are pregnant or breast-feeding.
Keep out of the sight and reach of children. Do not store above 25°C. Do not refrigerate.
Each 5 ml of this medicine contains:
133.5 mg Sodium Bicarbonate, 250 mg Sodium Alginate, 80 mg Calcium Carbonate. It also contains carbomer, sodium hydroxide, saccharin sodium, ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), butyl parahydroxybenzoate, isopropyl alcohol, peppermint oil, purified water. Contains sodium. See leaflet for further information. 93089692312 GB 23LL0594
Distributed by
Care, Linthwaite, Huddersfield, HD7 5QH
MA Holder and Manufacturer:
Pinewood Laboratories Ltd., Ballymacarbry, Clonmel, Co. Tipperary, Ireland. PL 04917/0027 UK

Peel Here

5 011309 113211

PACKAGE LEAFLET: INFORMATION FOR THE USER
Care Heartburn Relief Oral Suspension
Peppermint Flavour

sodium bicarbonate, sodium alginate, calcium carbonate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Always take this medicine exactly as described in the leaflet or as your doctor or pharmacist has told you.

What this medicine is and what it is used for

This medicine belongs to a group of medicines called 'reflux suppressants' which form a protective layer on top of the stomach contents to prevent stomach acid escaping from the stomach where it works into the food pipe causing pain and discomfort. This medicine is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy.

Before using this medicine

Do not take this product if:

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

Take special care before treatment with this medicine:

This medicine contains 286.5 mg (12.45 mmol) sodium (main component of cooking/table salt) in each 20 ml dose. This is equivalent to 14.3 % of the recommended maximum daily dietary intake of sodium for an adult. The maximum daily dose of this product is equivalent to 57.2 % of the WHO recommended maximum daily intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period, especially if you have been advised to follow a low salt (sodium) diet.

This medicine contains ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216) and butyl parahydroxybenzoate which may cause an allergic reaction (possibly delayed).

Taking other medicines


Do **not** take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparations, antifungals such as ketoconazole, digoxin and beta blockers (for heart conditions), penicillamine (for rheumatoid arthritis), glucocorticoids (for inflammatory and autoimmune disorders), neuroleptics (for mental

stick on area (no print here)

23LL0594

Braille Translation:

Care Heartburn Relief Oral Suspension



stick on area (no print here)

illness), thyroid hormones, chloroquine (for malaria), estramustine (for prostate cancer), bisphosphonates (for osteoporosis), ACE inhibitors (for high blood pressure), analgesics (for pain relief), antiepileptics, antipsychotics and lithium (for mental health problems).

How to take this medicine

To be taken by mouth, after meals and at bedtime. Shake bottle well before use.

Adults and children over 12 years:

Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 12 years:

Should be given only on medical advice.

After taking this medicine:

If symptoms persist after 7 days consult your doctor.

If you take too much of this product you may feel bloated and experience some abdominal discomfort. It is unlikely to cause any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

Possible side effects

Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, difficulty in

breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any side effects stop taking the product and consult your doctor or pharmacist.

Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

How to store this medicine

Keep all medicines out of the sight and reach of children.

Do not use after the expiry date shown on the pack (EXP month/year).

Do not store above 25°C.

Do not refrigerate.

PL 04917/0027

Revision date: 12/2023

93089692312 GB
23LL05944

120.00 mm

80.00 mm

Braille Translation:

Care Heartburn Relief Oral Suspension

C A R E H E A R T B U R N
 R E L I E F O R A L
 S U S P E N S I O N

Care +

Heartburn Relief

Oral Suspension

Sodium Bicarbonate,
Sodium Alginate,
Calcium Carbonate

Peppermint Flavour

SUGAR FREE

500 ml

Provides fast effective relief from Acid Indigestion & Heartburn

Please read this entire label before use.
Dosage: For oral use. Shake well before use.
Adults and children over 12 years:
 Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.
Children under 12 years:
 Should be given only on medical advice.
See leaflet for further information.
 This product is sugar free and gluten free. You can take this product if you are pregnant or breast-feeding.
Keep out of the sight and reach of children. Do not store above 25°C. Do not refrigerate.

Each 5 ml of this medicine contains:
 133.5 mg Sodium Bicarbonate, 250 mg Sodium Alginate, 80 mg Calcium Carbonate. It also contains carbomer, sodium hydroxide, saccharin sodium, ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216), butyl parahydroxybenzoate, isopropyl alcohol, peppermint oil, purified water. Contains sodium. See leaflet for further information.

Distributed by
 Care, Linthwaite, Huddersfield, HD7 5QH

MA Holder and Manufacturer:
 Pinewood Laboratories Ltd., Ballymacarbry, Clonmel, Co. Tipperary, Ireland. PL 04917/0027

Peel Here

5 011309 113211 >



PACKAGE LEAFLET: INFORMATION FOR THE USER
Care Heartburn Relief Oral Suspension
Peppermint Flavour

sodium bicarbonate, sodium alginate, calcium carbonate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Always take this medicine exactly as described in the leaflet or as your doctor or pharmacist has told you.

What this medicine is and what it is used for

This medicine belongs to a group of medicines called 'reflux suppressants' which form a protective layer on top of the stomach contents to prevent stomach acid escaping from the stomach where it works into the food pipe causing pain and discomfort. This medicine is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, which may occur, for example, following meals or during pregnancy.

Before using this medicine

Do not take this product if:

- You know you are allergic to any of the ingredients as very rarely difficulty in breathing and skin rashes have occurred (see further information for a full list of ingredients).

Take special care before treatment with this medicine:

This medicine contains 286.5 mg (12.45 mmol) sodium (main component of cooking/table salt) in each 20 ml dose. This is equivalent to 14.3 % of the recommended maximum daily dietary intake of sodium for an adult. The maximum daily dose of this product is equivalent to 57.2 % of the WHO recommended maximum daily intake for sodium. Talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period, especially if you have been advised to follow a low salt (sodium) diet.

This medicine contains ethyl parahydroxybenzoate (E214), propyl parahydroxybenzoate (E216) and butyl parahydroxybenzoate which may cause an allergic reaction (possibly delayed).

Taking other medicines

Do **not** take this product within two hours of taking other medicines by mouth as it can interfere with the action of some other medicines. This is especially important if you are taking antibiotics (tetracyclines and quinolones such as norfloxacin), iron preparations, antifungals such as ketoconazole, digoxin and beta blockers (for heart conditions), penicillamine (for rheumatoid arthritis), glucocorticoids (for inflammatory and autoimmune disorders), neuroleptics (for mental

23LL0594PW

stick on area (no print here)

stick on area (no print here)

illness), thyroid hormones, chloroquine (for malaria), estramustine (for prostate cancer), bisphosphonates (for osteoporosis), ACE inhibitors (for high blood pressure), analgesics (for pain relief), antiepileptics, antipsychotics and lithium (for mental health problems).

How to take this medicine

To be taken by mouth, after meals and at bedtime. Shake bottle well before use.

Adults and children over 12 years:

Take 10-20 ml (two to four 5 ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 12 years:

Should be given only on medical advice.

After taking this medicine:

If symptoms persist after 7 days consult your doctor.

If you take too much of this product you may feel bloated and experience some abdominal discomfort. It is unlikely to cause any harm, but please consult your doctor or pharmacist. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.

Possible side effects

Very rarely (less than 1 in 10,000 patients treated) an allergic reaction to the ingredients may occur. Symptoms of this may include skin rash, difficulty in

breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any side effects stop taking the product and consult your doctor or pharmacist.

Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

How to store this medicine

Keep all medicines out of the sight and reach of children.

Do not use after the expiry date shown on the pack (EXP month/year).

Do not store above 25°C.

Do not refrigerate.

PL 04917/0027

Revision date: 12/2023

93089692312 GB
23LL05944PW