

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

Tamsulosin Ascend 400 mcg modified-release capsule, hard

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each hard gelatin capsule contains as active ingredient tamsulosin hydrochloride 0.4 mg.

Excipient(s):

For a full list of excipients, see section 6.1.

### **3 PHARMACEUTICAL FORM**

Modified-release hard capsule

White to off white pellets filled in hard gelatin capsule size “1” Olive green opaque cap & orange opaque body imprinted with “TAM” on cap & “0.4 mg” on body in black ink.

## **4 CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Lower urinary tract symptoms (LUTS) associated with benign prostatic hyperplasia (BPH).

### **4.2 Posology and method of administration**

Oral use

One capsule daily, to be taken after breakfast or the first meal of the day.

The capsule must be swallowed whole and must not be crunched or chewed, as this interferes with the modified release of the active ingredient.

No dose adjustment is warranted in renal impairment. No dose adjustment is warranted in patients with mild to moderate hepatic insufficiency (see also 4.3 Contraindications).

#### *Paediatric population*

There is no relevant indication for use of tamsulosin in children.

The safety and efficacy of tamsulosin in children <18 years have not been established. Currently available data are described in section 5.1

### **4.3 Contraindications**

Hypersensitivity to tamsulosin hydrochloride, including drug-induced angioedema or to any of the excipients.

A history of orthostatic hypotension.

Severe hepatic insufficiency

#### **4.4 Special warnings and precautions for use**

As with other  $\alpha_1$ -adrenoceptors antagonists, a reduction in blood pressure can occur in individual cases during treatment with tamsulosin 0,4, as a result of which, rarely, syncope can occur. At the first signs of orthostatic hypotension (dizziness, weakness), the patient should sit or lie down until the symptoms have disappeared.

Before therapy with tamsulosin 0,4 is initiated, the patient should be examined in order to exclude the presence of other conditions, which can cause the same symptoms as benign prostatic hyperplasia. Digital rectal examination and, when necessary, determination of prostate specific antigen (PSA) should be performed before treatment and at regular intervals afterwards.

The treatment of patients with severe renal impairment (creatinine clearance of < 10 ml/min) should be approached with caution, as these patients have not been studied.

The 'Intraoperative Floppy Iris Syndrome' (IFIS, a variant of small pupil syndrome) has been observed during cataract and glaucoma surgery in some patients on or previously treated with tamsulosin hydrochloride. IFIS may increase the risk of eye complications during and after the operation.

Discontinuing tamsulosin hydrochloride 1-2 weeks prior to cataract or glaucoma surgery is anecdotally considered helpful, but the benefit of treatment discontinuation has not yet been established. IFIS has also been reported in patients who had discontinued tamsulosin for a longer period prior to surgery.

The initiation of therapy with tamsulosin hydrochloride in patients for whom cataract or glaucoma surgery is scheduled is not recommended. During pre-operative assessment, surgeons and ophthalmic teams should consider whether patients scheduled for cataract or glaucoma surgery are being or have been treated with tamsulosin in order to ensure that appropriate measures will be in place to manage the IFIS during surgery.

Tamsulosin hydrochloride should not be given in combination with strong inhibitors of CYP3A4 in patients with poor metaboliser CYP2D6 phenotype.

Tamsulosin hydrochloride should be used with caution in combination with strong and moderate inhibitors of CYP3A4 (see section 4.5).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have only been performed in adults.

No interactions have been seen when tamsulosin hydrochloride was given concomitantly with either atenolol, enalapril or theophylline. Concomitant cimetidine brings about a rise in plasma levels of tamsulosin, whereas furosemide a fall, but as levels remain within the normal range posology need not be adjusted.

In vitro, neither diazepam nor propranolol, trichlormethiazide, chlormadinon, amitriptyline, diclofenac, glibenclamide, simvastatin and warfarin change the free fraction of tamsulosin in human plasma. Neither does tamsulosin change the free fractions of diazepam, propranolol, trichlormethiazide and chlormadinon.

Diclofenac and warfarin, however, may increase the elimination rate of tamsulosin.

Concomitant administration of tamsulosin hydrochloride with strong inhibitors of CYP3A4 may lead to increased exposure to tamsulosin hydrochloride. Concomitant administration with ketoconazole (a known strong CYP3A4 inhibitor) resulted in an increase in AUC and  $C_{max}$  of tamsulosin hydrochloride by a factor of 2.8 and 2.2, respectively.

Tamsulosin hydrochloride should not be given in combination with strong inhibitors of CYP3A4 in patients with poor metaboliser CYP2D6 phenotype.

Tamsulosin hydrochloride should be used with caution in combination with strong and moderate inhibitors of CYP3A4.

Concomitant administration of tamsulosin hydrochloride with paroxetine, a strong inhibitor of CYP2D6, resulted in a  $C_{max}$  and AUC of tamsulosin that had increased by a factor of 1.3 and 1.6, respectively, but these increases are not considered clinically relevant.

Concurrent administration of other  $\alpha_1$ -adrenoceptor antagonists could lead to hypotensive effects

#### 4.6 Fertility, Pregnancy and lactation

Tamsulosin is not indicated for use in women.

Ejaculation disorders have been observed in short and long term clinical studies with tamsulosin. Events of ejaculation disorder, retrograde ejaculation and ejaculation failure have been reported in the post authorization phase.

#### 4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

However, patients should be aware of the fact that dizziness can occur.

#### 4.8 Undesirable effects

<b>System Organ Class</b>	<b>Common (&gt;1/100, &lt;1/10)</b>	<b>Uncommon (&gt;1,000, &lt;1/100)</b>	<b>Rare (&gt;1/10,000, &lt;1/1,000)</b>	<b>Very rare (&lt;1/10,000)</b>	<b>Not known (cannot be estimated from the available data)</b>
<b>Nervous systems disorders</b>	Dizziness (1.3%)	Headache	Syncope		
<b>Eye disorders</b>					Blurred vision* Visual impairment*
<b>Cardiac disorders</b>		Palpitations			
<b>Vascular</b>		Orthostatic			

<b>disorders</b>		hypotension			
<b>Respiratory, thoracic and mediastinal disorders</b>		Rhinitis			Epistaxis*
<b>Gastrointestinal disorders</b>		Constipation, diarrhoea, nausea, vomiting			Dry mouth*
<b>Skin and subcutaneous tissue disorders</b>		Rash, pruritus, urticaria	Angioedema	Stevens-Johnson syndrome	Erythema multiforme*, exfoliative dermatitis

\*Observed post-marketing

During cataract and glaucoma surgery a small pupil situation, known as Intraoperative Floppy Iris Syndrome (IFIS), has been associated with therapy of tamsulosin during post-marketing surveillance (see also section 4.4).

Post-marketing experience: In addition to the adverse events listed above, atrial fibrillation, arrhythmia, tachycardia and dyspnoea have been reported in association with tamsulosin use. Because these spontaneously reported events are from the worldwide post marketing experience, the frequency of events and the role of tamsulosin in their causation cannot be reliably determined.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: <http://www.mhra.gov.uk/yellowcard> or search for MHRA Yellow Card in the Google Play or Apple App Store.

## 4.9 Overdose

### Symptoms

Overdosage with tamsulosin hydrochloride can potentially result in severe hypotensive effects. Severe hypotensive effects have been observed at different levels of overdosing.

### Treatment

In case of acute hypotension occurring after overdosage cardiovascular support should be given. Blood pressure can be restored and heart rate brought back to normal by lying the patient down. If this does not help then volume expanders and, when necessary, vasopressors could be employed. Renal function should be monitored and general supportive measures applied. Dialysis is unlikely to be of help as tamsulosin is very highly bound to plasma proteins.

Measures, such as emesis, can be taken to impede absorption. When large quantities are involved, gastric lavage can be applied and activated charcoal and an osmotic laxative such as sodium sulphate, can be administered.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group:  $\alpha_1$ -adrenoceptor antagonist, ATC code: G04CA02.  
Preparations for the exclusive treatment of prostatic disease.

Mechanism of action:

Tamsulosin binds selectively and competitively to the postsynaptic  $\alpha_1$ -adrenoceptors, in particular to subtypes  $\alpha_{1A}$  and  $\alpha_{1D}$ . It brings about relaxation of prostatic and urethral smooth muscle.

Pharmacodynamic effects:

Tamsulosin 0,4 increases the maximum urinary flow rate. It relieves obstruction by relaxing smooth muscle in prostate and urethra thereby improving voiding symptoms.

It also improves the storage symptoms in which bladder instability plays an important role.

These effects on storage and voiding symptoms are maintained during long – term therapy. The need for surgery or catheterization is significantly delayed.

$\alpha_1$ -adrenoceptors antagonists can reduce blood pressure by lowering peripheral resistance. No reduction in blood pressure of any clinical significance was observed during studies with tamsulosin 0,4.

### ***Paediatric population***

A double-blind, randomized, placebo-controlled, dose ranging study was performed in children with neuropathic bladder. A total of 161 children (with an age of 2 to 16 years) were randomized and treated at 1 of 3 dose levels of tamsulosin (low [0.001 to 0.002 mg/kg], medium [0.002 to 0.004 mg/kg], and high [0.004 to 0.008 mg/kg]), or placebo.

The primary endpoint was number of patients who decreased their detrusor leak point pressure (LPP) to  $<40$  cm H<sub>2</sub>O based upon two evaluations on the same day. Secondary endpoints were: Actual and percent change from baseline in detrusor leak point pressure, improvement or stabilization of hydronephrosis and hydroureter and change in urine volumes obtained by catheterisation and number of times wet at time of catheterization as recorded in catheterisation diaries. No statistically significant difference was found between the placebo group and any of the 3 tamsulosin dose groups for either the primary or any secondary endpoints. No dose response was observed for any dose level.

## **5.2 Pharmacokinetic properties**

Absorption:

Tamsulosin hydrochloride is absorbed from the intestine and is almost completely bioavailable. Absorption of tamsulosin hydrochloride is reduced by a recent meal. Uniformity of absorption can be promoted by the patient always taking tamsulosin 0.4 mg after the usual breakfast. Tamsulosin shows linear kinetics.

After a single dose of tamsulosin 0.4 mg in the fed state, plasma levels of tamsulosin peak at around 6 hours and, in the steady state, which is reached by day 5 of multiple dosing,  $C_{max}$  in patients is about two thirds higher than that reached after a single dose. Although this was seen in elderly patients, the same finding would also be expected in young ones.

There is a considerable inter-patient variation in plasma levels both after single and multiple dosing.

#### Distribution:

In man, tamsulosin is about 99% bound to plasma proteins. The volume of distribution is small (about 0.2 l/kg).

#### Biotransformation:

Tamsulosin has a low first pass effect, being metabolized slowly. Most tamsulosin is present in plasma in the form of unchanged active substance. It is metabolized in the liver.

In rats, hardly any induction of microsomal liver enzymes was seen to be caused by tamsulosin.

In vitro results suggest that CYP3A4 and also CYP2D6 are involved in metabolism, with possible minor contributions to tamsulosin hydrochloride metabolism by other CYP isozymes. Inhibition of CYP3A4 and CYP2D6 drug metabolizing enzymes may lead to increased exposure to tamsulosin hydrochloride (see section 4.4 and 4.5).

None of the metabolites are more active than the original compound.

#### Elimination:

Tamsulosin and its metabolites are mainly excreted in the urine with about 9% of a dose being present in the form of unchanged active substance.

After a single dose of tamsulosin 0.4 mg after breakfast, and in the steady state in patients, elimination half-lives of about 10 and 13 hours respectively have been measured.

### **5.3 Preclinical safety data**

Single and repeat dose toxicity studies were performed in mice, rats and dogs. In addition, reproduction toxicity in rats, carcinogenicity in mice and rats and in vivo and in vitro genotoxicity were examined.

The general toxicity profile, as seen with high doses of tamsulosin is consistent with the known pharmacological actions of the  $\alpha$ 1-adrenoceptors antagonists.

At very high dose levels the ECG was altered in dogs. This response is considered to be not clinically relevant. Tamsulosin showed no relevant genotoxic properties.

Increased incidences of proliferative changes of mammary glands of female rats and mice have been reported. These findings, which are probably mediated by hyperprolactinemia and only occurred at high dose levels, are regarded as irrelevant.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### ***Capsule fill***

Microcrystalline Cellulose (Grade 101)

Talc

Methacrylic Acid – Ethyl Acrylate Copolymer (1:1) Dispersion 30 percent (contains sodium lauril sulfate and polysorbate 80)

Triacetin

Hypromellose

Calcium stearate

#### ***Capsule shell***

Gelatin

FD and C Blue 2 (E132)

Iron Oxide Red (E172)

Iron Oxide Yellow (E172)

Titanium Dioxide (E171)

#### ***Printing ink***

Shellac (E904)

Propylene glycol (E1520)

Ammonia Solution Concentrated (E527)

Potassium hydroxide (E525)

Black iron oxide (E172)

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

36 months

## **6.4 Special precautions for storage**

This product does not require any special storage conditions.

## **6.5 Nature and contents of container**

Tamsulosin 0.4 mg capsules are available in OPA/ALU/PVC-Aluminium blister packs containing 10's, 20's, 30's, 50's, 60's, 90's, 100's, 200's hard capsules

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Ascend Laboratories (UK) Limited  
Elsley Court,  
20-22 Great Titchfield Street London,  
W1W 8BE  
United Kingdom

## **8 MARKETING AUTHORISATION NUMBER(S)**

PL 43805/0149

## **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

15/11/2023

**10 DATE OF REVISION OF THE TEXT**